

The Department of Vermont Health Access Medical Policy

Subject: Apnea Monitor

Last Review: June 6, 2016

Revision 6: August 26, 2015

Revision 5: January 2, 2015

Revision 4: November 1, 2013

Revision 3: October 25, 2011

Revision 2: March 29, 2010

Revision 1: March 31, 2009

Original Effective: September 10, 2008

Description of Service or Procedure

An apnea monitor is a device which monitors abnormal cessation of breathing and abnormal cardiac status for high risk children. It contains an alarm function which is triggered when the measured rate (respiratory rate, slow heart rate, and/or fast heart rate) differs from the pre-determined respiratory or cardiac parameters set by the treating provider. It is important that parents complete cardiopulmonary resuscitation (CPR) and apnea response training.

Disclaimer

Coverage is limited to that outlined in Medicaid Rule that pertains to the beneficiary's aid category. Prior Authorization (PA) is only valid if the beneficiary is eligible for the applicable item or service on the date of service.

Medicaid Rule

[7102.2](#) Prior Authorization Determination

[7103](#) Medical Necessity

[7505.2](#) Covered Services: "Items of durable medical equipment that have been pre-approved for coverage are limited to: apnea monitors..."

Medicaid Rules can be found at <http://humanservices.vermont.gov/on-line-rules>



Coverage Position

An apnea monitor may be covered for beneficiaries:

- When this device is prescribed by a licensed medical provider, enrolled in the Vermont Medicaid program, operating within their scope of practice in accordance with Vermont State Practice Act, who is knowledgeable in the use of apnea monitors and who provides medical care to the beneficiary AND
- When the clinical guidelines below are met.

Coverage Criteria

An apnea monitor may be covered for beneficiaries (infants under age 1 year) who have one of the diagnoses/high risk conditions listed below:

EXAMPLES OF DIAGNOSES:

- Infants who experienced some combination of the following Apparent Life Threatening Episode(s) (ALTE):
 - apnea (central or occasional obstructive),
 - choking or gagging,
 - skin color change (usually cyanotic or pallid but occasionally erythematous or plethoric),
 - marked changes in muscle tone.
 - Continued use is considered medically necessary until the infant remain event-free for six weeks.
- Infants with tracheotomies that make them vulnerable to airway compromise.
- Infants with anatomical abnormalities that make them vulnerable to airway compromise.
- Infants with metabolic disorder affecting respiratory control.
- Infants with neurologic disorder affecting respiratory control.
- Infants with chronic lung disease (i.e. bronchopulmonary dysplasia) especially those requiring mechanical ventilation, positive airway pressure, or supplemental oxygen.
- Apnea unresponsive to treatment.
- Apnea of prematurity, defined as sudden cessation of breathing that lasts for at least 20 seconds or is accompanied by bradycardia or hypoxemia, or cyanosis in infants younger than 37 weeks' gestational age. Continued use is considered medically necessary until infants are past post-conception age of 43 weeks and are event-free for six weeks.
- Chronic respiratory failure on home ventilator support.
- Infants requiring home ventilation for potential airway obstruction.
- Preterm infant with bradycardia and/or desaturation.
- Infants diagnosed with pertussis with positive cultures, upon discharge from acute care facility. If monitored for pertussis, use of an apnea monitor is considered medically necessary for **up to one-month** post-diagnosis.
- Infants with gastroesophageal reflux disease that results in apnea, bradycardia or oxygen desaturation, until the infants remain event-free for six weeks.
- Apnea accompanied by marked hypotonia; use of an apnea monitor until the infants remain event-free for six weeks.
- Infants discharged home on a schedule of weaning narcotics.
- Infants with bradycardia on caffeine, theophylline, or similar agents, until event free for 2 weeks off medication

SPECIAL NOTATIONS:

1. Coverage should be discontinued when clinical evaluation shows that the condition(s) requiring a monitor have been resolved or stabilized as indicated by:
 - a. The beneficiary has been free of events requiring stimulation or resuscitation as defined above. OR
 - b. The beneficiary has experienced significant stressors such as respiratory illness or immunizations without apnea.
2. Prior authorization is required for age one year and older.
3. Home cardiorespiratory monitoring should not be prescribed to prevent sudden infant death syndrome (SIDS).
4. The monitor should be equipped with an event recorder.
5. The physician should establish a specific plan for periodic review and termination of the home monitor before initiating therapy

Clinical guidelines for repeat service or procedure

If the beneficiary is still under one year of age and requires an apnea monitor for an additional time, the same criteria apply as for the initial approval.

Type of service or procedure covered

Apnea monitor and related supplies and services.

Type of service or procedure not covered (this list may not be all inclusive)

Apnea monitor not covered for:

- Sibling of SIDS
- Diagnosis of gastroesophageal reflux

References

American Academy of Pediatrics (AAP) (2003). Committee on Fetus and Newborn. Apnea, sudden infant death syndrome, and home monitoring. *Pediatrics*, 111(4):914-7. Retrieved May 26, 2016, from: <http://pediatrics.aappublications.org/content/111/4/914.full.pdf+html>

American Academy of Pediatrics (AAP) (2008). Committee on Fetus and Newborn. Hospital discharge of the high risk neonate-proposed guidelines. *Pediatrics*, 122(5):1119-1126. Retrieved May 25, 2016, from: <http://pediatrics.aappublications.org/content/122/5/1119.full.pdf+html>

American Academy of Pediatrics (AAP) (2005). Task Force on Sudden Infant Death Syndrome. The changing concept of sudden infant death syndrome: Diagnostic coding shifts, controversies regarding the sleeping environment, and new variables to consider in reducing risk. *Pediatrics*, 116(5): 1245-1255. Retrieved May 25, 2016, from: <http://pediatrics.aappublications.org/content/116/5/1245.full.pdf+html>

Minnesota Neonatal Physicians. (2016). Apnea and bradycardia in infants <44 weeks post menstrual age. Retrieved May 26, 2016, from: http://mnneo.org/getattachment/For-Providers/Guidelines/Apnea_and_Bradycardia_in_Infants.pdf

Di Fiore, J.M., Arko, M.K., Miller M.J., Krauss, A., Betkerur, A, Zadell, A. et al. (2001). Cardiorespiratory events in preterm infants referred for apnea monitoring studies. *Pediatrics*, 108(6):1304-8. Retrieved May 25, 2016, from: <http://pediatrics.aappublications.org/cgi/reprint/108/6/1304>

Di Fiore, J.M., Arko, M.K., Whitehouse, M., Kimball, A., & Martin, R.J. (2005). Apnea is not prolonged by acid gastroesophageal reflux in preterm infants. *Pediatrics*, 116(5):1059-1063. Retrieved May25, 2016, from: <http://pediatrics.aappublications.org/content/116/5/1059.full.pdf+html>

Eichenwald, E., Aina, A., & Stark, A. (1997). Apnea frequency persists beyond term gestation in infants delivered at 24 to 28 weeks. *Pediatrics*, 100(3). Retrieved May25, 2016, from: <http://pediatrics.aappublications.org/content/100/3/354.full.pdf+html>

Ramanathan, R., Corwin, M., Hunt, C., Lister, G., Tinsley, L., Baird, T. et al. (2001). Cardiorespiratory events recorded on home monitors. Comparison of healthy infants with those at increases risk for SIDS. *Journal of the American Medical Association* 285(17). Retrieved May 25, 2016, from: <http://jama.ama-assn.org/cgi/content/full/285/17/2199>

Pediatric Gastroesophageal Reflux Clinical Practice Guidelines. (2009). *Journal of Pediatric Gastroenterol Nutrition*, 49(2). Retrieved May 25, 2016, from: <http://journals.lww.com/jpgn/Abstract/2009/10000/Notices.22.aspx>

RT Board Review (2003). Apnea Monitoring. Retrieved May 4, 2015, from: http://rtboardreview.com/public/other_procedures/apnea_monitoring.htm

Silvesti, J.M., Lister, G., Corwin, M.J., Smok-Pearsall, S.H., Baird, T., Crowell, D.H., Cantey-Kiser, J., Hunt, C.E., et al. (2005). Factors that influence use of a home cardiorespiratory monitor for infants. *Archives of Pediatrics & Adolescent Medicine*. 159, pp. 18-24.

Tieder, J.S., Altman, R., Bonkowsky, J.L., Brand, D.A., Claudius, I., Cunningham, D.J. et al. (2013). Management of Apparent Life-Threatening Events in Infants: A Systematic Review. *Journal of Pediatrics*, 163(1). Retrieved May 25, 2016, from: [http://www.jpeds.com/article/S0022-3476\(12\)01575-2/fulltext](http://www.jpeds.com/article/S0022-3476(12)01575-2/fulltext)

This document has been classified as public information.