



Department of Vermont Health Access Pharmacy Benefit Management Program

Vermont Preferred Drug List and Drugs Requiring Prior Authorization (includes clinical criteria)

EFFECTIVE
Version
Updated:01/01/2025

The Commissioner for Office of Vermont Health Access shall establish a pharmacy best practices and cost control program designed to reduce the cost of providing prescription drugs, while maintaining high quality in prescription drug therapies. The program shall include:

"A preferred list of covered prescription drugs that identifies preferred choices within therapeutic classes for particular diseases and conditions, including generic alternatives."

From Act 127 passed in 2002

The following pages contain:

- The therapeutic classes of drugs subject to the Preferred Drug List, the drugs within those categories and the criteria required for Prior Authorization (P.A.) of non-preferred drugs in those categories. The therapeutic classes of drugs which have clinical criteria for Prior Authorization may or may not be subject to a preferred agent.
- Within both categories there may be drugs or drug classes that are subject to Quantity Limit Parameters.

Therapeutic class criteria are listed alphabetically. Within each category the Preferred Drugs are noted in the left-hand columns. Representative non-preferred agents have been included and are listed in the right-hand column. Any drug not listed as preferred in any of the included categories requires Prior Authorization. Approval of non-preferred brand name products may require trial and failure of at least 2 different generic manufacturers. Drug samples, prescription discount programs, and patient assistance programs are not considered adequate justification for stabilization on non-preferred drugs. Drugs used for weight loss, drugs used to promote fertility, and drugs used for cosmetic purposes or hair growth are excluded from coverage under the Vermont Medicaid Pharmacy program.

Optum (formally Change Healthcare) PRESCRIBER Call Center: PA Requests Tel: 1-844-679-5363; Fax: 1-844-679-5366 Note: Fax requests are responded to within 24 hrs.	Optum (formally Change Healthcare) PHARMACY Call Center: PA Requests Tel: 1-844-679-5362 Available for assistance with claims processing	Optum Assc Dir. Account Manager: Michael Ouellette, RPh E-Mail: michael_ouellette@optum.com
	DVHA Pharmacy Unit Staff: Tel: 802-241-0140 Fax: 802-241-0268 E-Mail: ahs.dvhaph@vermont.gov	

This is not an all-inclusive list of available covered drugs and includes only managed categories. Unless otherwise stated, the listing of a particular brand or generic name includes all dosage forms of that drug. NR indicates a new drug that has not yet been reviewed by the P&T Committee.

Drugs highlighted in yellow denote a change in PDL status.

To search the PDL, press CTRL + F

Contents

ACNE AGENTS	6
ADHD AND NARCOLEPSY CATAPLEXY MEDICATIONS	7
ALLERGEN IMMUNOTHERAPY	10
ALPHA1-PROTEINASE INHIBITORS	11
ALZHEIMER'S MEDICATIONS	11
ANALGESICS	13
ANKYLOSING SPONDYLITIS: INJECTABLES	18
ANTI-ANXIETY: ANXIOLYTICS	19
ANTICOAGULANTS	20
ANTICONVULSANTS	21
ANTIDEPRESSANTS	23
ANTI-DIABETICS	27
ANTI-EMETICS	31
ANTI-HYPERTENSIVES	33
ANTI-INFECTIVES ANTIBIOTICS	38
ANTI-INFECTIVES ANTIFUNGAL	43
ANTI-INFECTIVES ANTIMALARIALS	45
ANTI-PARASITICS	45
ANTI-INFECTIVES ANTI-VIRALS	46
MIGRAINE THERAPY: PREVENTATIVE TREATMENTS	48
MIGRAINE THERAPY: ACUTE TREATMENTS	50
ANTI-PSYCHOTIC ATYPICAL & COMBINATIONS (CHILDREN < 18 YEARS OLD)	52
ANTI-PSYCHOTIC ATYPICAL & COMBINATIONS (ADULTS > 18 YEARS OLD)	54
ANTI-PSYCHOTIC: TYPICALS	58
ANTIRETROVIRAL THERAPY HUMAN IMMUNODEFICIENCY VIRUS (HIV)	59
BILE SALTS AND BILIARY AGENTS	62
BONE RESORPTION INHIBITORS	64
BOTULINUM TOXINS	65
BPH AGENTS	66
BULK POWDERS	66

CARDIAC GLYCOSIDES	67
CLOSTRIDIUM DIFFICILE (C.diff) AGENTS	67
CUSHING’S DISEASE	68
GASTROINTESTINAL AGENTS: BOWEL PREP AGENTS, CONSTIPATION/DIARRHEA, IRRITABLE BOWEL SYNDROME-CONSTIPATION (IBS-C), IRRITABLE BOWEL SYNDROME-DIARRHEA (IBS-D), SHORT BOWEL SYNDROME, OPIOID INDUCED CONSTIPATION	69
CONTINUOUS GLUCOSE MONITORS	71
CONTRACEPTIVES	72
CORONARY VASODILATORS/ANTIANGINALS/SINUS NODE INHIBITORS	76
CORTICOSTEROIDS: ORAL	77
COUGH AND COLD PREPARATIONS	77
CYSTIC FIBROSIS MEDICATIONS	78
DERMATOLOGICAL AGENTS	79
DESMOPRESSIN: INTRANASAL/ORAL	85
DIABETIC TESTING SUPPLIES	86
ENDOMETRIOSIS/UTERINE FIBROIDS AGENTS	86
EPINEPHRINE: SELF-ADMINISTERED	87
ESTROGENS: VAGINAL	87
GASTROINTESTINAL	88
GAUCHER’S DISEASE MEDICATIONS	92
GOUT AGENTS	93
GROWTH STIMULATING AGENTS	93
hATTR TREATMENTS	95
HEMATOPOIETICS	97
HEMOPHILIA TREATMENTS	98
HEPATITIS C AGENTS	102
HEREDITARY ANGIOEDEMA MEDICATIONS	103
HIDRADENITIS SUPPURATIVA	104
HYPERKALEMIA AGENTS	104
HYPOTHYROID AGENTS	105
IDIOPATHIC PULMONARY FIBROSIS (IPF)	105
IMMUNOLOGIC THERAPIES FOR ASTHMA	106

IMMUNOSUPPRESSANTS, ORAL	110
CRYOPYRIN ASSOCIATED PERIODIC SYNDROMES (CAPS) AND PERIODIC FEVER SYNDROME (PFS)	111
IRON CHELATING AGENTS	112
LIPOTROPICS	112
MISCELLANEOUS	115
MOOD STABILIZERS	126
MOVEMENT DISORDERS	127
MULTIPLE SCLEROSIS MEDICATIONS	127
MUSCLE RELAXANTS, SKELETAL	129
MUSCULAR DYSTROPHY AGENTS	130
NEUROGENIC ORTHOSTATIC HYPOTENSION	132
NEUROPATHIC PAIN & FIBROMYALGIA AGENTS	132
NUTRITIONALS, LIQUID ORAL SUPPLEMENTS	133
ONCOLOGY: DRUGS (select)	134
OPHTHALMICS	134
OTIC ANTI-INFECTIVES/ANTI-INFLAMMATORIES	140
OVER THE COUNTER (OTC) MEDICATIONS	140
PANCREATIC ENZYME PRODUCTS	140
PARATHYROID AGENTS	140
PARKINSON'S MEDICATIONS	141
PLATELET INHIBITORS	143
PLATELET STIMULATING AGENTS	143
PSEUDOBULBAR AFFECT AGENTS	144
PSORIASIS	145
PULMONARY AGENTS	146
PULMONARY ARTERIAL HYPERTENSION MEDICATIONS	153
RENAL DISEASE: PHOSPHATE BINDERS	155
RESTLESS LEG SYNDROME MEDICATIONS	155
RHEUMATOID, JUVENILE & PSORIATIC ARTHRITIS: IMMUNOMODULATORS	156
SICKLE CELL DISEASE THERAPIES	157
SEDATIVE/HYPNOTICS	158

SMOKING CESSATION THERAPIES	159
SUBSTANCE USE DISORDER TREATMENTS	160
TESTOSTERONE REPLACEMENT THERAPY	161
URINARY ANTISPASMODICS	163
VAGINAL ANTI-INFECTIVES	163
VASOPRESSIN RECEPTOR ANTAGONIST	164
VITAMINS: PRENATAL MULTIVITAMINS	164

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

ACNE AGENTS

ORAL AGENTS		
<p>AMNESTEEM (isotretinoin) capsules CLARAVIS (isotretinoin) capsules ZENATANE (isotretinoin) capsules</p>	<p>Absorica® (isotretinoin) capsules Isotretinoin capsules</p>	<p>Absorica, Isotretinoin: patient has had a documented side effect, allergy, or treatment failure with at least two isotretinoin preferred products.</p>
TOPICAL AGENTS		
<p><u>BENZOYL PEROXIDE PRODUCTS</u> BENZOYL PEROXIDE 2.5%, 5%, 10%G; 3%, 5%, 10% CL; 5.3%, 9.8% F</p> <p><u>CLINDAMYCIN PRODUCTS</u> CLINDAMYCIN 1% S, G, L, P (compare to Cleocin-T)</p> <p><u>ERYTHROMYCIN PRODUCTS</u> ERYTHROMYCIN 2% S, G</p> <p><u>SODIUM SULFACETAMIDE PRODUCTS</u> KLARON® (sodium sulfacetamide 10% L)</p> <p><u>COMBINATION PRODUCTS</u> ERYTHROMYCIN / BENZOYL PEROXIDE CLINDAMYCIN/BENZOYL PEROXIDE (compare to Benzaclin®) G</p> <p><u>OTHER</u> C=cream, CL=cleanser, E=emulsion, F=foam, G=gel, L=lotion, O=ointment, P=pads, S=solution, W=wash, B=bar</p>	<p>Benzol Peroxide 5%, 10%L</p> <p>Clindacin (clindamycin) 1% CL, P, Swab Clindamycin 1%F Clindamycin 1%G (compare to Clindagel) 75mL bottle Cleocin-T® (clindamycin) 1% L</p> <p>Erygel® (erythromycin 2% Ery (erythromycin 2%) P</p> <p>Sodium Sulfacetamide 10% L Sodium Sulfacetamide/Sulfur CL, C, P, E Sodium Sulfacetamide/Sulfur W Sumaxin® (sulfacetamide/sulfur L, P, W)</p> <p>Benzaclin® (clindamycin/benzoyl peroxide) Benzamycin® (erythromycin/benzoyl peroxide) Cabtreo® (clindamycin phosphate/benzoyl peroxide/adapalene) Clindamycin/Benzoyl Peroxide Pump Onexton® (clindamycin/benzoyl peroxide)</p> <p>Dapsone 5%, 7.5% G</p> <p>All other brands any topical acne anti-infective medication</p>	<p>Single ingredient products: patient has had a documented side effect, allergy, or treatment failure with two preferred products including one from the same sub-category, if there is one available. If a product has an AB rated generic, there must have been a trial of the generic.</p> <p>Benzaclin, Benzamycin: patient must have a documented intolerance to the generic equivalent.</p> <p>Sodium Sulfacetamide Products: patient has had a documented side effect, allergy, or treatment failure with two preferred products, one of which must be Klaron lotion.</p> <p>Clindamycin/Benzoyl peroxide pump, Onexton: there must be a clinically compelling reason why clindamycin/benzoyl peroxide gel cannot be used.</p> <p>Cabtreo: Patient must try and have failed individual preferred products.</p> <p>Limitations: Kits with non-drug products are not covered</p>
TOPICAL – ANDROGEN RECEPTOR INHIBITORS		
<p>All products require PA</p>	<p>Winlevi® (clascoterone) 1% C</p>	<p>Winlevi: patient has had a documented side effect, allergy, or treatment failure with two preferred topical acne agents.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
TOPICAL - RETINOIDS		
<p>AVITA® (tretinoin) ADAPALENE 0.1% G, 0.3% G DIFFERIN® (adapalene) 0.1% G RETIN-A® (tretinoin) 0.025%, 0.05%, 0.1% C; 0.01%, 0.025% G</p> <p><i>C= cream, G=gel, L=lotion</i></p>	<p>Adapalene (compare to Differin®) 0.1% C Adapalene/Benzoyl Peroxide 0.1-2.5% G Altreno™ (tretinoin) 0.05% L Arazlo® (tazarotene) 0.045% L Atralin® (tretinoin) 0.05% G Clindamycin/tretinoin 1.2-0.025% G Fabiator® (tazarotene) 0.1% F Plixda® (adapalene) 0.1% swabs Retin-A Micro® (tretinoin microsphere) 0.04%, 0.06%, 0.08%, 0.1% G Tazarotene (compare to Tazorac®) 0.1% C Tretinoin (compare to Retin-A®) 0.025%, 0.05%, 0.1% C; 0.01%, 0.025% G Tretinoin microsphere (compare to Retin-A Micro®) 0.1%, 0.04% Twynéo® (tretinoin/benzoyl peroxide) 0.1%-3% C</p>	<p>Altreno, Atralin, Retin-A Micro, Tretinoin, Tretinoin microsphere: diagnosis or indication is acne vulgaris, actinic keratosis, or rosacea AND patient has had a documented side effect, allergy, or treatment failure with a preferred topical tretinoin product (Avita or Retin-A®).</p> <p>Adapalene Cream: patient has had a documented side effect, allergy, or treatment failure with adapalene gel.</p> <p>Arazlo, Fabiatar, Tazarotene: patient has had a documented side effect or treatment failure with a preferred topical tretinoin product and adapalene.</p> <p>Adapalene/benzoyl peroxide gel, Clindamycin/tretinoin gel, Twynéo: patient has had a documented side effect or treatment failure on combination therapy with the separate ingredients of the combination product</p> <p>Plixda: patient has had a documented side effect, allergy, or treatment failure with brand Differin AND a generic adapalene product.</p> <p>Limitations: Coverage of topical retinoid products will not be approved for cosmetic use (wrinkles age spots, etc.) (i.e. Avage, Renova, Tri-Luma).</p>
TOPICAL - ROSACEA		
<p>AZELAIC ACID 15% G FINACEA 15% F METRONIDAZOLE 0.75% C, G, L</p> <p><i>C=cream, F=Foam, G=gel, L=lotion</i></p>	<p>All brand metronidazole products (MetroCream® 0.75% C, MetroGel® 1% G, MetroLotion® 0.75% L, Noritate® 1% C etc.) Ivermectin (compare to Soolanta®) 1% C Metronidazole 1% G Rhofade® (oxymetazoline) 1% C</p>	<p>Brand name metronidazole products, Metronidazole 1% gel (generic): diagnosis or indication is rosacea AND patient has had a documented side effect, allergy, or treatment failure with a preferred generic topical metronidazole product. If a product has an AB rated generic, there must have also been a trial of the generic formulation.</p> <p>Ivermectin, Rhofade: the patient has had a documented side effect, allergy, or treatment failure with 2 preferred topical rosacea agents.</p> <p>Limitations: The use of Mirvaso (brimonidine topical gel) for treating skin redness is considered cosmetic. Medications used for cosmetic purposes are excluded from coverage. Mirvaso topical gel has not been shown to improve any other symptom of rosacea (e.g. pustules, papules, flushing, etc.) or to alter the course of the disease.</p>
ADHD AND NARCOLEPSY CATAPLEXY MEDICATIONS		
SHORT/INTERMEDIATE ACTING STIMULANTS		
<p>AMPHETAMINE/DETRIOAMPHETAMINE (compare to Adderall®)</p>	<p>Adderall® (amphetamine/dextroamphetamine) Amphetamine Sulfate (compare to Evekeo)</p>	<p>Clinical Criteria for ALL non-preferred drugs: patient has a diagnosis of ADD, ADHD or narcolepsy AND patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>DEXMETHYLPHENIDATE (compare to Focalin®) METHYLPHENIDATE (compare to Ritalin®) tablets, solution METHYLPHENIDATE SR (compare to Ritalin® SR) PROCENTRA® (dextroamphetamine sulfate) 1 mg/ml oral solution</p>	<p>Desoxyn® (methamphetamine) Dextroamphetamine sulfate 1 mg/ml oral solution Dextroamphetamine IR (Zenedi 5 or 10 mg, formerly Dexedrine®) Evekeo® (amphetamine sulfate) Evekeo® ODT (amphetamine sulfate) Focalin® (dexmethylphenidate) Methamphetamine (compare to Desoxyn®) Methylphenidate (compare to Ritalin®) chewable tablets Methylin® (compare to Ritalin®) solution Ritalin® (methylphenidate) Zenedi® (dextroamphetamine IR) 2.5 mg, 7.5 mg, 15 mg, 20 mg, 30 mg tablets</p>	<p>justification for stabilization.) OR patient meets additional clinical criteria outlined below. Adderall, Focalin, Methylin, Ritalin: the patient must have had a documented intolerance to the preferred generic equivalent. Methamphetamine and Desoxyn: Given the high abuse potential of methamphetamine and Desoxyn, the patient must have a diagnosis of ADD, ADHD or narcolepsy and have failed all preferred treatment alternatives. In addition, for approval of brand name Desoxyn, the patient must have had a documented intolerance to generic methamphetamine. Methylphenidate chewable tablets: patient has a documented intolerance to methylphenidate and Methylin solution. Evekeo ODT, Dextroamphetamine oral solution: patient has a medical necessity for a non-solid oral dosage form. (e.g. swallowing disorder). AND the patient has a documented intolerance Procentra oral solution. Amphetamine Sulfate, Dextroamphetamine IR, Zenedi, Evekeo: the patient has had a documented side-effect, allergy, or treatment failure of at least 2 preferred agents (If a product has an AB rated generic, there must have been a trial of the generic.)</p>
LONG ACTING STIMULANTS		
<p><u>METHYLPHENIDATE PRODUCTS</u> <u>ORAL</u> CONCERTA® (methylphenidate SA OSM IR/ER, 22:78%) DEXMETHYLPHENIDATE SR 24 HR IR/ER, 50:50% (compare to Focalin XR®) FOCALIN® XR (dexmethylphenidate SR 24 HR) METHYLPHENIDATE CR, IR/ER, 30:70% (compare to Metadate CD®) METHYLPHENIDATE SR 24 HR, IR/ER, 50:50% (compare to Ritalin LA®) QUILLICHEW ER™ (methylphenidate IR/ER, 30:70%) chewable tablets RITALIN LA® (methylphenidate SR 24 HR, IR/ER, 50:50%) <u>ORAL SUSPENSION</u> QUILLIVANT XR® (methylphenidate IR/ER, 20:80%) <i>QTY LIMIT:</i> 1 bottle/Rx (60ml, 120ml, 150ml) 2 bottles/Rx (180ml)</p>	<p>Adhansia® XR (methylphenidate IR/ER 20:80%) <i>QTY LIMIT:</i> 1 capsule/day Aptensio® XR (methylphenidate DR 24HR IR/ER, 40:60%) Azstarys™ (serdexmethylphenidate/ dexmethylphenidate) Cotempla® XR (methylphenidate IR/ER 25:75%) ODT Jornay PM™ (methylphenidate ER) capsules <i>QTY LIMIT:</i> 1 capsule/day Methylphenidate DR 24HR IR/ER, 40:60% (compare to Aptensio®XR) Methylphenidate SA OSM IR/ER, 22:78% (compare to Concerta®) Relexxii® (methylphenidate ER OSM) IR/ER, 22:78%</p>	<p>Clinical criterial for ALL non-preferred drugs: the patient has a diagnosis of ADD, ADHD or narcolepsy AND has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization) OR meets the additional clinical criteria outlined below. Azstarys, Adhasia XR, Cotempla XR ODT, Jornay PM: patient has had a documented side-effect, allergy, or treatment failure on 3 preferred long-acting Methylphenidate products. Aptensio XR, Methylphenidate DR 40:60: patient has had a documented side effect, allergy, or treatment failure on two preferred long-acting Methylphenidate products. For approval of Methylphenidate DR 40:60, the patient must also have a documented intolerance to brand Aptensio XR. Methylphenidate SA OSM: the patient must have a documented intolerance to brand Concerta. Relexxi: Both Concerta and methylphenidate SA OSM must be on a long-term backorder and unavailable from the manufacturer.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>TRANSDERMAL</u> All products require PA</p> <p><u>AMPHETAMINE PRODUCTS</u> <u>ORAL</u> AMPHETAMINE/DEXTROAMPHETAMINE SR 24 HR, IR/ER, 50:50% (compare to Adderall XR®) DEXTROAMPHETAMINE 24 HR SR (compare to Dexedrine CR®) VYVANSE® (lisdexamfetamine) capsule <i>QTY LIMIT: 1 cap /day</i></p> <p><u>TRANSDERMAL</u> All products require PA</p>	<p>Daytrana® (methylphenidate patch) <i>QTY LIMIT: 1 patch/day</i> Methylphenidate patch (compare to Daytrana®) <i>QTY LIMIT: 1 patch/day</i></p> <p>Adderall XR® (amphetamine/dextroamphetamine SR 24 HR, IR/ER, 50:50%) Adzenys XR® ODT (amphetamine SR 24 HR, IR/ER, 50:50%) <i>QTY LIMIT: 1 cap/day</i> Adzenys ER™ suspension (amphetamine SR 24 HR, IR/ER, 50:50%) Dyanavel™ suspension (amphetamine/dextroamphetamine SR) <i>QTY LIMIT: 240ml/30days</i> Dyanavel® XR (amphetamine/dextroamphetamine SR) chewable tablet Dexedrine CR® (dextroamphetamine 24 HR SR) Lisdexamfetamine (compare to Vyvanse®) capsule <i>QTY LIMIT: 1 cap/day</i> Mydayis® (mixed amphetamine salts) extended-release capsules Vyvanse® (lisdexamfetamine) chewable tablet <i>QTY LIMIT: 1 tab/day</i></p> <p>Xelstry™ (dextroamphetamine patch) <i>QTY LIMIT: 1 patch/day</i></p>	<p>Daytrana patch, Methylphenidate patch: patient has a documented medical necessity for a specialty non-oral dosage form AND for approval of generic Methylphenidate patch, the patient must have a documented intolerance to brand Daytrana.</p> <p>Adderall XR: Patient must have a documented intolerance to generic product. Adzenys XR ODT, Adzenys ER suspension, Dyanavel XR chewable tablet, Vyvanse Chew: Patient must have a documented side effect, allergy, or treatment failure to Dyanavel XR suspension. Dexedrine CR, Mydayis: patient must have a documented intolerance to two preferred amphetamine products. For approval of brand Dexedrine CR, the patient must also have a documented intolerance to the generic equivalent. Dyanavel XR Suspension: patient must have medical necessity for a non-solid oral dosage form. Lisdexamfetamine: patient must have a documented intolerance to Brand Vyvanse. Xelstry™: patient has a documented medical necessity for a specialty non-oral dosage form.</p>
MISCELLANEOUS		
<p>ARMODAFINIL (compare to Nuvigil®) <i>QTY LIMIT: 50 mg = 2 tabs/day</i> 150 mg/200 mg/250 mg = 1 tab/day, Max days supply = 30 days ATOMOXETINE (compare to Strattera®) <i>QTY LIMIT: 10, 18, 25 and 40 mg = 2 capsules/day</i> 60, 80 and 100 mg = 1 capsule/day FDA maximum recommended dose = 100 mg/day CLONIDINE ER <i>QTY LIMIT: 4 tabs/day</i> GUANFACINE ER (Intuniv®) MODAFINIL (compare to Provigil®) <i>QTY LIMIT: 100 mg = 1.5 tablets/day; 200 mg = 2 tablets/day</i></p>	<p>Intuniv® (guanfacine extended release) tablet <i>QTY LIMIT: 1 tablet/day</i> Nuvigil® (armodafinil) <i>QTY LIMIT: 50 mg = 2 tablets/day; 150 mg/200 mg/250 mg = 1 tablet/day, Max days supply = 30 days</i> Provigil® (modafinil) <i>QTY LIMIT: 100 mg = 1.5 tablets/day; 200 mg = 2 tablets/day</i> Maximum Daily Dose = 400 mg, Max day supply = 30 days</p>	<p>Intuniv, Nuvigil, Provigil, Strattera: patient must have a documented intolerance to the generic equivalent. Qelbree: The patient has had a documented side effect, allergy, contraindication, or treatment failure to two preferred stimulant OR atomoxetine Sunosi: patient has had a documented side effect, allergy, or treatment failure to 2 preferred agents (may be stimulant or non-stimulant) Wakix patient has no known risk factors for increased QT prolongation (e.g. cardiac arrhythmias, symptomatic bradycardia, hypokalemia, or congenital prolongation of the QT interval) AND medication is not being used in combination with other drugs known to prolong the QT interval (e.g.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>Maximum Daily Dose = 400 mg, Max day supply = 30 days</p> <p>Preferred After Clinical Criteria Are Met QELBREE® (viloxazine hydrochloride) ER capsule <i>QTY LIMIT:</i> 100 mg = 1 capsule/day 150 mg = 2 capsules/day 200 mg = 3 capsules/FDA maximum recommended dose = 600 mg/day</p>	<p>FDA maximum recommended dose = 600 mg/day</p> <p>Strattera® (atomoxetine) <i>QTY LIMIT:</i> 10, 18, 25 and 40 mg = 2 capsules/day 60, 80 and 100 mg = 1 capsule/day</p> <p>FDA maximum recommended dose = 100 mg/day</p> <p>Sunosi® (solriamfetol) tablet <i>QTY LIMIT:</i> 1 tablet/day</p> <p>FDA maximum recommended dose = 150 mg/day</p> <p>Wakix® (pitolisant) tablet <i>QTY LIMIT:</i> 2 tablets/day</p> <p>FDA maximum recommended dose = 35.6 mg/day</p> <p>Xyrem® (sodium oxybate) oral solution <i>QTY LIMIT:</i> 540 ml/30 days</p> <p>Xywav™ (calcium, magnesium, potassium, and sodium oxybates) solution <i>QTY LIMIT:</i> 9 g (18 mL)/day</p>	<p>antipsychotics, erythromycin, tricyclic antidepressants) AND patient has had a documented side effect, allergy, or treatment failure to at least 3 agents (may be preferred or non-preferred; may be stimulant or non-stimulant), one of which must be Sunosi.</p> <p>Xyrem, Xywav: patient has had a documented side effect, allergy, or treatment failure to 2 preferred agents (may be stimulant or non-stimulant) and Sunosi AND patient has been enrolled in the REMS program AND for approval of Xywav, the patient must have a documented intolerance to Xyrem.</p>

ALLERGEN IMMUNOTHERAPY

<p>All products require PA</p>	<p>Grastek® (Timothy Grass Pollen Extract) <i>QTY LIMIT:</i> 1 tablet/day</p> <p>Odactra® (House Dust Mite Allergen Extract) <i>QTY LIMIT:</i> 1 tablet/day</p> <p>Oralair® (Sweet Vernal/Orchard/Perennial Rye/Timothy/Kentucky Blue Grass Mixed Pollen Allergen Extract) <i>QTY LIMIT:</i> 1 tablet/day</p> <p>Palforzia® (peanut allergen powder-dnfp)</p> <p>Ragwitek® (Short Ragweed Pollen Allergen Extract) <i>QTY LIMIT:</i> 1 tablet/day</p>	<p>Grastek, Oralair, Ragwitek:</p> <ul style="list-style-type: none"> • The patient's age is FDA approved for the given indication AND • Treatment must start 12 weeks before expected onset of pollen season and only after confirmed by positive skin test or in vitro testing for specific IgE antibodies to the relevant allergen AND • Patient must have an auto-injectable epinephrine on hand. <p>Odactra:</p> <ul style="list-style-type: none"> • The patient's age is FDA approved for the given indication AND • Diagnosis is confirmed by positive skin test or in vitro testing for IgE antibodies to <i>Dermatophagoides farinae</i> or <i>Dermatophagoides pteronyssinus</i> house dust mites AND • Patient must have an auto-injectable epinephrine on-hand <p>Palforzia:</p> <ul style="list-style-type: none"> • Patient age ≥ 4 years and ≤ 17 years for initial dose escalation or ≥ 4 years for up-dosing and maintenance • The prescriber is an allergist or immunologist • Prescriber must provide the testing to show that the patient is allergic to peanuts • Patient must not have a recent history of uncontrolled asthma, eosinophilic esophagitis, or other eosinophilic GI disease. • Prescriber, pharmacy, and patient must be registered with the REMS
--------------------------------	---	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

		<ul style="list-style-type: none"> program • Patient must have an auto-injectable epinephrine on-hand • Initial approval will be granted for 6 months and includes approval for initial dose escalation and Up Dosing. Approval for Up Dosing may be extended if the patient was unable to tolerate all the dose levels at 2-week intervals. • For approval of Maintenance Dosing (300mg daily), pharmacy records will be evaluated to assess compliance with once daily therapy and ensure no level was missed during Up Dosing. Documentation must be provided attesting that the patient has not experienced any treatment restricting adverse events (e.g. systemic allergic reactions, severe anaphylaxis).
--	--	--

ALPHA1-PROTEINASE INHIBITORS

<p>All products require PA</p>	<p>Aralast NP[®] Glassia[®] Prolastin-C[®] Zemaira[®] **Maximum days supply per fill for all drugs is 14 days**</p>	<p>Criteria for Approval: The indication for use is treatment of alpha1 - proteinase inhibitor deficiency-associated lung disease when all of the following criteria are met: Patient's alpha1 -antitrypsin (ATT) concentration < 80 mg per dl [or < 11 micromolar] AND patient has obstructive lung disease as defined by a forced expiratory volume in one second (FEV1) OF 30 - 65% of predicted or a rapid decline in lung function defined as a change in FEV1 of > 120 mL/year. AND medication is being administered intravenously (inhalation administration will not be approved) AND patient is a non-smoker OR patient meets above criteria except lung function has deteriorated beneath above limits while on therapy.</p>
--------------------------------	---	--

ALZHEIMER'S MEDICATIONS

CHOLINESTERASE INHIBITORS

<p>DONEPEZIL (compare to Aricept[®]) tablet 5 mg and 10 mg <i>QTY LIMIT:</i> 1 tablet/day DONEPEZIL ODT (compare to Aricept[®] ODT) <i>QTY LIMIT:</i> 1 tablet/day GALANTAMINE tablet RIVASTIGMINE (compare to Exelon[®]) capsule <i>QTY LIMIT:</i> 2 capsules/day</p> <p><u>SOLUTION</u> All products require PA</p>	<p>Aricept[®] (donepezil) Tablet <i>QTY LIMIT:</i> 1 tablet/day Donepezil (compare to Aricept[®]) Tablet 23 mg Galantamine ER capsule (compare to Razadyne[®] ER) Razadyne ER[®] (galantamine) capsule</p> <p>Galantamine (compare to Razadyne[®]) Oral Solution</p>	<p>Donepezil 23mg Tablet, Galantamine ER Capsule, Razadyne ER Capsule: the patient has been started and stabilized on the requested medication (Note: samples are not considered adequate justification for stabilization) OR patient had a documented side effect, allergy, or treatment failure to a preferred cholinesterase inhibitor.</p> <p>Adlarity: medical necessity for a specialty dosage form has been provided AND the patient had a documented side effect, allergy, or treatment failure to Exelon patch.</p> <p>Aricept: the patient has a documented intolerance to the generic product.</p> <p>Galantamine Oral Solution, Rivastigmine patch: medical necessity for a specialty dosage form has been provided. AND for approval of rivastigmine patch the patient has a documented intolerance to brand Exelon patch.</p>
--	---	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<u>TRANSDERMAL</u>		
EXELON® (rivastigmine transdermal) Patch <i>QTY LIMIT</i> : 1 patch/day	Adlarity® (donzpezil) patch <i>QTY LIMIT</i> : 12 patches/84 days Rivastigmine (compare to Exelon®) patch <i>QTY LIMIT</i> : 1 patch/day	
IMMUNOGLOBULIN GAMMA 1 (IgG1) MONOCLONAL ANTIBODY		
All products require PA	Leqembi® (lecanemab-irmb) IV solution	Leqembi: <ul style="list-style-type: none"> • Patient is 50 years of age or older • Prescriber has assessed and documented baseline disease severity utilizing an objective measure/tool (e.g., MMSE, Alzheimer's Disease Assessment Scale-Cognitive Subscale [ADAS-Cog-13], Alzheimer's Disease Cooperative Study-Activities of Daily Living Inventory-Mild Cognitive, Impairment version [ADCS-ADL-MCI], Clinical Dementia Rating-Sum of Boxes [CDR-SB]). • Patient has mild cognitive impairment (MCI) due to Alzheimer's Disease or mild Alzheimer's dementia as evidenced by the following: <ul style="list-style-type: none"> ○ Clinical Dementia Rating (CDR) Global Score of 0.5 ○ Objective evidence of cognitive impairment at screening ○ MMSE score between 22 and 30 ○ PET scan is positive for amyloid beta plaque OR Cerebrospinal fluid (CSF) test is positive for amyloid • Patient has had a recent (within 1 year) brain MRI prior to initiating treatment and prescriber attests to a repeat brain MRI as directed in the labeling (to the 5th, 7th, and 14th infusion for Leqembi). • Patient does not have any of the following within 1 year of treatment initiation: pretreatment localized superficial siderosis, 10 or more brain microhemorrhages, or brain hemorrhage >1 cm • For re-approval, the patient must have responded to therapy compared to pre-treatment baseline as evidenced by improvement, stabilization, or slowing in cognitive or functional impairment AND patient has not progressed to moderate or severe disease (there is insufficient evidence in moderate or severe AD).
NMDA RECEPTOR ANTAGONIST		
MEMANTINE Tablets MEMANTINE XR Oral capsule <i>QTY LIMIT</i> : 1 capsule/day	Memantine oral solution	Namenda: Patient has a documented intolerance to the generic.
	Namenda® (memantine) tablet	Memantine Oral Solution: medical necessity for a specialty dosage form has been provided.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

CHOLINESTERASE INHIBITOR/NMDA COMBINATION

All products require PA	Namzatic [®] (donepezil/memantine) Capsule <i>QTY LIMIT</i> : 1 capsule/day	Namzatic : Clinically compelling reason why the individual ingredients of donepezil and memantine cannot be used.
-------------------------	---	--

ANALGESICS

MISCELLANEOUS: TOPICAL AND TRANSDERMAL PATCH

LIDOCAINE 3% Cream LIDOCAINE 4% OTC Patch LIDOCAINE 4% cream LIDOCAINE 5% Ointment LIDOCAINE 5% patch <i>QTY LIMIT</i> : 3 patches/day LIDOCAINE/PRILOCAINE 2.5-2.5% Cream	Qutenza [®] Patch (capsaicin 8 %) <i>QTY LIMIT</i> : 4 patches/90 days Ztlido [™] Patch (lidocaine 1.8%) <i>QTY LIMIT</i> : 3 patches/day (Note: Please refer to Analgesics: COX IIs and NSAIDs for topical NSAIDS)	Qutenza, Ztlido : diagnosis or indication is post-herpetic neuralgia AND patient has had a documented side effect, allergy, treatment failure or contraindication to 2 drugs in the tricyclic antidepressant (TCA) class and/or anticonvulsant class as well as Lidocaine 5% patch. OR patient has a medical necessity for transdermal formulation (ex. dysphagia, inability to take oral medications) AND patient has had a documented side effect, allergy, treatment failure or contraindication to lidocaine 5% patch.
--	---	---

OPIOIDS: SHORT ACTING

ACETAMINOPHEN W/CODEINE (compare to Tylenol [®] w/codeine) (age >12 years) BUTALBITAL COMP. W/ CODEINE (age >12 years) CODEINE SULFATE (age >12 years) ENDOCET [®] (oxycodone w/ acetaminophen) HYDROCODONE (plain, w/acetaminophen, or w/ibuprofen) (some exceptions apply) <i>QTY LIMIT</i> : Hydrocodone/APAP 12 tablets/day HYDROMORPHONE tablets (compare to Dilaudid [®]) MORPHINE SULFATE OXYCODONE (plain) OXYCODONE (w/acetaminophen, w/aspirin or w/ibuprofen) <i>QTY LIMIT</i> : Oxycodone/APAP 12 tablets/day TRAMADOL <i>QTY LIMIT</i> : 8 tablets/day (Age ≥ 16)	Acetaminophen w/hydrocodone: <i>all branded products</i> <i>QTY LIMIT</i> : = 12 tablets/day Acetaminophen w/oxycodone: <i>all branded products</i> <i>QTY LIMIT</i> : = 12 tablets/day Actiq [®] (fentanyl lozenge on a stick: 200 mcg, 400 mcg, 600 mcg, 800 mcg, 1200 mcg, 1600 mcg) Apadaz [®] (benzhydrocodone/APAP) <i>QTY LIMIT</i> : 12 tablets/day Benzhydrocodone/APAP (compare to Apadaz [®]) <i>QTY LIMIT</i> : 12 tablets/day Butorphanol Nasal Spray <i>QTY LIMIT</i> : 2 bottles/month Demerol (meperidine) Dilaudid [®] (hydromorphone) tablets Dilaudid-5 [®] (hydromorphone) oral solution Fentanyl citrate transmucosal (compare to Actiq [®]) Fentora [®] (fentanyl citrate buccal tablets)	Note : The initial fill for all short-acting opiates will be limited to 50 Morphine Milligram Equivalents (MME) and 7-day supply for patients ≥ 18 years of age OR 24 MME and 3-day supply for patients ≤ 17 years of age. Butorphanol Nasal Spray : documented site effect, allergy, treatment failure, or contraindication to codeine, hydrocodone, morphine, & oxycodone (all 4 generic entities) as single or combination products. OR is unable to use tablet or liquid formulations. Actiq, Fentanyl transmucosal, Fentora : indication of cancer breakthrough pain AND patient is opioid tolerant AND is on a long acting opioid formulation AND is 18 years of age or older (Actiq 16 years of age or older) AND prescriber is registered in the Transmucosal Immediate Release Fentanyl (TIRF) Risk Evaluation and Mitigation Strategy (REMS) Access program AND member has had a documented treatment failure with or intolerance to 2 of the following 3 immediate release treatment options: morphine, hydromorphone or oxycodone. OR is unable to use tablet or liquid formulations AND if the request is for brand name Actiq, member has a documented intolerance to generic fentanyl transmucosal. Dilaudid - 5 Oral Solution, Hydromorphone Oral Solution : member has had a documented side effect, allergy or treatment failure with oxycodone oral
--	---	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>TRAMADOL/APAP <i>QTY LIMIT:</i> 8 tablets/day (Age ≥18)</p> <p>**NOTE: As of 5/1/21, a completed safety checklist must be submitted for new patients exceeding 90 MME per day, and existing patients exceeding 120 MME per day (applies to any combination of short and/or long acting opioids)**</p> <p>Note: The FDA restricts the use of prescription codeine pain and cough medicines in children. Prior authorization is required for patients <12 years of age.</p>	<p>Hydromorphone oral solution (compare to Dilaudid-5®) Meperidine <i>QTY LIMIT:</i> 30 tablets/5-day supply per 30 days Oxycodone (plain) capsules Oxymorphone (compare to Opana®) Pentazocine w/naloxone Seglentis® (celecoxib/tramadol) oral tablet Tramadol oral solution 5mg/ml</p>	<p>solution and morphine oral solution OR has been started and stabilized on another dosage form of hydromorphone AND if the request is for the branded product, patient has a documented intolerance to the generic product.</p> <p>Oxycodone (generic) Capsules: member has a documented intolerance to generic oxycodone tablets.</p> <p>Seglentis: The patient has a documented side effect, allergy, or treatment failure with two or more preferred agents AND the patient is unable to take the individual components separately</p> <p>Tramadol Oral Solution: patient has a medical necessity for a non-solid oral dosage form. (e.g. swallowing disorder).</p> <p>Ultracet: member has a documented intolerance to the generic formulation</p> <p>Other Short acting Opioids: member has had a documented side effect, allergy, or treatment failure to at least 3 medications not requiring prior approval. (If a product has an AB rated generic, one trial must be the generic.)</p> <p>PA requests to exceed daily cumulative MME limits:</p> <ul style="list-style-type: none"> • Non-Opioid alternatives (up to a maximum dose recommended by the FDA) and Non-Pharmacological Treatments have been considered, and any appropriate treatments are documented in the patient’s medical records. Such treatments may include, but are not limited to: NSAIDs, Acetaminophen, Acupuncture, Chiropractic, Physical Therapy. • Vermont Prescription Monitoring System (VPMS) has been queried. • Patient education and informed consent have been obtained, and a Controlled Substance Treatment Agreement is included in the patient’s medical record. • A reevaluation of the effectiveness and safety of the patient’s pain management plan, including an assessment of the patient’s adherence to the treatment regimen is completed no less than once every 90 days. • Patient has a valid prescription for or states they are in possession of naloxone. • Patients in nursing homes, receiving or eligible for hospice services, or those with chronic pain associated with cancer or cancer treatment are exempt from these requirements. <p>Limitations: APAP containing products: daily doses that result in > 4 grams of acetaminophen/day will reject for PA</p>
OPIOIDS: LONG ACTING		
<p><u>TRANSDERMAL</u> BUTRANS (buprenorphine) TRANSDERMAL SYSTEM <i>QTY LIMIT:</i> 4 patches/28 days (Maximum 28-day fill) FENTANYL PATCH (compare to Duragesic®) <i>QTY LIMIT:</i> 12 mcg/hr, 25 mcg/hr, 50 mcg/hr = 15</p>	<p>Buprenorphine patch (compare to Butrans®) <i>QTY LIMIT:</i> 4 patches/28 days (Maximum 28-day Fill) Fentanyl patch 37.5 mcg/hr, 62.5 mcg/hr, 87.5 mcg/hr</p>	<p>CLINICAL CONSIDERATIONS: Long-acting opioid dosage forms are intended for use in opioid tolerant patients only. These tablet/capsule/topical medication strengths may cause fatal respiratory depression when administered to patients not previously exposed to opioids. LA opioids should be prescribed for patients with a diagnosis or condition that requires a continuous, around-the-clock analgesic. LA opioids should be reserved for use in patients for whom alternative treatment options (e.g., non-opioid</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>patches/30 days, 75 mcg/hr, 100 mcg/hr = 30 patches/30 days</p> <p><u>BUCCAL</u> All products require PA</p> <p><u>ORAL</u> MORPHINE SULFATE CR 12 hr tablet (compare to MS Contin[®]) <i>QTY LIMIT</i>: 90 tablets/strength/30 days</p>	<p>Belbuca[®] (buprenorphine hcl buccal film) <i>QTY LIMIT</i>: 56 films/28 days (Maximum 28-day fill)</p> <p>Conzip[®] (tramadol ER biphasic release) capsule <i>QTY LIMIT</i>: 1 capsule/day</p> <p>Hydromorphone XR tablet <i>QTY LIMIT</i>: 30 tablets/30 days (8 mg, 12 mg, 16 mg tabs)</p> <p>Methadone 5 mg, 10 mg tablets</p> <p>Methadone oral solution (no PA required for patient less than 1 year old)</p> <p>Methadone oral concentrate 10 mg/ml</p> <p>Morphine sulfate SR 24hr capsule <i>QTY LIMIT</i>: 60 capsules/strength/30 days</p> <p>Morphine sulfate SR beads 24hr capsule <i>QTY LIMIT</i>: 30 capsules/strength/30 days</p> <p>MS Contin[®] (morphine sulfate CR 12 hr) tablets <i>QTY LIMIT</i>: 90 tablets/strength/30 days</p> <p>Oxymorphone ER <i>QTY LIMIT</i>: 60 tablets/strength/30 days</p> <p>Tramadol SR <i>QTY LIMIT</i>: 1 tablet/day</p> <p>Tramadol ER biphasic-release[®] capsule <i>QTY LIMIT</i>: 150 mg = 1 capsule/day</p> <p>Tramadol ER biphasic-release tablet (formerly Ryzolt[®]) <i>QTY LIMIT</i>: 1 tablet/day</p>	<p>analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain. LA opioids are NOT intended for use as 'prn' analgesic. LA opioids are NOT indicated for pain in the immediate post-operative period (the first 12-24 hours following surgery) or if the pain is mild, or not expected to persist for an extended period of time. LA opioids are not intended to be used in a dosage frequency other than FDA approved regimens. Patients should not be using other extended-release opioids prescribed by another physician. Prescribers should consult the VPMS (Vermont Prescription Monitoring System) to review a patient's Schedule II - IV medication use before prescribing long-acting opioids.</p> <p>Oral Non-Preferred (except methadone & tramadol containing products): the patient has had a documented side effect, allergy, or at least two preferred products. (If a product has an AB rated generic, there must have been a trial of the generic).</p> <p>Belbuca Films, Buprenorphine Patch: the patient has had a documented intolerance to Butrans patches</p> <p>Fentanyl patches 37.5mcg/hr, 62.5mcg/hr, 87.5mcg/hr: provider must submit clinical rationale detailing why the patient is unable to use a combination of the preferred strengths.</p> <p>Methadone Tablet: patient has had a documented side effect, allergy, or treatment failure to morphine sulfate CR 12 hr tablets AND the initial methadone daily dose does not exceed 30mg (Note: Methadone products, when used for treatment of opioid addiction in detoxification or maintenance programs, shall be dispensed ONLY by certified opioid treatment programs as stipulated in 42 CFR 8.12, NOT retail pharmacy.) Note: Methadone 40mg dispersible tablet not approved for retail dispensing.</p> <p>Methadone Liquid: Patient must have a medical necessity for an oral liquid (i.e. swallowing disorder, inability to take oral medications) AND the initial daily dose does not exceed 30mg OR patient has been started and stabilized on the requested oral liquid medication. (Note: Methadone products, when used for treatment of opioid addiction in detoxification or maintenance programs, shall be dispensed ONLY by certified opioid treatment programs as stipulated in 42 CFR 8.12, NOT retail pharmacy.)</p> <p>Conzip, Tramadol ER biphasic-release Capsule, Tramadol ER biphasic-release Tablet, Tramadol ER/SR: member has had a documented treatment failure to a preferred short-acting tramadol product. In addition, for approval of tramadol ER biphasic-release capsule or tablet or the patient must have a documented intolerance to generic tramadol ER/SR.</p> <p>OxyContin: Patient has diagnosis of cancer, sickle cell disease, or is on hospice/palliative care OR patient was previously maintained on a long-acting oxycodone formulation OR clinical justification is provided for why patient cannot use preferred products</p> <p>Hysingla ER: Available with PA for those unable to tolerate any preferred medications. All requests will go to the DVHA Medical Director for approval.</p> <p>PA requests to exceed daily cumulative MME limits:</p> <ul style="list-style-type: none"> • Non-Opioid alternatives (up to a maximum dose recommended by the

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>ORAL, ABUSE-DETERRENT FORMULATIONS</u> All products require PA</p> <p>**NOTE: As of 5/1/21, a completed safety checklist must be submitted for new patients exceeding 90 MME per day, and existing patients exceeding 120 MME per day (applies to any combination of short and/or long acting opioids)**</p>	<p>Hysingla ER® (hydrocodone bitartrate) <i>QTY LIMIT:</i> 1 tablet/ day</p> <p>OxyContin® (Oxycodone ER) <i>QTY LIMIT:</i> 90 tablets/strength/30 days</p>	<p>FDA) and Non-Pharmacological Treatments have been considered, and any appropriate treatments are documented in the patient’s medical records. Such treatments may include, but are not limited to: NSAIDs, Acetaminophen, Acupuncture, Chiropractic, Physical Therapy.</p> <ul style="list-style-type: none"> • Vermont Prescription Monitoring System (VPMS) has been queried. • Patient education and informed consent have been obtained, and a Controlled Substance Treatment Agreement is included in the patient’s medical record. • A reevaluation of the effectiveness and safety of the patient’s pain management plan, including an assessment of the patient’s adherence to the treatment regimen is completed no less than once every 90 days. • Patient has a valid prescription for or states they are in possession of naloxone. • Patients in nursing homes, receiving or eligible for hospice services, or those with chronic pain associated with cancer or cancer treatment are exempt from these requirements.
NSAIDS		
<p><u>ORAL SINGLE AGENT</u> DICLOFENAC POTASSIUM DICLOFENAC SODIUM ETODOLAC FLURBIPROFEN IBUPROFEN INDOMETHACIN INDOMETHACIN ER KETOPROFEN KETOROLAC <i>QTY LIMIT:</i> 20 doses/5 day supply every 90 day MECLOFENAMATE SODIUM MEFANAMIC ACID capsules MELOXICAM tabs NABUMETONE NAPROXEN 250 mg, 375 mg, 500 mg NAPROXEN ENTERIC COATED 375 mg, 500 mg NAPROXEN SODIUM 275mg, 550mg NAPROXEN SODIUM OTC 220 mg</p>	<p>Cambia® (diclofenac potassium) packet for oral solution <i>QTY LIMIT:</i> 9 packets/month</p> <p>Daypro® (oxaprozin) Etodolac ER</p> <p>Feldene® (piroxicam) Fenoprofen 400 mg cap Fenoprofen 600 mg tab</p> <p>Indocin® (indomethacin) suspension Ketoprofen ER Lofena™ (diclofenac) tablet Meloxicam capsule (compare to Vivlodex®)</p> <p>Nalfon® (fenoprofen) 400 mg capsules</p> <p>Naprelan® (naproxen sodium ER) Naproxen oral suspension Naproxen sodium ER Naproxen suspension 125mg/5ml Relafen® DS (nabumetone) Zipsor® (diclofenac potassium)</p>	<p>Arthrotec, diclofenac/misoprostol, Duexis: patient has a documented side effect or treatment failure to 2 or more preferred generic NSAIDs OR patient is not a candidate for therapy with a preferred generic NSAID mono-therapy due to one of the following: patient is 60 years of age or older, Patient has a history of GI bleed, Patient is currently taking an oral corticosteroid, Patient is currently taking methotrexate AND for approval of diclofenac/misoprostol, the patient must have a documented intolerance to brand Arthrotec</p> <p>Cambia: drug is being prescribed for treatment of acute migraine attacks AND patient has had a documented side effect or treatment failure to 2 or more preferred generic NSAIDs, one of which must be generic diclofenac OR drug is being prescribed for treatment of acute migraine attacks AND patient has a requirement for an oral liquid dosage form (i.e. swallowing disorder, inability to take oral medications) AND patient has had a documented side effect or treatment failure with the generic ibuprofen suspension.</p> <p>Celebrex: patient has had a documented intolerance to generic celecoxib.</p> <p>Pennsaid: patient has had a documented side effect or inadequate response to Diclofenac gel or topical solution.</p> <p>Diclofenac Patch, Licart: patient has had a documented side effect or inadequate response to Diclofenac gel or topical solution AND patient has a documented intolerance to brand Flector Patch.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>OXAPROZIN (compare to Daypro®) PIROXICAM (compare to Feldene®) SULINDAC</p> <p><u>ORAL</u> <u>COX-II Selective</u> CELECOXIB <i>QTY LIMIT: 2 caps/day</i></p> <p><u>INJECTABLE</u> KETOROLAC Injection (formerly Toradol®) <i>QTY LIMIT: 1 dose per fill</i></p> <p><u>NASAL SPRAY</u> All products require PA</p> <p><u>TOPICAL</u> DICLOFENAC (compare to Voltaren®) gel 1% DICLOFENAC 1.5 % Topical Solution</p> <p><u>TRANSDERMAL</u> Flector® (diclofenac) 1.3 % Patch <i>QTY LIMIT: 2 patches/day</i></p> <p><u>NSAID/ANTI-ULCER</u> All products require PA</p> <p>Note: Please refer to “Dermatological: Actinic Keratosis Therapy” for Solaraze® or Diclofenac 3% Gel</p>	<p>Zorvolex® (diclofenac) Capsules <i>QTY LIMIT: 3 capsules/day</i></p> <p>Celebrex® (celecoxib) capsule <i>QTY LIMIT: 2 caps/day</i> Elyxyb™ (celecoxib) oral solution</p> <p>Sprix® (ketorolac) Nasal Spray <i>QTY LIMIT: 5 bottles/5 days – once every 90 days</i></p> <p>Pennsaid® (diclofenac) 2% Topical Solution</p> <p>Diclofenac (compare to Flector®) 1.3% Patch <i>QTY LIMIT: 2 patches/day</i> Licart® (diclofenac epolamine) 1.3% Patch <i>QTY LIMIT: 1 patch/day</i></p> <p>Arthrotec® (diclofenac sodium w/misoprostol) Diclofenac sodium w/misoprostol (compare to Arthrotec®)</p> <p>Duexis® (ibuprofen/famotidine) <i>QTY LIMIT: 3 tablets/day</i> Ibuprofen/famotidine (compare to Duexis®) <i>QTY LIMIT: 3 tablets/day</i> Naproxen/esomeprazole (compare to Vimovo®)</p> <p>Vimovo® (naproxen/esomeprazole) <i>QTY LIMIT: 2 tablets/day</i></p>	<p>Duexis, Ibuprofen/famotidine, naproxen/esomeprazole, Vimovo: patient is unable to take the individual components separately AND for approval of ibuprofen/famotidine or naproxen/esomeprazole, the patient must have a documented intolerance to the brand name equivalent.</p> <p>Elyxyb: drug is being prescribed for treatment of acute migraine attacks AND patient has had a documented side effect or treatment failure to 2 or more preferred generic NSAIDs, one of which must be generic celecoxib OR drug is being prescribed for treatment of acute migraine attacks AND patient has a requirement for an oral liquid dosage form (i.e. swallowing disorder, inability to take oral medications) AND patient has had a documented side effect or treatment failure with the generic ibuprofen suspension.</p> <p>Lofena, Zipsor, Zorvolex: patient has had a documented side effect, allergy, or treatment failure to 4 or more preferred generic NSAIDs, one of which must be generic diclofenac.</p> <p>Meloxicam Capsule: patient has had a documented side effect, allergy, or treatment failure to 4 or more preferred generic NSAIDs, one of which must be generic meloxicam tablet.</p> <p>Naproxen suspension: patient has a requirement for an oral liquid dosage form (i.e. swallowing disorder, inability to take oral medications) AND patient has had a documented side effect or treatment failure with generic ibuprofen suspension.</p> <p>Relafen DS: patient has had a documented side effect, allergy, or treatment failure to 4 or more preferred generic NSAIDs, one of which must be generic nabumetone.</p> <p>Sprix: indication or diagnosis is moderate to moderately severe pain. AND patient has had a documented inadequate response or intolerance to generic ketorolac tablets. OR patient has a documented medical necessity for the specialty dosage form (i.e. inability to take medication orally (NPO)).</p> <p>All other PA requiring NSAIDs: patient has had a documented side effect or treatment failure to 2 or more preferred generic NSAIDs. (If a product has an AB rated generic, one trial must be the generic.) AND if the request is for a non-preferred extended-release formulation, the patient has not been able to adhere to the dosing schedule of the immediate release formulation resulting in significant clinical impact.</p>

ANKYLOSING SPONDYLITIS: INJECTABLES

Length of Authorization: Initial PA 3 months; 12 months thereafter

Preferred After Clinical Criteria Are Met

INJECTABLE

ADALIMUMAB-ADBM (compare to Cyltezo®) biosimilar to Humira®

AVSOLA® (infliximab-axxq) biosimilar to Remicade®

ENBREL® (etanercept)

QTY LIMIT: 50 mg = 4 syringes/28 days,
25 mg = 8 syringes/28 days

HUMIRA® (adalimumab)

QTY LIMIT: 2 syringes/28 days

INFLECTRA® (infliximab-dyyb) biosimilar to Remicade®

TALTZ® (ixekizumab)

QTY LIMIT: 80 mg prefilled syringe or autoinjector = 2/28 days for the first month and 1/28 days subsequently

ORAL

XELJANZ® (tofacitinib) tablet

QTY LIMIT: 2 tablets/day

XELJANZ® XR (tofacitinib) tablet

QTY LIMIT: 1 tablet/day

Maximum 30 days supply

XELJANZ® (tofacitinib) oral solution

Abrilada™ (adalimumab-afzb) biosimilar to Humira®
Adalimumab-adaz (compare to Hyrimoz®) biosimilar to Humira®

Adalimumab-fkjp (compare to Hulio®) biosimilar to Humira®

Amjevita™ (adalimumab-atto) biosimilar to Humira®

Cimzia® (certolizumab pegol)

QTY LIMIT: 1 kit/28 days (starter X 1, then regular)

Cosentyx® (secukinumab)

Cyltezo® (adalimumab-adbm) biosimilar to Humira®

Hadlima™ (adalimumab-bwvd) biosimilar to Humira®

Hulio® (adalimumab-fkjp) biosimilar to Humira®

Hyrimoz® (adalimumab-adaz) biosimilar to Humira®

Idacio® (adalimumab-aacf) biosimilar to Humira®

Remicade® (infliximab)

Renflexis™ (infliximab-abda) biosimilar to Remicade®

Simlandi® (adalimumab-ryvk) biosimilar to Humira®

Simponi® (golimumab) Subcutaneous

QTY LIMIT: 50 mg prefilled syringe or autoinjector = 1/28 days

Simponi® (golimumab) Aria Vial

Yuflyma® (adalimumab-aaty) biosimilar to Humira®

Yusimry™ (adalimumab-aqvh) biosimilar to Humira®

Rinvoq® (upadactinib) extended release tablet

QTY LIMIT: 1 tablet/day

Clinical Criteria:

For all drugs: patient has a diagnosis of ankylosing spondylitis (AS) and has already been stabilized on the medication being requested. OR patient has a confirmed diagnosis of AS, and conventional NSAID treatment and DMARD therapy (e.g. methotrexate therapy) resulted in an adverse effect, allergic reaction, inadequate response, or treatment failure. If methotrexate is contraindicated, another DMARD should be tried.

Additional criteria for Cimzia, Simponi, Taltz, Xeljanz, Xeljanz XR: the patient had a trial and failure or contraindication to a preferred TNF Inhibitor.

Additional criteria for Simponi Aria: The patient meets the FDA approved age AND the patient has not responded adequately to Simponi subcutaneous AND The prescriber must provide a clinically valid reason why at least 2 preferred agents cannot be used.

Additional criteria for Cosentyx: the patient had a trial and failure or contraindication to a preferred TNF Inhibitor and Taltz.

Additional Criteria for Humira Biosimilars: the patient must be unable to use Humira.

Additional criteria for Remicade, Renflexis: the prescriber must provide a clinically valid reason why at least 2 preferred agents cannot be used, and the patient must be unable to use Avsola or Inflectra.

Additional Criteria for Rinvoq: the prescriber must provide a clinically valid reason why at least 2 preferred agents cannot be used, one of which must be Xeljanz or Xeljanz XR.

* Patients with documented diagnosis of active axial involvement should have a trial with two NSAIDs, but a trial with DMARD is not required. If no active axial skeletal involvement, then NSAID trial and a DMARD trial are required (unless otherwise contraindicated).

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

ANTI-ANXIETY: ANXIOLYTICS

BENZODIAZEPINE

CHLORDIAZEPOXIDE (formerly Librium[®])
 CLONAZEPAM (compare to Klonopin[®])
QTY LIMIT: 4 tabs/day except 2 mg.
 2 mg = 3 tabs/day
 CLONAZEPAM ODT
QTY LIMIT: 4 tabs/day except 2 mg.
 2 mg = 3 tabs/day
 DIAZEPAM (compare to Valium[®])
 LORAZEPAM (compare to Ativan[®])
QTY LIMIT: 4 tablets/day
 OXAZEPAM

Alprazolam (compare to Xanax[®])
QTY LIMIT: 4 tablets/day
 Alprazolam ER, Alprazolam XR[®] (compare to Xanax XR[®])
QTY LIMIT: 2 tablets/day
 Alprazolam ODT
QTY LIMIT: 3 tablets/day
 Alprazolam Intensol[®] (alprazolam concentrate)
 Ativan[®] (lorazepam)
QTY LIMIT: 4 tablets/day
 Clorazepate tabs (compare to Tranxene T[®])
 Diazepam Intensol[®] (diazepam concentrate)
 Klonopin[®] (clonazepam)
QTY LIMIT: 4 tabs/day except 2 mg.
 2 mg = 3 tabs/day
 Lorazepam Intensol[®] (lorazepam concentrate)
 Loreev XR[™] (lorazepam extended release)
 Tranxene T[®] (clorazepate tablets)
 Valium[®] (diazepam)
 Xanax[®] (alprazolam)
QTY LIMIT: 4 tablets/day
 Xanax XR[®] (alprazolam XR)
QTY LIMIT: 2 tablets/day

Non-preferred Benzodiazepines (except for Alprazolam ODT, Intensol Products, and Loreev XR): patient has a documented side effect, allergy, or treatment failure to at least 2 preferred benzodiazepine medications. (If a product has an AB rated generic, there must also be a trial of the generic formulation.)
Alprazolam ODT: patient has a documented side effect, allergy, or treatment failure to at least 2 preferred benzodiazepine medications. (If a product has an AB rated generic, there must also be a trial of the generic formulation). OR patient has a medical necessity for disintegrating tablet administration (i.e. inability to swallow tablets) AND patient has a documented side effect, allergy or treatment failure to clonazepam ODT.
Alprazolam Intensol, Diazepam Intensol, and Lorazepam Intensol: patient has a medical necessity for the specialty dosage form (i.e. swallowing disorder). AND the medication cannot be administered by crushing oral tablets.
Loreev XR: The patient is receiving a stable dose of lorazepam tablets, evenly divided, three times daily AND medical reasoning for use beyond convenience or enhanced compliance is provided.

NON-BENZODIAZEPINE

BUSPIRONE (formerly Buspar[®])
 HYDROXYZINE HYDROCHLORIDE (formerly Atarax[®])
 HYDROXYZINE PAMOATE (compare to Vistaril[®])
 (all strengths except 100 mg)
 MEPROBAMATE

Hydroxyzine Pamoate (100 mg strength ONLY)
 (compare to Vistaril[®])
 Vistaril[®] (hydroxyzine pamoate)

Hydroxyzine Pamote 100mg strength ONLY: patient is unable to use generic 50 mg capsules.
Vistaril: patient has a documented intolerance to the generic formulation.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

ANTICOAGULANTS

ORAL		
<p><u>VITAMIN K ANTAGONIST</u> WARFARIN</p> <p><u>DIRECT THROMBIN INHIBITOR</u> PRADAXA® (dabigatran etexilate) capsule</p> <p><u>FACTOR XA INHIBITOR</u> ELIQUIS® (apixaban) XARELTO® (rivaroxaban) tablet</p> <p><u>Preferred After Clinical Criteria Are Met</u> XARELTO® (rivaroxaban) 2.5 mg tablet</p>	<p>Dabigatran Etexilate (compare to Pradaxa®) capsule Pradaxa® (dabigatran etexilate) oral pellets</p> <p>Savaysa® (edoxaban) <i>QTY LIMIT: 1 tablet/daily</i> Xarelto® (rivaroxaban) oral suspension</p>	<p>Dabigatran: the patient must have a documented intolerance to brand name Pradaxa</p> <p>Pradaxa pellets: patient has a medical necessity for a non-solid oral dosage form and prescriber has provided a clinically valid reason why Xarelto suspension cannot be used.</p> <p>Savaysa: creatinine clearance is documented to be < 95 ml/min AND prescriber has provided another clinically valid reason why generic warfarin, Pradaxa, Xarelto or Eliquis cannot be used. A yearly creatinine clearance is required with renewal of PA request</p> <p>Xarelto suspension: patient has a medical necessity for a non-solid oral dosage form (e.g. swallowing disorder).</p> <p>Xarelto 2.5 mg: Patient has a diagnosis of chronic coronary artery disease (CAD) or peripheral artery disease (PAD) AND medication is being used concurrently with aspirin.</p>
INJECTABLE		
<p><u>UNFRACTIONATED HEPARIN INJECTABLE</u> HEPARIN</p> <p><u>LOW MOLECULAR WEIGHT HEPARINS INJECTABLE</u> ENOXAPARIN (compare to Lovenox®) <i>QTY LIMIT: 2 syringes/day calculated in ml volume</i></p> <p><u>SELECTIVE FACTOR XA INHIBITON INJECTABLE</u> All products require PA</p>	<p>Fragmin® (dalteparin) Lovenox® (enoxaparin) <i>QTY LIMIT: 2 syringes/day calculated in ml volume</i></p> <p>Arixtra® (fondaparinux) Fondaparinux (compare to Arixtra®)</p>	<p>Arixtra, Fondaparinux, Lovenox and Fragmin: patient has a documented intolerance to generic enoxaparin AND if the request is for brand Arixtra, the patient must also have a documented intolerance to generic fondaparinux.</p>

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

ANTICONVULSANTS

ORAL

CARBAMAZEPINE tablets (compare to Tegretol[®])
CARBAMAZEPINE capsules (compare to Carbatrol[®])
CARBAMAZEPINE extended release (compare to Tegretol XR[®])
CELONTIN[®] (methsuxamide)
CLOBAZAM (compare to Onfi[®])
QTY LIMIT: 10 mg = 3 tabs/day, 20 mg = 2 tabs/day, oral suspension = 16mL/day (40mg/day)
CLONAZEPAM (compare to Klonopin[®])
QTY LIMIT: 4 tablets/day
CLONAZEPAM ODT (formerly Klonopin Wafers[®])
QTY LIMIT: 4 tablets/day
DIAZEPAM (compare to Valium[®])
DILVALPROEX SODIUM capsules (compare to Depakote Sprinkles[®])
DIVALPROEX SODIUM (compare to Depakote[®])
DIVALPROEX SODIUM ER (compare to Depakote ER[®])
EPITOL (carbamazepine)
ETHOSUXAMIDE (compare to Zarontin[®])
GABAPENTIN 100 mg, 300 mg, 400 mg capsules, 600 mg, 800 mg tablets, 250 mg/5 ml oral solution (compare to Neurontin[®])
LACOSAMIDE (compare to Vimpat[®]) tabs, solution
LAMOTRIGINE chew tabs (compare to Lamictal[®] chew tabs)
LAMOTRIGINE tabs (compare to Lamictal[®] tabs)
LEVETIRACETAM tabs (compare to Keppra[®] tabs)
LEVETIRACETAM oral solution (compare to Keppra[®] oral solution)
LEVETIRACETAM ER (compare to Keppra XR[®])
LYRICA[®] (pregabalin) capsules
QTY LIMIT: 3 capsules/day
LYRICA[®] (pregabalin) oral solution
OXCARBAZEPINE tablets (compare to Trileptal[®])

Aptiom[®] (eslicarbazepine acetate)
QTY LIMIT: 200, 400 = 1 tab/day
600 mg, 800 mg = 2 tabs/day
Banzel[®] (rufinamide)
QTY LIMIT: 400 mg = 8 tabs/day, 200 mg = 16 tabs/day
Banzel[®] (rufinamide) oral suspension
QTY LIMIT: 80 ml/day (3,200 mg/day)
Briviact[®] (brivaracetam) tablets, oral suspension
Carbatrol[®] (carbamazepine) capsules
Clorazepate (compare to Tranxene-T[®]) tablets
Depakote[®] (divalproex sodium)
Depakote ER[®] (divalproex sodium)
Depakote Sprinkles[®] (divalproex sodium caps)
Diacomit[®] (stiripentol)
Dilantin[®] (phenytoin) chewable tablets, capsules, suspension
Elepsia[™] (levetiracetam) extended release
Eprontia[™] (topiramate) oral solution
Felbamate (compare to Felbatol[®])
Fintepla[®] (fenfluramine) oral solution
Felbatol[®] (felbamate)
Fycompa[®] (perampanel) tablets
QTY LIMIT: 1 tablet/day
Keppra[®]* (levetiracetam) tablets, oral solution
Keppra XR[®] (levetiracetam extended release)
Klonopin[®] (clonazepam)
QTY LIMIT: 4 tablets/day
Lamictal[®] tabs (lamotrigine tabs)
Lamictal[®] chew tabs (lamotrigine chew tabs)
Lamictal ODT[®] (lamotrigine orally disintegrating tablets)
Lamictal XR[®] tablets (lamotrigine extended release)
Lamotrigine ER (compare to Lamictal XR[®])
Lamotrigine ODT (compare to Lamictal ODT[®])

Criteria for approval of ALL non-preferred drugs: patient has been started and stabilized on the requested medication (Note: samples are not considered adequate justification for stabilization.) OR patient meets additional criteria outlined below.

Aptiom: the diagnosis is adjunctive therapy of partial-onset seizures and the patient has had a documented side effect, allergy, treatment failure/inadequate response or a contraindication to at least TWO preferred anticonvulsants, one of which is oxcarbazepine.

Banzel, Rufinamide: diagnosis or indication is treatment of Lennox-Gastaut Syndrome. AND patient has had a documented side effect, allergy, treatment failure/inadequate response or a contraindication to at least TWO preferred anticonvulsants used for the treatment of Lennox-Gastaut syndrome (topiramate, lamotrigine, valproic acid) AND for approval of the oral suspension, patient must have medical necessity for a specialty dosage form AND for approval of generic rufinamide, the patient must have a documented intolerance to brand Banzel.

Briviact: the diagnosis is adjunctive therapy of partial-onset seizures and the patient has had a documented side effect, allergy, treatment failure/inadequate response, or a contraindication to at least TWO preferred anticonvulsants, one of which is levetiracetam.

Carbatrol, Depakote, Depakote ER, Depakote Sprinkles, Dilantin, Keppra tablets or oral solution, Klonopin, Klonopin Wafers, Lamictal tablets or chew tablets, Mysoline, Neurontin capsules, tablets, solution, Onfi, Phenytek, Tegretol tablets, Topamax tabs, Topamax sprinkles, Trileptal tablets, Trileptal oral suspension, Vimpat, Zarontin: patient has had a documented intolerance to the generic equivalent of the requested medication.

Clorazepate, Fycompa, Tranxene-T: diagnosis is adjunctive therapy of partial-onset seizures OR diagnosis is adjunctive therapy for primary generalized tonic-clonic seizures (Fycompa only) AND the patient has had a documented side effect, allergy, treatment failure, inadequate response, or a contraindication to at least TWO preferred anticonvulsants. AND for approval of Tranxene-T the patient must have a documented intolerance to the generic equivalent.

Diacomit: Diagnosis or indication is treatment of Dravet Syndrome AND neutrophil and platelet counts have been obtained prior to starting therapy and are monitored periodically thereafter AND Patient is unable to tolerate or has had an inadequate response to valproate and clobazam AND medication will be used concurrently with clobazam. **Note:** There are no clinical data to support the use of Diacomit as monotherapy.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>OXCARBAZEPINE oral suspension (compare to Trileptal[®])</p> <p>PHENYTOIN (compare to Dilantin[®])</p> <p>PHENYTOIN EX cap (compare to Phenytek[®])</p> <p>PREGABALIN capsules (compare to Lyrica) <i>QTY LIMIT: 3 capsules/day</i></p> <p>PRIMIDONE (compare to Mysoline[®])</p> <p>TEGRETOL[®] (carbamazepine) suspension</p> <p>TEGRETOL XR[®] (carbamazepine)</p> <p>TOPIRAMATE tabs (compare to Topamax[®] tabs)</p> <p>TOPIRAMATE sprinkle caps (compare to Topamax[®] Sprinkles)</p> <p>VALPROIC ACID</p> <p>ZONISAMIDE</p> <p><u><i>Preferred After Clinical Criteria Are Met</i></u></p> <p>EPIDIOLEX[®] (cannabidiol) oral solution</p>	<p>Libervant (diazepam) buccal film <i>QTY LIMIT: 10 films/month</i></p> <p>Motpoly XR[™] (lacosamide ER) capsule <i>QTY LIMIT: 100 mg capsules = 1/day, 150 mg and 200 mg capsules = 2/day</i></p> <p>Mysoline[®] (primidone)</p> <p>Neurontin[®] (gabapentin) capsules, tablets and solution</p> <p>Onfi[®] (clobazam) Oral Suspension 2.5 mg/ml <i>QTY LIMIT: 16 ml/day</i></p> <p>Onfi[®] (clobazam) Tablets <i>QTY LIMIT: 10 mg = 3 tabs/day, 20 mg = 2 tabs/day</i></p> <p>Oxtellar[®] XR (oxcarbazepine ER) tablet</p> <p>Pregabalin oral solution (compare to Lyrica[®])</p> <p>Qudexy[®] XR (topiramate) capsules</p> <p>Rufinamide (compare to Banzel[®]) tablet, oral suspension <i>QTY LIMIT: 400 mg = 8 tabs/day, 200 mg = 16 tabs/day, oral suspension = 80 ml/day (3200 mg/day)</i></p> <p>Sabril[®] (vigabatrin)</p> <p>Spritam[®] (levetiracetam) tablets for oral suspension</p> <p>Sympazan[®] (clobazam) films</p> <p>Tegretol[®] (carbamazepine) tablets</p> <p>Tiagabine (compare to Gabitril[®])</p> <p>Topamax[®] (topiramate) tablets</p> <p>Topamax[®] (topiramate) Sprinkle Capsules</p> <p>Topiramate ER sprinkle capsules (compare to Qudexy[®] XR)</p> <p>Topiramate SR 24hr (compare to Trokendi[®]) capsules <i>QTY LIMIT: 200 mg = 2 caps/day, all other strengths = 1 cap/day</i></p> <p>Tranxene-T[®] (clorazepate) tablets</p> <p>Trileptal[®] tablets (oxcarbazepine)</p> <p>Trileptal[®] oral suspension (oxcarbazepine)</p> <p>Trokendi XR[®] (topiramate SR 24hr) capsules <i>QTY LIMIT: 200 mg = 2 caps/day, all other strengths = 1 cap/day</i></p> <p>Vigabatrin (compare to Sabril[®])</p> <p>Vimpat[®] (lacosamide) tablets, oral solution</p> <p>Xcopri[®] (cenobamate) tablets <i>QTY LIMIT: 200 mg = 2 tabs/day, all other strengths = 1 tab/day</i></p>	<p>Eprontia, Zonisade: The patient has a medical necessity for specialty dosage form.</p> <p>Epidiolex: The patient is unable to tolerate or has had an inadequate response to at least 2 of the following medications: clobazam, levetiracetam, valproate, lamotrigine, topiramate, rufinamide, or felbamate Note: This is processed via automated (electronic step therapy)</p> <p>Felbamate, Felbatol: patient information/consent describing aplastic anemia and liver injury has been completed AND diagnosis is adjunctive therapy of partial-onset seizures or Lennox-Gastaut seizures and the patient has had a documented side effect, allergy, treatment failure/inadequate response or a contraindication to at least THREE preferred anticonvulsants. Additionally, if brand is requested, the patient has a documented intolerance to the generic product.</p> <p>Fintepla: Diagnosis or indication is treatment of Dravet Syndrome or Lennox-Gastaut Syndrome AND patient has had a documented side effect, allergy, treatment failure/inadequate response or contraindication to at least two preferred anticonvulsants and Epidiolex AND prescriber, pharmacy and patient are registered with the REMS programs AND for reapproval, the patient must have a documented decrease from baseline in seizure frequency per 28 days.</p> <p>Elepsia XR, Keppra XR, Lamictal XR, Lamotrigine ER, Motpoly XR, Oxtellar XR, Qudexy XR, Topiramate ER, Topiramate SR, Trokendi XR: patient has been unable to be compliant with or tolerate twice daily dosing of the immediate release product. Additionally, if brand Elepsia XR, Keppra XR or Lamictal XR is requested, the patient has a documented intolerance to the generic product. If topiramate ER sprinkle caps are requested, the patient must have a documented intolerance to Qudexy XR.</p> <p>Lamictal ODT, Lamotrigine ODT: medical necessity for a specialty dosage form has been provided AND lamotrigine chewable tabs cannot be used. For approval of brand Lamictal ODT, the patient must have a documented intolerance to the generic equivalent.</p> <p>Libervant: The diagnosis is intermittent episodes of frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) which are distinct from the patient's usual seizure pattern AND Patient is within FDA indicated age range AND the medication is prescribed by or in consultation with a neurologist AND the patient must have a documented intolerance to diazepam rectal gel.</p> <p>Pregabalin oral solution: the patient is unable to use pregabalin capsules (i.e. swallowing disorder) AND has a documented intolerance to brand Lyrica solution.</p> <p>Spritam: medical necessity for a specialty dosage form has been provided AND patient must have a documented intolerance to levetiracetam oral solution.</p> <p>Sympazan: diagnosis or indication is adjunctive treatment of refractory epilepsy (may include different types of epilepsy) AND patient has had a documented side effect, allergy, treatment failure, inadequate response or a contraindication to at least TWO preferred anticonvulsants AND prescriber must provide a clinically compelling reason why the patient is unable to use Clobazam tablets AND Clobazam suspension.</p> <p>Tiagabine generic: The diagnosis is adjunctive therapy of focal-onset seizures and the patient has had a documented side effect, allergy, treatment failure/inadequate response or a contraindication to at least TWO preferred anticonvulsants.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
	Zarontin [®] (ethosuximide) Zonisade [™] (zonisamide) suspension Ztalmy [®] (ganaxolone) suspension <i>QTY LIMIT</i> : 36 mL/day	<p>Sabril, Vigabatrin: prescriber and patient are registered with the REMS program AND diagnosis is infantile spasms OR patient is ≥2 years old and the indication is adjunctive therapy in refractory complex partial seizures and failure of THREE other preferred anticonvulsants. For approval of Vigabatrin, the patient must have a documented side effect, allergy, contraindication or treatment failure with brand Sabril.</p> <p>Xcopri: the diagnosis is adjunctive therapy of partial-onset seizures AND the patient is ≥ 18 years of age AND the patient has had a documented side effect, allergy, treatment failure/inadequate response or a contraindication to at least TWO preferred anticonvulsants AND for reapproval, the patient must have a documented decrease from baseline in seizure frequency per 28 days.</p> <p>Ztalmy: Diagnosis or indication is for the treatment of seizures associated with cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder (CDD) confirmed by genetic testing (results must be submitted) AND patient has had a documented side effect, allergy, treatment failure, inadequate response or a contraindication to at least TWO preferred anticonvulsants AND for reapproval, the patient must have a documented decrease from baseline in seizure frequency per 28 days.</p> <p>PA Requests to Exceed QTY LIMIT for clonazepam/clonazepam ODT or Klonopin: all requests will be referred to the DVHA Medical Director for review unless the patient has been started and stabilized on the requested quantity for treatment of a seizure disorder.</p>
NASAL		
NAYZILAM [®] (midazolam) nasal spray (age ≥ 12 years) <i>QTY LIMIT</i> : 10 units/30 days VALTOCO [®] (diazepam) nasal spray (age ≥ 6 years) <i>QTY LIMIT</i> : 20 units/30 days		
RECTAL		
DIAZEPAM (compare to Diastat [®]) rectal gel		
ANTIDEPRESSANTS		
MAO INHIBITORS		
PHENELZINE SULFATE (compare to Nardil [®]) FDA maximum recommended dose = 90 mg/day TRANYLCPROMINE FDA maximum recommended dose = 60 mg/day	Emsam [®] (selegiline) <i>QTY LIMIT</i> : 1 patch/day Marplan [®] (isocarboxazid) Nardil [®] (phenylzine) FDA maximum recommended dose = 90 mg/day	<p>Marplan: patient has been started and stabilized on the requested medication (Note: samples are not considered adequate justification for stabilization). OR patient has had a documented side effect, allergy, or treatment failure to phenelzine and tranylcypromine.</p> <p>Nardil: patient has had a documented intolerance to generic equivalent product.</p> <p>Emsam: patient has had a documented side effect, allergy, or treatment failure with at least 3 antidepressants from 2 of the major antidepressant classes (Miscellaneous, SNRIs, SSRIs, and Tricyclic Antidepressants). OR patient is unable to tolerate oral medication.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

MISCELLANEOUS

<p>BUPROPION SR (compare to Wellbutrin SR®) FDA maximum recommended dose = 400mg/day</p> <p>BUPROPION XL (compare to Wellbutrin XL®) 150 mg, 300 mg FDA maximum recommended dose = 450 mg/day</p> <p>BUPROPION FDA maximum recommended dose = 450 mg/day</p> <p>MAPROTILINE FDA maximum recommended dose = 225 mg/day</p> <p>MIRTAZAPINE (compare to Remeron®) FDA maximum recommended dose = 45 mg/day</p> <p>MIRTAZAPINE RDT (compare to Remeron Sol-Tab®) FDA maximum recommended dose = 45 mg/day</p> <p>TRAZODONE HCL (formerly Desyrel®) FDA maximum recommended dose = 600 mg/day</p> <p>VILAZODONE (compare to Viibryd®) Tablet (Age ≥ 18 years) <i>QTY LIMIT:</i> 1 tablet/day FDA maximum recommended dose = 40 mg/day</p>	<p>Aplenzin® (bupropion hydrobromide) ER tablets <i>QTY LIMIT:</i> 1 tablet/day</p> <p>Auvelity™ (bupropion/dextromethorphan) <i>QTY LIMIT:</i> 2 tablets/day</p> <p>Bupropion XL 450mg (compare to Forfivo XL®) <i>QTY LIMIT:</i> 1 tablet/day FDA maximum recommended dose = 450 mg/day</p> <p>Forfivo XL® (bupropion SR 24hr) 450 mg tablet <i>QTY LIMIT:</i> 1 tablet/day FDA maximum recommended dose = 450 mg/day</p> <p>Nefazodone FDA maximum recommended dose = 600 mg/day</p> <p>Remeron® (mirtazapine) FDA maximum recommended dose = 45 mg/day</p> <p>Remeron Sol Tab® (mirtazapine RDT) FDA maximum recommended dose = 45 mg/day</p> <p>Spravato® (esketamine) nasal spray <i>QTY LIMIT:</i> not to exceed FDA recommended dose and frequency for corresponding timeframe</p> <p>Trintellix® (vortioxetine) Tablet <i>QTY LIMIT:</i> 1 tablet/day</p> <p>Wellbutrin SR® (bupropion SR) FDA maximum recommended dose = 400 mg/day</p> <p>Viibryd® (vilazodone) Tablet (Age ≥ 18 years) <i>QTY LIMIT:</i> 1 tablet/day FDA maximum recommended dose = 40 mg/day</p> <p>Wellbutrin XL® (bupropion XL) FDA maximum recommended dose = 450 mg/day</p> <p>Zulresso™ (brexanolone) intravenous solution</p> <p>Zurzuvae™ (zuranolone) capsule FDA maximum recommended dose = 50 mg/day Max approval of 14 days</p>	<p>Criteria for approval for ALL non-preferred drugs: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR The patient meets additional criteria as outlined below.</p> <p>Aplenzin, Auvelity: The patient is ≥ 18 years of age AND The patient has had a documented side effect, allergy, or inadequate response to at least 3 different antidepressants from the SSRI, SNRI and/or Miscellaneous Antidepressant categories (may be preferred or non-preferred), one of which must be bupropion.</p> <p>Bupropion XL 450mg, Forfivo XL: The patient is unable to take the equivalent dose as generic bupropion XL (150mg & 300mg) AND for approval of brand, the patient must have a documented intolerance to the generic equivalent.</p> <p>Nefazodone: The patient is ≥ 18 years of age AND The patient has had a documented side effect, allergy, or inadequate response to at least 3 different antidepressants from the SSRI, SNRI and/or Miscellaneous Antidepressant categories (may be preferred or non-preferred)</p> <p>Remeron, Remeron SolTab, Wellbutrin SR, and Wellbutrin XL: The patient has had a documented intolerance to the generic formulation of the requested medication.</p> <p>Spravato: <i>Diagnosis is treatment resistant depression:</i> the patient is ≥ 18 years of age AND medication is being used as adjunct treatment with an oral antidepressant AND the patient has a documented treatment failure (defined by at least 8 weeks of therapy) with at least 2 different antidepressants from the SSRI, SNRI, and/or Miscellaneous Antidepressant categories (may be preferred or non-preferred) AND the healthcare site and patient are enrolled in the Spravato® REMS program. Initial approval will be granted for 3 months. For re-approval after 3 months, the patient must have documented improvement in symptoms. <i>Diagnosis is Major Depressive Disorder (MDD) with Acute Suicidal Ideation or Behavior:</i> the patient is ≥ 18 years of age AND the medication is being used as adjunct treatment with an oral antidepressant AND the healthcare site and patient are enrolled in the Spravato® REMS program. Approval will be granted for 4 weeks.</p> <p>Trintellix: The patient is ≥ 18 years of age AND The diagnosis or indication is MDD AND The patient has had a documented side effect, allergy, or inadequate response (defined by at least 8 weeks of therapy) to at least 2 different antidepressants from the SSRI, SNRI, and/or Miscellaneous Antidepressant categories (may be preferred or non-preferred).</p> <p>Viibryd: Patient is ≥ 18 years of age AND The patient has had a documented intolerance to generic vilazodone.</p>
--	--	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>Zulresso: Patient is ≥ 18 years of age and ≤ 6 months postpartum AND patient has a diagnosis of postpartum depression (PPD) with documented onset of symptoms occurring in the third trimester or within 4 weeks of delivery AND the patient has a documented treatment failure (defined by at least 8 weeks of therapy) with two different oral antidepressants unless contraindicated or documentation shows that the severity of depression would place the health of the mother or infant at significant risk AND the pharmacy, patient, and healthcare facility are enrolled in the REMS program. Note: Zulresso™ will be approved as a medical benefit ONLY and will NOT be approved if billed through pharmacy point of sale.</p> <p>Zurzuvae: Patient is ≥ 18 years of age and ≤ 6 months postpartum AND patient has a diagnosis of postpartum depression (PPD) with documented onset of symptoms occurring in the third trimester or within 4 weeks of delivery AND the patient has a documented treatment failure (defined by at least 8 weeks of therapy) with two different oral antidepressants unless contraindicated or documentation shows that the severity of depression would place the health of the mother or infant at significant risk AND the patient has been instructed not to drive a motor vehicle or engage in other potentially hazardous activities requiring complete mental alertness, such as operating machinery, until at least 12 hours after administration for the duration of the 14-day treatment course.</p> <p>Note: After a 4-month lapse in use of a non-preferred agent for a mental health indication, or if there is a change in therapy, a lookback through claims information will identify the need to re-initiate therapy following the PDL and clinical criteria.</p>

SNRI		
<p>DULOXETINE (compare to Cymbalta®) capsule <i>QTY LIMIT:</i> 2 capsules/day FDA maximum recommended dose = 120 mg/day (MDD and GAD), 60 mg/day all others</p> <p>VENLAFAXINE ER capsule (compare to Effexor XR®) <i>QTY LIMIT:</i> 37.5 mg and 75 mg = 1 capsule/day FDA maximum recommended dose = 225 mg/day</p> <p>VENLAFAXINE IR tablet FDA maximum recommended dose = 225 mg/day</p>	<p>Cymbalta® (duloxetine) capsule <i>QTY LIMIT:</i> 2 capsules/day FDA maximum recommended dose = 120 mg/day (MDD and GAD), 60 mg/day all others</p> <p>Desvenlafaxine base SR <i>QTY LIMIT:</i> 50 mg tablet only = 1 tablet/day FDA maximum recommended dose = 400 mg/day</p> <p>Desvenlafaxine succinate ER (compare to Pristiq®) <i>QTY LIMIT:</i> 50 mg tablet only = 1 tablet/day FDA maximum recommended dose = 400 mg/day</p> <p>Drizalma® (duloxetine) sprinkle capsule <i>QTY LIMIT:</i> 2 capsules/day FDA maximum recommended dose = 120 mg/day (MDD and GAD), 60 mg/day all others</p> <p>Effexor XR® (venlafaxine XR) capsule <i>QTY LIMIT:</i> 37.5 mg and 75 mg = 1 capsule/day FDA maximum recommended dose = 225 mg/day</p> <p>Fetzima® (levomilnacipran ER) capsule <i>QTY LIMIT:</i> 1 capsule/day FDA maximum recommended dose = 120 mg/day</p>	<p>Criteria for approval of ALL non-preferred drugs: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR The patient meets additional criteria as outlined below.</p> <p>Venlafaxine ER tablet (generic), Effexor XR Capsule (brand), Desvenlafaxine ER succinate, Pristiq: The patient has had a documented intolerance to generic venlafaxine ER caps AND if the request is for Pristiq, the patient has a documented intolerance to the generic.</p> <p>Desvenlafaxine SR (base), Fetzima: The patient has had a documented side effect, allergy, or inadequate response to at least 2 different antidepressants AND The patient has had a documented intolerance with generic desvenlafaxine succinate ER.</p> <p>Cymbalta, Drizalma: There must be a clinically compelling reason why the dosing needs cannot be accomplished with generic duloxetine.</p> <p>Note: After a 4-month lapse in use of a non-preferred agent for a mental health indication, or if there is a change in therapy, a lookback through claims information will identify the need to re-initiate therapy following the PDL and clinical criteria.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
	Fetzima [®] (levomilnacipran ER) capsule titration pack <i>QTY LIMIT</i> : 1 pack per lifetime FDA maximum recommended dose = 120 mg/day Pristiq [®] (desvenlafaxine succinate SR) <i>QTY LIMIT</i> : 50 mg tablet only = 1 tablet/day FDA maximum recommended dose = 400 mg/day Venlafaxine ER [®] tablet <i>QTY LIMIT</i> : 37.5 mg and 75 mg = 1 tablet/day FDA maximum recommended dose = 225 mg/day	

SSRIs

CITALOPRAM (compare to Celexa [®]) tablets, solution FDA maximum recommended dose = 40 mg/day ESCITALOPRAM (compare to Lexapro [®]) tablets FDA maximum recommended dose = 20mg/day FLUOXETINE (compare to Prozac [®]) capsules, tablets, solution FDA maximum recommended dose = 80 mg/day FLUVOXAMINE FDA maximum recommended dose = 300 mg/day PAROXETINE hydrochloride tablet (compare to Paxil [®]) FDA maximum recommended dose = 60 mg/day SERTRALINE (compare to Zoloft [®]) tablet, solution FDA maximum recommended dose = 200 mg/day,	Brisdelle [®] (paroxetine mesylate) <i>QTY LIMIT</i> : 1 capsule/day Celexa [®] (citalopram) FDA maximum recommended dose = 40 mg/day Citalopram capsule <i>QTY LIMIT</i> : 1 capsule/day Escitalopram solution FDA maximum recommended dose = 20 mg/day Fluoxetine 90 mg FDA maximum recommended dose = 90 mg/week Fluvoxamine CR <i>QTY LIMIT</i> : 2 capsules/day FDA maximum recommended dose = 300 mg/day Lexapro [®] (escitalopram) <i>QTY LIMIT</i> : 5 mg and 10 mg tablets = 1.5 tabs/day FDA maximum recommended dose = 20mg/day Paroxetine mesylate (compare to Brisdelle [®]) <i>QTY LIMIT</i> : 1 capsule/day Paroxetine CR (compare to Paxil CR [®]) FDA maximum recommended dose = 75 mg/day Paxil [®] (paroxetine) FDA maximum recommended dose = 60 mg/day Paxil [®] suspension (paroxetine) FDA maximum recommended dose = 60 mg/day Paxil CR [®] (paroxetine CR) FDA maximum recommended dose = 75 mg/day Pexeva [®] (paroxetine) FDA maximum recommended dose = 60 mg/day Prozac [®] (fluoxetine) FDA maximum recommended dose = 80 mg/day Sertraline capsule 150 mg, 200 mg <i>QTY LIMIT</i> : 1 capsule/day	<p>Celexa, Fluvoxamine CR, Lexapro, Paxil tablet, Pexva, Paroxetine CR, Paxil CR, Prozac, Zoloft: The patient had a documented side effect, allergy, or treatment failure with 2 preferred SSRIs. One trial must be the generic formulation or IR formulation if CR formulation requested.</p> <p>Brisdelle, Paroxetine mesylate: The indication for use is moderate to severe vasomotor symptoms (VMS) associated with menopause. AND The patient has tried and failed generic paroxetine hydrochloride.</p> <p>Paxil suspension, Escitalopram solution: The patient has a requirement for an oral liquid dosage form. AND The patient had a documented side effect, allergy, or treatment failure with 2 preferred liquid SSRI formulations.</p> <p>Fluoxetine 90mg: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR The patient failed and is not a candidate for daily fluoxetine. AND The prescriber provides clinically compelling rationale for once-weekly dosing.</p> <p>Citalopram capsules, Sertraline capsules: Prescriber must provide a clinically compelling reason why the patient is unable to use tablets.</p> <p>Note: After a 4-month lapse in use of a non-preferred agent for a mental health indication, or if there is a change in therapy, a lookback through claims information will identify the need to re-initiate therapy following the PDL and clinical criteria.</p>
---	--	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
	Zoloft® (sertraline) <i>QTY LIMIT:</i> 25 mg and 50 mg tablets = 1.5 tabs/day FDA maximum recommended dose = 200 mg/day	
TRICYCLICS		
AMITRIPTYLINE FDA maximum recommended dose = 300 mg/day CLOMIPRAMINE DOXEPIN capsules, solution IMIPRAMINE FDA maximum recommended dose = 300 mg/day NORTRIPTYLINE (compare to Pamelor®) NORTRIPTYLINE Oral Solution	Amoxapine Anafranil® (clomipramine) Imipramine Pamoate capsules Desipramine (compare to Norpramin®) Norpramin® (desipramine) Pamelor® (nortriptyline) Protriptyline Trimipramine (compare to Surmontil®)	<p>Criteria for approval of ALL non-preferred drugs: patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR the patient meets additional criteria as outlined below.</p> <p>Imipramine Pamoate: The patient has had a documented side effect, allergy, or treatment failure to 3 preferred TCAs, one of which must be imipramine tablets.</p> <p>Desipramine: The patient has had a documented side effect, allergy, or treatment failure to nortriptyline.</p> <p>All other non-preferred agents: The patient has had a documented side effect, allergy, or treatment failure to 2 or more preferred TCAs. One trial must be the AB rated generic formulation if available</p> <p>Limitation: Chlordiazepoxide/amitriptyline and amitriptyline/perphenazine combinations are not covered. Generic agents may be prescribed separately.</p>
ANTI-DIABETICS		
ALPHA-GLUCOSIDASE INHIBITORS		
ACARBOSE	Miglitol	Miglitol: Patient must have a documented side effect, allergy or treatment failure to acarbose.
BIGUANIDES & COMBINATIONS		
<u>SINGLE AGENT</u> METFORMIN METFORMIN XR <u>COMBINATION</u> GLIPIZIDE/METFORMIN GLYBURIDE/METFORMIN	Glumetza® (metformin ER modified release) Metformin ER modified release (compare to Glumetza®) Metformin oral solution (compare to Riomet®) Metformin ER Osmotic Riomet® (metformin oral solution)	<p>Glumetza, Metformin ER mod release, Metformin ER osmotic: patient has had a documented intolerance to generic metformin XR (if product has an AB rated generic, there must have been a trial of the generic)</p> <p>Metformin oral solution, Riomet: prescriber provides documentation of medical necessity for the specialty dosage form (i.e. inability to swallow tablets, dysphagia)</p>
CD3 MONOCLONAL ANTIBODY		
All products require PA	Tziel™ (teplizumab-mzwv) vial for IV infusion	<p>Tziel:</p> <ul style="list-style-type: none"> • Patient is ≥ 8 years of age • Patient has Stage 2 Type 1 Diabetes as documented by the following: <ul style="list-style-type: none"> ○ Patient has at least 2 positive pancreatic islet cell autoantibodies (Glutamic acid decarboxylase 65 (GAD) autoantibodies, Insulin autoantibody (IAA), Insulinoma-associated antigen 2 autoantibody

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>(IA-2A), Zinc transporter 8 autoantibody (ZnT8A), or Islet cell autoantibody (ICA)</p> <ul style="list-style-type: none"> ○ Dysglycemia without overt hyperglycemia, as demonstrated by at least one of the following: Fasting plasma glucose 110-125 mg/dL, 2-hour postprandial glucose 140-199 mg/dL, or Postprandial glucose level at 30, 60, or 90 minutes > 200 mg/dL ● Patient does not have any of the following: <ul style="list-style-type: none"> ○ Lymphocyte count less than 1,000 lymphocytes/mcL ○ Hemoglobin less than 10 g/dL ○ Platelet count less than 150,000 platelets/mcL ○ Absolute neutrophil count less than 1,500 neutrophils/mcL ○ Elevated ALT or AST greater than 2 times the upper limit of normal (ULN) ○ Bilirubin greater than 1.5 times ULN ● Patient has received all age-appropriate vaccines prior to starting Tzield (Live-attenuated vaccines should be administered at least 8 weeks prior to treatment. Inactivated vaccines or mRNA vaccines should be administered at least 2 weeks prior to treatment).
DIPEPTIDYL PEPTIDASE (DPP-4) INHIBITORS		
<p><u>SINGLE AGENT</u> JANUVIA® (sitagliptin) TRADJENTA® (linagliptin)</p> <p><u>COMBINATION</u> JANUMET® (sitagliptin/metformin) JANUMET XR® (sitagliptin/metformin ER) JENTADUETO® (linagliptin/metformin) JENTADUETO® XR (linagliptin/metformin ER)</p>	<p>Alogliptan <i>QTY LIMIT</i>: 1 tab/day Zituvio™ (sitagliptin)</p> <p>Alogliptin/metformin <i>QTY LIMIT</i>: 1 tab/day Alogliptin/pioglitazone <i>QTY LIMIT</i>: 1 tab/day Kombiglyze XR® (saxagliptin/metformin ER) <i>QTY LIMIT</i>: 1 tab/day</p>	<p>Alogliptan, Zituvio: patient has had a documented side effect, allergy OR treatment failure with two preferred DPP-4 agents.</p> <p>Alogliptin/metformin, Kombiglyze XR: patient has had a documented side effect, allergy OR treatment failure with at least one preferred DPP-4 combination agent.</p> <p>Alogliptin/pioglitazone: patient has had a documented side effect, allergy, OR treatment failure with at least one preferred DPP-4 agent used in combination with pioglitazone.</p>
HYPOGLYCEMIA TREATMENTS		
<p>BAQSIMI® (glucagon nasal powder) 3mg <i>QTY LIMIT</i>: 2 devices/28 days GLUCAGEN® HYPOKIT® (glucagon for injection) 1mg ZEGALOGUE® (dasiglucagon SC injection) 0.6 mg <i>QTY LIMIT</i>: 2 prefilled syringes or auto-injectors/28 days</p>	<p>Glucagon emergency kit Gvoke™ (glucagon SC injection) prefilled syringe, auto-injector 0.5mg, 1mg</p>	<p>Glucagon Emergency Kit, Gvoke: Patient has recurrent episodes of symptomatic or severe hypoglycemia (<55 mg/dL) requiring the assistance of another individual AND the preferred formulations would not be suitable alternatives.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

INSULINS

RAPID-ACTING INJECTABLE

FIASP® (insulin aspart)
HUMALOG® (insulin lispro)
INSULIN ASPART (compare to Novolog®)
INSULIN LISPRO (compare to Humalog®)

SHORT-ACTING INJECTABLE

HUMULIN R® U-500

INTERMEDIATE-ACTING INJECTABLE

All products require PA

LONG-ACTING ANALOGS INJECTABLE

LANTUS® (insulin glargine)
TOUJEO® (insulin glargine)
TOUJEO® MAX (insulin glargine)

MIXED INSULINS INJECTABLE

HUMULIN 70/30® (NPH/Regular)
NOVOLOG MIX 70/30® (Protamine/Aspart)
HUMALOG MIX 50/50® (Protamine/Lispro)
HUMALOG MIX 75/25® (Protamine/Lispro)
INSULIN ASPART PROTAMINE/ASPART 70/30
(compare to Novolog Mix 70/30®)

Admelog® (insulin lispro)
Afrezza® Inhaled (insulin human)
Apidra® (insulin glulisine)
Humalog (insulin lispro) Kwikpen U-200
Lyumjev® (insulin lispro-aabc)
Novolog® (insulin aspart)

Humulin R® (Regular) U-100
Novolin R® (Regular) U-100
Humulin N® (NPH)
Novolin N® (NPH)

Basaglar® (insulin glargine)
Insulin Degludec (compare to Tresiba®)
Insulin Glargine (compare to Lantus®)
Insulin Glargine-yfgn (compare to Semglee®)
Rezvoglar™ (insulin glargine-aglr)
Semglee® (insulin glargine-yfgn)
Tresiba® (insulin degludec)

Novolin 70/30® (NPH/Regular)

Non-preferred rapid-acting insulin: Patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR patient has had a documented side effect, allergy OR treatment failure to two preferred formulations of rapid-acting insulin.

Additional criteria for Lyumjev: Patient has had a documented side effect, allergy, or treatment failure to Fiasp.

Humulin N/R, Novolin N/R: patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR patient has a documented treatment failure to at least one preferred agent.

Non-preferred long-acting insulin products: Patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR had a documented side effect, allergy or treatment failure to one preferred long-acting insulin products.

Insulin Degludec: Patient has had a documented side effect, allergy or treatment failure to Tresiba.

Novolin 70/30: patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR patient has had a documented side effect, allergy or treatment failure to two preferred mixed insulin formulations.

AFREZZA INHALED INSULIN:

- Baseline PFT with FEV1 ≥ 70 % predicted
- Patient does not have underlying lung disease (Asthma, COPD)
- Patient is a non-smoker or has stopped smoking more than six months prior to starting Afrezza
- Patient is currently using a long-acting insulin
- Patient has failed to achieve HbA1c goal (defined as ≤ 7%) on a short-acting insulin in combination with a long-acting insulin
- Initial approval is for 3 months and improved glycemic control must be documented for further approvals

MEGLITINIDES

NATEGLINIDE
REPAGLINIDE

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

PEPTIDE HORMONES: GLP-1 RECEPTOR AGONISTS

<p><i>Preferred After Clinical Criteria Are Met</i></p> <p>SINGLE AGENTS Rybelsus® (semaglutide) tablets <i>QTY LIMIT: 1 tablet/day</i></p> <p>TRULICITY® (dulaglutide) <i>QTY LIMIT: 12 pens/84 days</i></p> <p>VICTOZA® (liraglutide) <i>QTY LIMIT: 9 pens/90 days</i></p> <p>COMBINATION AGENTS All products require PA</p> <p>AMYLINOMIMETICS All products require PA</p>	<p>Adlyxin® (lixisenatide) Bydureon® BCise™ (exenatide extended-release) <i>QTY LIMIT: 12 pens/84 days</i></p> <p>Byetta® (exenatide) <i>QTY LIMIT: 3 pens/90 days</i></p> <p>Mounjaro™ (tirzepatide) <i>QTY LIMIT: 4 pens/28 days</i></p> <p>OZEMPIC® (semaglutide) <i>QTY LIMIT: 9mL/84 days</i></p> <p>Soliqua® (insulin glargine/lixisenatide) <i>QTY LIMIT: 3 pens/25 days</i></p> <p>Xultophy® (insulin degludec/liraglutide)</p> <p>Symlin® (pramlintide)</p>	<p>Clinical criteria for all drugs: patient has a diagnosis of Type 2 Diabetes Mellitus</p> <p>Ozempic: Patient has a documented side effect, allergy, contraindication, or treatment failure with Trulicity. Treatment failure is defined as <1% reductio in HbA1c after 12 weeks at the maximally tolerated dose.</p> <p>Additional criteria for Adlyxin/Byetta/Bydureon BCise, Mounjaro: patient has a documented side effect, allergy, contraindication, or treatment failure with one preferred GLP-1 Receptor Agonist AND Ozempic. Treatment failure is defined as <1% reduction in HbA1c after 12 weeks at the maximally tolerated dose.</p> <p>Soliqua/Xultophy: patient has a documented side effect, allergy, contraindication, or treatment failure with at least one preferred GLP-1 Receptor Agonist used in combination with Lantus. Treatment failure is defined as < 1% reduction in HbA1c after 12 weeks at the maximally tolerated dose.</p> <p>Symlin: patient is at least 18 years of age AND patient is on insulin.</p>
---	---	--

SODIUM-GLUCOSE CO-TRANSPORTER 2 (SGLT2) INHIBITORS AND COMBINATIONS

<p>SINGLE AGENTS FARXIGA® (dapagliflozin) JARDIANCE (empagliflozin)</p> <p>COMBINATIONS AGENTS SYNJARDY® (empagliflozin/metformin) SYNDJARDY® XR (empagliflozin/metformin ER) <i>QTY LIMIT: 1 tab/day</i></p>	<p>Dapagliflozin (compare to Farxiga) Invokana® (canagliflozin) Steglatro® (ertugliflozin) <i>QTY LIMIT: 1 tab/day</i></p> <p>Glyxambi® (empagliflozin/ linagliptin) <i>QTY LIMIT: 1 tab/day</i></p> <p>Invokamet® (canagliflozin/metformin) Invokamet® XR (canagliflozin/metformin ER) Qtern® (dapagliflozin/saxagliptin) Segluromet® (ertugliflozin/metformin) <i>QTY LIMIT: 2 tabs/day</i> Steglujan® (ertugliflozin/sitagliptin) <i>QTY LIMIT: 1 tab/day</i></p> <p>Trijardy® XR (empagliflozin/linagliptin/metformin ER) Xigduo XR® (dapagliflozin/metformin ER) <i>QTY LIMIT: 5/1000 mg = 2 tabs/day, all other strengths = 1 tab/day</i></p>	<p>Dapagliflozin, Invokana, Steglatro: Patient has a documented side effect, allergy, or contraindication to two preferred SGLT2 inhibitors.</p> <p>Invokamet, Invokamet XR, Segluromet, Xigduo XR: The patient has documentation of a failure of therapy with a preferred SGLT2 inhibitor used in combination with metformin/metformin XR.</p> <p>Glyxambi/Qtern/Steglujan: The patient has documentation of a failure of therapy with the combination of a preferred SGLT2 inhibitor plus a preferred DPP-4 inhibitor</p> <p>Trijardy XR: patient has documentation of a failure of therapy with a preferred SGLT2 inhibitor, a preferred DDP-4 inhibitor and metformin/metformin XR used in combination.</p>
--	--	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
SULFONYLUREAS 2ND GENERATION		
GLIMEPIRIDE (compare to Amaryl [®]) GLIPIZIDE (compare to Glucotrol [®]) GLIPIZIDE ER (compare to Glucotrol XL [®]) GLYBURIDE GLYBURIDE MICRONIZED	Amaryl [®] (glimepiride) Glucotrol XL [®] (glipizide ER) Glynase [®] (glyburide micronized)	Criteria for Approval: Patient must have a documented side effect, allergy or treatment failure to two preferred sulfonylureas. If a product has an AB rated generic, one trial must be the generic.
THIAZOLIDINEDIONES & COMBINATIONS		
PIOGLITAZONE (compare to Actos [®]) <u>COMBINATION</u> All products require PA	Actos [®] (pioglitazone) Actoplus Met [®] (pioglitazone/metformin) Duetact [®] (pioglitazone/glimepiride) <i>QTY LIMIT:</i> 1 tablet/day Pioglitazone/Glimepiride (compare to Duetact [®]) <i>QTY LIMIT:</i> 1 tablet/day Pioglitazone/Metformin (compare to Actoplus Met)	Actos: the patient has a documented intolerance to the generic equivalent. Actoplus Met, Duetact, Pioglitazone/Metformin, Pioglitazone/Glimepiride: patient is unable to take as the individual separate agents AND if the request is for Actoplus Met or Duetact, the patient has had a documented intolerance to the generic equivalent.
ANTI-EMETICS		
5HT3 ANTAGONISTS: Length of Authorization: 6 months for chemotherapy or radiotherapy; 3 months for hyperemesis gravidarum, 1 time for prevention of post-op nausea/vomiting: see clinical criteria. Monthly quantity limits apply, PA required to exceed.		
GRANISETRON injection ONDANSETRON injection ONDANSETRON tablet ONDANSETRON ODT ONDANSETRON oral solution 4mg/5mL PALONOSETRON injection	Akynzeo [®] (nutupitant/palonosetron) Granisetron 1 mg tablets <i>QTY LIMIT:</i> 6 tabs/28 days Sancuso [®] 3.1 mg/24 hr transdermal patch (granisetron) <i>QTY LIMIT:</i> 4 patches/28 days Sustol [®] (granisetron) injection 10 mg/0.4ml <i>QTY LIMIT:</i> 4 injections/28 days	Akynzeo: Has a diagnosis of nausea and vomiting associated with cancer chemotherapy AND patient has a documented side effect, allergy, or treatment failure of a regimen consisting of a 5-HT3 antagonist, an NK1 antagonist, and dexamethasone. Granisetron tablets: The patient has had a documented side effect, allergy, or treatment failure to generic ondansetron. Sancuso: patient has a diagnosis of nausea and vomiting associated with cancer chemotherapy. AND prescriber provides documentation of medical necessity for the transdermal formulation. OR patient has had a documented side effect, allergy, or treatment failure with generic ondansetron. Sustol: Patient has a diagnosis of nausea and vomiting associated with cancer chemotherapy or radiotherapy AND prescriber provides documentation of medical necessity for the specialty dosage form (i.e. inability to swallow

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>tablets, dysphagia) AND the patient has a documented side effect, allergy, or treatment failure with Ondansetron injection and Sancuso transdermal.</p> <p>CRITERIA FOR APPROVAL to Exceed QTY LIMIT: Granisetron: For nausea and vomiting associated with chemotherapy, 2 tablets for each day of chemotherapy and 2 tablets for 2 days after completion of chemotherapy may be approved. OR For nausea and vomiting associated with radiation therapy, 2 tablets for each day of radiation may be approved. Sancuso: For nausea and vomiting associated with chemotherapy, 1 patch for each chemotherapy cycle may be approved.</p>
MISCELLANEOUS (PREGNANCY)		
<p>DICLEGIS® (10 mg doxylamine succinate and 10 mg pyridoxine hydrochloride) DR tablet <i>QTY LIMIT:</i> 4 tablets/day</p>	<p>Bonjesta® (20 mg doxylamine succinate and 20 mg pyridoxine hydrochloride ER tablet) <i>QTY LIMIT:</i> 2 tablets/day Doxylamine succinate/pyridoxine hydrochloride DR tablet (compare to Diclegis®) <i>QTY LIMIT:</i> 4 tablets/day</p>	<p>Bonjesta, Doxylamine/Pyridoxone: patient has a documented intolerance to Diclegis.</p>
NKI ANTAGONISTS		
<p>APONVIE® (aprepitant) injection CINVANTI® (aprepitant) injection EMEND® (fosaprepitant) injection</p> <p><u>Preferred After Clinical Criteria Are Met</u> EMEND® (aprepitant) 80 mg <i>QTY LIMIT:</i> 2 caps/28 days EMEND® (aprepitant) Tri-fold Pack <i>QTY LIMIT:</i> 1 pack/28 days</p>	<p>Aprepitant (compare to Emend®) 40 mg <i>QTY LIMIT:</i> 1 cap/28 days Aprepitant (compare to Emend®) 80 mg <i>QTY LIMIT:</i> 2 caps/28 days Aprepitant (compare to Emend®) 125 mg <i>QTY LIMIT:</i> 1 cap/28 days Emend® (aprepitant) oral suspension</p>	<p>Aprepitant, Emend (aprepitant): medication will be prescribed by an oncology practitioner. AND patient requires prevention of nausea and vomiting associated with moderate to highly emetogenic cancer chemotherapy. AND The requested quantity does not exceed one 125 mg and two 80 mg capsules OR one Tri-Fold Pack per course of chemotherapy. Patients with multiple courses of chemotherapy per 28 days will be approved quantities sufficient for the number of courses of chemotherapy. For approval of generic aprepitant, the patient must have a documented intolerance to brand Emend.</p> <p>Emend oral suspension: medication will be prescribed by an oncology practitioner AND patient requires prevention of nausea and vomiting associated with moderate to highly emetogenic cancer chemotherapy AND patient has a documented medical necessity for the specialty dosage form (e.g. swallowing disorder)</p>
THC DERIVATIVES		
<p>All products require PA</p>	<p>Dronabinol (compare to Marinol®) Marinol® (dronabinol)</p>	<p>Dronabinol/Marinol: patient has a diagnosis of chemotherapy-induced nausea/vomiting AND patient has had a documented side effect, allergy, or</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		treatment failure to at least 2 antiemetic agents, of which, one must be a preferred 5HT3 receptor antagonist. If the request is for Marinol, the patient must additionally have a documented intolerance to generic dronabinol. OR patient has a diagnosis of HIV/AIDS associated anorexia. AND patient has had an inadequate response, adverse reaction, or contraindication to meggestrol acetate. If the request is for Marinol, the patient must additionally have a documented intolerance to generic dronabinol.
ANTI-HYPERTENSIVES		
ACE INHIBITORS		
BENAZEPRIL (compare to Lotensin®) ENALAPRIL (compare to Vasotec®) tablet ENALAPRIL oral solution (age ≤ 12 years old) FOSINOPRIL LISINOPRIL (compare to Zestril®) QUINAPRIL (compare to Accupril®) RAMIPRIL (compare to Altace®) TRANDOLAPRIL	Accupril® (quinapril) Altace® (Ramipril) Captopril Enalapril oral solution (age > 12 years old) Epaned® (enalapril) oral solution Lotensin® (benazepril) Moexepiril Perindopril Qbrelis® (Lisinopril) 1mg/ml solution Vasotec® (enalapril) Zestril® (lisinopril)	<p>Enalapril (Patients > 12 years old), Epaned Oral Solution: patient has a requirement for an oral liquid dosage form (i.e. swallowing disorder) AND for approval of Epaned, the patient must have a documented intolerance to the generic equivalent.</p> <p>Qbrelis Oral Solution: patient has a requirement for an oral liquid dosage form (i.e. swallowing disorder, inability to take oral medications) AND has a side effect, allergy, or treatment failure to Epaned oral solution.</p> <p>Other ACE Inhibitors: patient has had a documented side effect, allergy, or treatment failure to all available preferred generic ACEI. If a medication has an AB rated generic, there must have been a trial of the generic formulation.</p>
ACE INHIBITOR W/ HYDROCHLOROTHIAZIDE		
BENAZEPRIL/HYDROCHLOROTHIAZIDE (compare to Lotensin HCT®) ENALAPRIL/HYDROCHLOROTHIAZIDE (compare to Vaseretic®) FOSINOPRIL/HYDROCHLOROTHIAZIDE LISINOPRIL/HYDROCHLOROTHIAZIDE (compare to Zestoretic®) QUINAPRIL/HYDROCHLOROTHIAZIDE (compare to Accuretic®)	Accuretic® (quinapril/HCTZ) Lotensin HCT® (benazepril/HCTZ) Vaseretic® (enalapril/HCTZ) Zestoretic® (lisinopril/HCTZ)	<p>ACE Inhibitor/Hydrochlorothiazide combinations: patient has had a documented side effect, allergy, or treatment failure to all available preferred generic ACEI/Hydrochlorothiazide combination. If a medication has an AB rated generic, there must have been a trial of the generic formulation.</p>
ACE INHIBITOR W/CALCIUM CHANNEL BLOCKER		
AMLODIPINE/BENAZEPRIL (compare to Lotrel®)	Lotrel® amlodipine/(benazepril) Trandolapril/Verapamil ER	<p>Lotrel: The patient has had a documented side effect, allergy, or treatment failure to the generic formulation.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		Trandolapril/Verapamil ER: The patient has had a documented side effect, allergy, or treatment failure to amlodipine/benazepril AND the patient is unable to take as the individual separate agents.
ANGIOTENSIN RECEPTOR BLOCKERS (ARBs)		
CANDESARTAN IRBESARTAN (compare to Avapro®) LOSARTAN (compare to Cozaar®) OLMESARTAN (compare to Benicar®) TELMISARTAN (compare to Micardis®) VALSARTAN (compare to Diovan®)	Avapro® (irbesartan) Benicar® (olmesartan) Cozaar® (losartan) Diovan® (valsartan) Edarbi® (azilsartan) Tablet <i>QTY LIMIT:</i> 1 tablet/day Micardis® (telmisartan)	Avapro, Benicar, Cozaar, Diovan, Edarbi, and Micardis: Patient has had a documented side effect, allergy, or treatment failure with TWO preferred Angiotensin Receptor Blocker (ARB) or ARB combinations. AND If brand name product with generic available, the patient has had a documented intolerance with the generic product.
ANGIOTENSIN RECEPTOR BLOCKER/DIURETIC COMBINATIONS		
IRBESARTAN/HYDROCHLOROTHIAZIDE (compare to Avalide®) LOSARTAN/HYDROCHLOROTHIAZIDE (compare to Hyzaar®) OLMESARTAN/HYDOCHLOROTHIAZIDE (compare to Benicar HCT®) TELMISARTAN/HYDROCHLOROTHIAZIDE (compare to Micardis HCT®) VALSARTAN/HYDROCHLOROTHIAZIDE (compare to Diovan HCT®)	Avalide® (irbesartan/hydrochlorothiazide) Benicar HCT® (olmesartan/hydrochlorothiazide) Candesartan/hydrochlorothiazide Diovan HCT® (valsartan/hydrochlorothiazide) Edarbyclor® (azilsartan/chlorthalidone) Tablet <i>QTY LIMIT:</i> 1 tablet/day Hyzaar® (losartan/hydrochlorothiazide) Micardis HCT® (telmisartan/hydrochlorothiazide)	Avalide, Benicar HCT, Candesartan/HCTZ, Diovan HCT, Edarbyclor, Hyzaar, Micardis HCT and Telmisartan/HCTZ: patient has had a documented side effect, allergy, or treatment failure with a preferred ARB/Hydrochlorothiazide combination AND If brand name product with generic available, the patient has had a documented intolerance with the generic product.
ANGIOTENSIN RECEPTOR BLOCKER/CALCIUM CHANNEL BLOCK COMBINATIONS		
OLMESARTAN/AMLODIPINE (compare to Azor®) VALSARTAN/AMLODIPINE (compare to Exforge®) <i>QTY LIMIT:</i> 1 tablet/day	Azor® (olmesartan/amlodipine) <i>QTY LIMIT:</i> 1 tablet/day Amlodipine/telmisartan <i>QTY LIMIT:</i> 1 tablet/day Exforge® (valsartan/amlodipine) <i>QTY LIMIT:</i> 1 tablet/day	Azor, Amlodipine/Telmisartan, Exforge, Olmesartan/amlodipine: The patient has had a documented side effect, allergy, or treatment failure to a preferred ARB/CCB combination product AND if brand name product with generic available, the patient has had a documented intolerance with the generic equivalent.
ANGIOTENSIN RECEPTOR BLOCKER/CALCIUM CHANNEL BLOCKER/HCTZ COMBO		
VALSARTAN/AMLODIPINE/HCTZ (compare to Exforge HCT®) <i>QTY LIMIT:</i> 1 tablet/day	Exforge HCT® (amlodipine/valsartan/hydrochlorothiazide) <i>QTY LIMIT:</i> 1 tablet/day	Exforge HCT, Olmesartan/amlodipine/HCTZ, Tribenzor: patient has had a documented side effect, allergy, or treatment failure to Valsartan/amlodipine/HCTZ.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
	<p>Olmesartan/amlodipine/hydrochlorothiazide (compare to Tribenzor®) <i>QTY LIMIT</i>: 1 tablet/day</p> <p>Tribenzor® (amlodipine/olmesartan/hydrochlorothiazide) <i>QTY LIMIT</i>: 1 tablet/day</p>	
BETA BLOCKERS		
<p><u>SINGLE AGENT</u></p> <p>ACEBUTOLOL</p> <p>ATENOLOL (compare to Tenormin®)</p> <p>BISOPROLOL FUMARATE</p> <p>BYSTOLIC® (nebivolol)</p> <p>CARVEDILOL (compare to Coreg®)</p> <p>LABELALOL</p> <p>METOPROLOL TARTRATE (compare to Lopressor®)</p> <p>METOPROLOL SUCCINATE XL (compare to Toprol XL®)</p> <p>NADOLOL</p> <p>NEBIVOLOL (compare to Bystolic®)</p> <p>PINDOLOL</p> <p>PROPRANOLOL</p> <p>PROPRANOLOL ER (compare to Inderal LA®)</p> <p>SOTALOL (compare to Betapace®, Betapace AF®)</p> <p><u>BETA-BLOCKER/DIURETIC COMBINATION</u></p> <p>ATENOLOL/CHLORTHALIDONE (compare to Tenoretic®)</p> <p>BISOPROLOL/HYDROCHLOROTHIAZIDE (compare to Ziac®)</p> <p>METOPROLOL/HYDROCHLOROTHIAZIDE</p>	<p>Betapace® (sotalol)</p> <p>Betapace AF® (sotalol)</p> <p>Betaxolol</p> <p>Carvedilol CR (compare to Coreg®) <i>QTY LIMIT</i>: 1 tablet/day</p> <p>Coreg® (carvedilol)</p> <p>Coreg CR® (carvedilol CR) <i>QTY LIMIT</i>: 1 tablet/day</p> <p>Corgard® (nadolol)</p> <p>Hemangeol® oral solution (propranolol)</p> <p>Inderal LA® (propranolol ER)</p> <p>Inderal XL® (propranolol SR)</p> <p>Innopran XL® (propranolol SR)</p> <p>Kaspargo Sprinkle™ (metoprolol succinate XL)</p> <p>Lopressor® (metoprolol tartrate)</p> <p>Sorine® (sotalol)</p> <p>Tenormin® (atenolol)</p> <p>Timolol</p> <p>Toprol XL® (metoprolol succinate XL)</p> <p>Nadolol/bendroflumethiazide</p> <p>Tenoretic® (atenolol/chlorthalidone)</p> <p>Ziac® (bisoprolol/HCTZ)</p>	<p>Non-preferred drugs (except as noted below) patient has had a documented side effect, allergy, or treatment failure to at least three preferred drugs. (If a medication has an AB rated generic, one trial must be the generic formulation.)</p> <p>Carvedilol CR, Coreg CR: <i>Indication: Heart Failure:</i> patient has been started and stabilized on the medication. (Note: Samples are not considered adequate justification for stabilization.) OR patient has had a documented side effect, allergy, or treatment failure to metoprolol SR or bisoprolol. AND patient has been unable to be compliant with or tolerate twice daily dosing of carvedilol IR.</p> <p><i>Indication: Hypertension:</i> patient has been started and stabilized on the medication. (Note: Samples are not considered adequate justification for stabilization.) OR patient has had a documented side effect, allergy, or treatment failure to 3 (three) preferred anti-hypertensive beta-blockers.</p> <p>Hemangeol: indication for use is the treatment of proliferating infantile hemangioma and the patient is initiating treatment at ages 5 weeks to 5 months AND patient has had a documented side effect, allergy, or treatment failure with propranolol generic solution.</p> <p>Kaspargo: patient is unable to take a solid oral dosage form and has a treatment failure with an immediate release oral solution or crushed tablets.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

MAJOR ADVERSE CARDIOVASCULAR EVENTS (MACE) REDUCTION

<p>All products require PA</p>	<p>Wegovy (semaglutide)</p>	<p>Wegovy:</p> <ul style="list-style-type: none"> • Patient has BMI > 27 kg/m², and is not being used for weight loss only • Patient has history of at least one of the following: <ul style="list-style-type: none"> ○ Stroke ○ Myocardial Infarction ○ Symptomatic peripheral arterial disease • Patient does not have diagnosis of diabetes, end stage renal disease/dialysis, or NYHA class IV heart failure • Patient has received counseling on chronic weight management (increased physical activity and a reduce calorie diet) and will continue to follow a treatment plan • Initial approval will be for 6 months, for reapproval: Patient must continue to follow a reduced calorie diet and increased physical activity plan AND patient has shown a documented weight loss of > 5% of baseline body weight OR continued to maintain initial 5% weight loss.
--------------------------------	-----------------------------	--

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

CALCIUM CHANNEL BLOCKERS		
<p><u>SINGLE AGENT</u> <u>DIHYDROPYRIDINES</u> AMLODIPINE (compare to Norvasc[®]) FELODIPINE ER NIFEDIPINE IR (compare to Procardia[®]) NIFEDIPINE SR osmotic (compare to Procardia[®] XL) NIFEDIPINE SR (compare to Adalat[®] CC)</p> <p><u>NON-DIHYDROPYRIDINES</u> CARTIA[®] XT (diltiazem SR, compare to Cardizem[®] CD) DILT-XR[®] (diltiazem SR) DILTIAZEM (compare to Cardizem[®]) DILTIAZEM ER 24-hour capsules (compare to Tiazac[®]) DILTIAZEM SR 24-hour capsules (compare to Cardizem[®] CD) DILTIAZEM SR 24-hour tablets TAZTIA[®] XT (diltiazem ER, compare to Tiazac[®]) VERAPAMIL (compare to Calan[®]) VERAPAMIL CR (compare to Calan SR[®]) VERAPAMIL SR 120 mg, 180 mg, 240 mg, and 360 mg (compare to Verelan[®])</p> <p>Note: Please refer to the Anti-Hypertensives: Angiotensin Receptor Blockers (ARBs) PDL category for ARB/CCB combination therapies</p>	<p>Isradipine Katerzia[®] (amlodipine) oral suspension Levamlodipine Nicardipine Nimodipine Norliqva[®] (amlodipine) oral solution Nisoldipine ER (compare to Sular[®]) Norvasc[®] (amlodipine) Nymalize[®] (nimodipine) Oral Solution Procardia[®] (nifedipine IR) Procardia XL[®] (nifedipine SR osmotic) Sular[®] (nisoldipine)</p> <p>Calan[®] SR (verapamil CR) Cardizem[®] (diltiazem) Cardizem[®] CD (diltiazem SR) Cardizem[®] LA (diltiazem SR) Diltiazem ER 12-hour capsules Diltiazem ER/Matzin LA (compare to Cardizem[®] LA) Tiazac[®] (diltiazem ER) Verapamil SR 100 mg, 200 mg, 300mg (compare to Verelan PM[®]) Verelan[®] (verapamil SR 120 mg, 180 mg, 240 mg and 360 mg) Verelan[®] PM (100 mg, 200 mg and 300 mg)</p>	<p>Criteria for approval (except as noted below:) patient has had a documented side effect, allergy, or treatment failure to at least three preferred drugs. (If a medication has an AB rated generic, one trial must be the generic formulation.)</p> <p>Katerzia: patient has a medical necessity for a specialty dosage form (i.e. dysphagia, swallowing disorder).</p> <p>Norliqva, Nymalize: patient has a medical necessity for a specialty dosage form (i.e. dysphagia, swallowing disorder) and the patient has a had a documented side effect, allergy, or treatment failure to Katerzia.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
CENTRAL ALPHA AGONISTS		
<p><u>ORAL TABLETS</u> CLONIDINE IR Tablets (compare to Catapres®) GUANFACINE IR Tablets (compare to Tenex®)</p> <p><u>TRANSDERMAL</u> CLONIDINE Transdermal Patch <i>QTY LIMIT</i>: 1 patch/7 days</p>	Methyldopa Tablets	Methyldopa: The patient has a documented side effect, allergy, or contraindication to two preferred central alpha agonists.
GANGLIONIC BLOCKERS		
All products require PA	Vecamyl® (mecamylamine) tablet	Vecamyl tabs: Patient has a diagnosis of moderately severe or severe hypertension AND patient has tried and failed, intolerant to, or contraindicated to at least THREE different antihypertension therapies of different mechanism of actions.
RENIN INHIBITOR		
All products require PA	<p><u>SINGLE AGENT</u> Aliskiren (compare to Tekturna®) <i>QTY LIMIT</i>: 1 tablet/day Tekturna® (aliskiren) <i>QTY LIMIT</i>: 1 tablet/day</p> <p><u>COMBINATIONS</u> Tekturna HCT® (aliskiren/hydrochlorothiazide) <i>QTY LIMIT</i>: 1 tablet/day</p>	<p>Aliskiren, Tekturna: patient is NOT a diabetic who will continue on therapy with an ACEI or ARB AND patient has a diagnosis of hypertension. AND patient has had a documented side effect, allergy, or treatment failure with an angiotensin Receptor Blocker (ARB).</p> <p>Tekturna HCT: the patient must meet criteria as listed above for Tekturna and is unable to use the individual separate agents.</p>
ANTI-INFECTIVES ANTIBIOTICS		
AMINOGLYCOSIDES		
NEOMYCIN SULFATE	Arikayce® (amikacin inhalation suspension) <i>QTY LIMIT</i> : 28 vials (235.2 mL)/28 days	Arikayce: Patient is ≥ 18 years of age AND indication for use is treatment of <i>Mycobacterium avium complex</i> (MAC) lung disease AND patient has not achieved negative sputum cultures after a minimum of 6 consecutive months of a multidrug background regimen therapy (e.g. macrolide, rifampin, & ethambutol) within the past 12 months. Note: Initial approval will be granted for 6 months. For re-approval, the patient must have documentation of clinical improvement AND 3 consecutive monthly negative sputum cultures.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
CEPHALOSPORINS 1ST GENERATION		
<p><u>CAPSULES/TABLETS</u> CEFADROXIL capsules CEPHALEXIN capsules</p> <p><u>SUSPENSION</u> CEFADROXIL suspension CEPHALEXIN suspension</p> <p>IV drugs are not managed at this time</p>	<p>Cefadroxil tablets Cephalexin tablets</p>	<p>Cephadroxil tabs: patient has had a documented intolerance to cefadroxil generic capsules.</p> <p>Cephalexin Tabs: patient has had a documented intolerance to cephalexin generic capsules.</p>
CEPHALOSPORINS 2ND GENERATION		
<p><u>CAPSULES/TABLETS</u> CEFACTOR capsule CEFPROZIL tablet CEFUROXIME tablet</p> <p><u>SUSPENSION</u> CEFPROZIL suspension</p> <p>IV drugs are not managed at this time</p>	<p>Cefaclor[®] ER tablet</p>	<p>Cefaclor ER Tabs: patient has had a documented intolerance to cefaclor capsules.</p>
CEPHALOSPORINS 3RD GENERATION		
<p><u>CAPSULES/TABLETS</u> CEFDINIR CAPSULE CEFPODOXIME TABLET</p> <p><u>SUSPENSION</u> CEFDINIR suspension</p> <p>IV drugs are not managed at this time</p>	<p>Cefixime capsule</p> <p>Cefixime suspension Cefpodoxime proxetil suspension</p>	<p>Cefixime capsule: patient is completing a course of therapy which was initiated in the hospital. OR patient has had a documented side effect or treatment failure to cefdinir or cefpodoxime.</p> <p>Cefpodoxime Proxetil Susp, Cefixime Susp: patient is completing a course of therapy which was initiated in the hospital. OR patient has had a documented side effect or treatment failure to cefdinir suspension.</p>
CLINDAMYCIN DERIVATIVES		
<p>CLINDAMYCIN (compare to Cleocin[®]) capsules CLINDAMYCIN (compare to Cleocin[®]) oral Solution</p>	<p>Cleocin (clindamycin) Capsules Cleocin[®] Ped (clindamycin) oral solution</p>	<p>Cleocin: the patient has a documented intolerance to the generic equivalent.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
NITROIMIDAZOLE ANTIMICROBIAL		
METRONIDAZOLE	Likmez™ (metronidazole oral suspension)	Likmez: patient has a medical necessity for a non-solid oral dosage form.
OXAZOLIDINONES		
<p>LINEZOLID (compare to Zyvox®) tablets <i>QTY LIMIT:</i> 56 tablets per 28 days</p> <p>IV form of this medication not managed at this time</p>	<p>Linezolid (compare to Zyvox®) suspension <i>QTY LIMIT:</i> 60 ml/day, maximum 28 days supply</p> <p>Sivextro® (tedizolid) <i>QTY LIMIT:</i> 1 tab/day</p> <p>Zyvox® (linezolid) tablets <i>QTY LIMIT:</i> 56 tablets per 28 days</p> <p>Zyvox® (linezolid) suspension <i>QTY LIMIT:</i> 60 ml/day, maximum 28 days supply</p>	<p>Linezolid suspension, Zyvox suspension: The patient must have medical necessity for a liquid formulation (i.e. swallowing disorder). AND if the request is for generic linezolid suspension, the patient has a documented intolerance to brand Zyvox suspension.</p> <p>Sivextro: Patient has been started on Sivextro in the hospital and will be finishing the course of therapy in an outpatient setting OR patient has a documented blood, tissue, sputum, or urine culture that is positive for Vancomycin-Resistant Enterococcus (VRE) species. OR patient has a documented blood, sputum, tissue, or urine culture that is positive for Methicillin Resistant Staphylococcus species AND patient has had a documented treatment failure with linezolid, trimethoprim/sulfamethoxazole, clindamycin, doxycycline, or minocycline OR there is a clinically valid reason that the patient cannot be treated with one of those agents.</p> <p>Zyvox tablets: The patient has a documented intolerance to generic linezolid tablets.</p>
PENICILLINS (ORAL)		
<p><u>SINGLE ENTITY AGENTS</u></p> <p><u>NATURAL PENICILLINS</u></p> <p>PENICILLIN V POTASSIUM tablets, oral solution</p> <p><u>PENICILLINASE-RESISTANT PENICILLINS</u></p> <p>DICLOXACILLIN Capsules</p> <p><u>AMINOPENICILLINS</u></p> <p>AMOXICILLIN capsules, tablets, chewable tablets, suspension</p> <p>AMPICILLIN capsules, suspension</p> <p><u>COMBINATION PRODUCTS</u></p> <p>AMOXICILLIN/CLAVULANATE tablets, chewable tablets, suspension</p>	<p>Amoxicillin/clavulanate ER tablets</p>	<p>Amoxicillin/Clavulanate ER: prescriber must provide a clinically valid reason for the use of the requested medication.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
QUINOLONES		
<p>CIPROFLOXACIN (compare to Cipro®) tabs CIPRO® (ciprofloxacin) oral suspension LEVOFLOXACIN (compare to Levaquin®) tabs, solution MOXIFLOXACIN tabs</p> <p>IV drugs are not managed at this time</p>	<p>Baxdela™ (delafloxacin) Cipro® (ciprofloxacin) tabs Levaquin® (levofloxacin) tabs, solution Ofloxacin</p>	<p>Cipro, Levaquin: the patient has had a documented intolerance to the generic equivalent. Baxdela: patient is completing a course of therapy with the requested medication that was initiated in the hospital OR patient is ≥ 18 years of age AND has a confirmed diagnosis of acute bacterial skin and skin structure infection (ABSSSI) AND current culture and sensitivity (C&S) report shows isolated pathogen is a gram-positive or gram-negative organism susceptible to delafloxacin (If obtaining a C&S report is not feasible, provider must submit documentation.) AND member has a documented treatment failure, intolerance or contraindication to 2 preferred antibiotics, one of which must be a fluoroquinolone AND duration of therapy does not exceed 14 days. Ofloxacin: patient has had a documented side effect, allergy, or treatment failure with two preferred fluoroquinolones</p>
RIFAMYCINS		
<p>All products require PA</p>	<p>Aemcolo® (rifamycin) delayed release tablets <i>QTY LIMIT: 12 tablets, max of 3 days</i> Xifaxan® (rifaximin) 200 mg tablets <i>QTY LIMIT: depends on indication</i> Xifaxan® (rifaximin) 550 mg tablets <i>QTY LIMIT: depends on indication</i></p>	<p>Aemcolo: patient has a diagnosis of traveler’s diarrhea caused by noninvasive strains of Escherichia coli AND Patient has had a documented side effect, allergy, treatment failure or contraindication with a fluoroquinolone or azithromycin. Xifaxan: Critical for Approval Based on Indication: Hepatic Encephalopathy (Xifaxan 550 mg Tablets Only): patient has a diagnosis of hepatic encephalopathy. AND Patient has had a documented side effect, allergy, treatment failure or contraindication to lactulose. AND Quantity limit is 2 tablets/day (550 mg tablets only). Traveler’s Diarrhea (Xifaxan 200 mg Tablets Only): patient has a diagnosis of traveler’s diarrhea caused by noninvasive strains of Escherichia coli. AND Patient has had a documented side effect, allergy, treatment failure or contraindication with a fluoroquinolone or azithromycin. AND Quantity limit is 9 tablets/RX (200 mg tablets only). Small Intestinal Bacterial Overgrowth (Xifaxan 550 mg or 200 mg Tablets): patient has a diagnosis of SIBO AND Quantity limit is 1,200 mg to 1,650mg/day for 14 days; maximum of 3 courses will be approved. Irritable Bowel Syndrome (Xifaxan 550 mg or 200 mg Tablets): patient has a diagnosis of irritable bowel syndrome without constipation or with symptoms of bloating. Quantity limit is 1,200 mg to 1,650 mg/day for 14 days; maximum of 3 courses will be approved. Inflammatory Bowel Disease: Crohn’s Disease (Xifaxan 550 mg or 200 mg Tablets): patient has a diagnosis of Crohn’s Disease. AND Patient has had a documented side effect, allergy, treatment failure or contraindication to one of the following: 6-mercaptopurine, azathioprine, corticosteroids, or</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>methotrexate. AND Quantity limit is 600 mg to 1,600 mg/day.</p> <p>Clostridium difficile Diarrhea (Xifaxan 200 mg Tablets): patient has a diagnosis of C. difficile diarrhea. AND Patient has had a documented side effect, allergy, treatment failure or contraindication to vancomycin AND Quantity limit is 1200mg/day.</p>
TETRACYCLINES		
<p>DOXYCYCLINE MONOHYDRATE 50 MG, 100 MG capsules, tablets</p> <p>DOXYCYCLINE HYCLATE 20MG tablets</p> <p>DOXYCYCLINE HYCLATE 100 MG capsules, tablets</p> <p>DOCYCYCLINE HYCLATE 50MG capsules</p> <p>DOXYCYCLINE MONOHYDRATE suspension 25 MG/5ML</p> <p>MINOCYCLINE 50 MG, 100 MG capsules</p>	<p>Demeclocycline 150mg, 300mg tabs</p> <p>Doryx (doxycycline hyclate) delayed release tabs</p> <p>Doxycycline hyclate delayed release tabs</p> <p>Doxycycline monohydrate 40mg cap</p> <p>Doxycycline 75mg, 150mg caps, tabs</p> <p>Minolira® ER (minocycline extended release) tablet</p> <p>QTY LIMIT: 1 tablet/day</p> <p>Minocycline 50 mg, 75 mg, 100 mg tabs</p> <p>Minocycline ER tablets (compare to Solodyn®)</p> <p>Nuzuira® (omadacycline) tabs</p> <p>QTY LIMIT: Max 14-day supply</p> <p>Solodyn®(minocycline) tabs ER</p> <p>Tetracycline 250 mg, 500 mg cap</p> <p>Vibramycin® (doxycycline hyclate) cap</p> <p>All other brands</p>	<p>Non-preferred doxycycline/minocycline products (except as listed below): patient has had a documented side effect, allergy, or treatment failure with a preferred doxycycline/minocycline. If a product has an AB rated generic, the trial must be the generic formulation.</p> <p>Nuzuira: patient has been started on intravenous or oral omadacycline in the hospital and will be finishing the course of therapy in an outpatient setting OR the patient has a diagnosis of community-acquired bacterial pneumonia (CABP) or acute bacterial skin and skin structure infections (ABSSSI) AND the patient has had a documented treatment failure with two preferred antibiotics (from any class) OR the provider submits clinical rationale as to why the preferred agents would not be appropriate for the patient.</p> <p>Minolira ER/Solodyn: patient is ≥ 12 years of age AND indication is to treat non-nodular inflammatory lesions of acne vulgaris AND patient has had a documented side effect, allergy, or treatment failure with a preferred minocycline. Note: no effect has been demonstrated on non-inflammatory acne lesions.</p> <p>Tetracycline: patient has had a documented side effect, allergy, or treatment failure with at least two preferred products OR the indication for use is the treatment of H. Pylori infection.</p>
ANTI-INFECTIVES ANTIFUNGAL		
ALLYLAMINES		
<p>TERBINAFINE tabs (compare to Lamisil®)</p> <p>QTY LIMIT: 30 tablets/month (therapy limit of 90 days)</p> <p>GRISEOFULVIN MICROSIZED Suspension</p>	<p>Griseofulvin Microsize Tablets</p> <p>Griseofulvin Ultramicrosized Tablets</p>	<p>Griseofulvin Microsize Tabs/Griseofulvin Ultramicrosized: patient has had a documented side effect, allergy, or treatment failure with terbinafine tablets and a preferred formulation of griseofulvin.</p>
AZOLES		
<p>FLUCONAZOLE (compare to Diflucan®) tabs, suspension</p> <p>CLOTRIMAZOLE Troche (compare to Mycelex®)</p>	<p>Cresemba® (isavuconazonium) caps</p> <p>Diflucan® (fluconazole) tabs, suspension</p> <p>Itraconazole (compare to Sporanox®) caps, solution</p>	<p>Cresemba: patient is completing a course of therapy that was initiated in the hospital OR patient has a diagnosis of mucormycosis OR patient has a diagnosis of invasive aspergillosis and has had a documented side effect, allergy, contraindication, or treatment failure with voriconazole.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>IV drugs are not managed at this time.</p>	<p>Ketoconazole tabs Noxafil[®] (posaconazole) oral suspension</p> <p>Noxafil[®] (posaconazole) DR Tablets <i>QTY LIMIT</i>: 93 tablets/30 days Noxafil[®] (posaconazole) DR Powder packets Oravig[®] (miconazole) 50 mg buccal tablet Posaconazole DR Tablets (compare to Noxafil[®]) <i>QTY LIMIT</i>: 93 tablets/30 days Posaconazole oral suspension (compare to Noxafil[®])</p> <p>Sporanox[®] (itraconazole) caps, solution Tolsura[®] (itraconazole) caps <i>QTY LIMIT</i>: 4 caps/day</p> <p>VFend[®] (voriconazole) tabs, suspension Vivjoa[®] (oteseconazole) caps Voriconazole (compare to VFend[®]) tabs, suspension</p>	<p>Ketoconazole/Itraconazole 100mg cap/Itraconazole Solution/Sporanox: patient has a documented side-effect, allergy, or treatment failure to at least ONE of the preferred medications OR patient is completing a course of therapy that was initiated in the hospital. For approval of Sporanox[®] capsules, the patient must have a documented intolerance to generic itraconazole. For approval of Itraconazole solution, the patient must have a medical necessity for a liquid dosage form.</p> <p>Limitations: Coverage of Onychomycosis agents will NOT be approved solely for cosmetic purposes.</p> <p>Tolsura: patient has a diagnosis of aspergillosis intolerant of or refractory to Amphotericin B therapy AND patient has a documented intolerance to both generic itraconazole and voriconazole OR patient has a diagnosis of blastomycosis or histoplasmosis AND the patient has a documented intolerance to itraconazole.</p> <p>Voriconazole/Vfend: Patient has a diagnosis of invasive aspergillosis. OR patient is completing a course of therapy with the requested medication that was initiated in the hospital. OR patient has a documented side-effect, allergy, or treatment failure to ONE of the preferred medications AND itraconazole. AND For approval of Vfend[®], the patient must have a documented intolerance to generic voriconazole. AND For approval of voriconazole suspension, the patient must have a medical necessity for a liquid dosage form.</p> <p>Noxafil tablet, Posaconazole tablet, Noxafil powder packets: patient has a diagnosis of HIV/immunocompromised status (neutropenia secondary to chemotherapy, hematopoietic stem cell transplant recipients) AND medication is being used for the prevention of invasive Aspergillosis/ Candida infections. OR patient is completing a course of therapy with the requested medication that was initiated in the hospital. Approval of powder packets will be limited to patients ≤ 12 years of age and < 40kg.</p> <p>Noxafil oral suspension, posaconazole oral suspension:</p> <ul style="list-style-type: none"> • Patient is completing a course of therapy with the requested medication that was initiated in the hospital OR • Patient has a diagnosis of HIV/immunocompromised status (neutropenia secondary to chemotherapy, hematopoietic stem cell transplant recipients) AND medication is being used for the prevention of invasive Aspergillosis/ Candida infections OR • Patient is being treated for oropharyngeal candidiasis and has a documented side-effect, allergy, or treatment failure to fluconazole and itraconazole. <p>Diflucan (brand): For approval of Diflucan brand name product, the patient must have a documented intolerance to generic fluconazole.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>Oravig: The indication for use is treatment of oropharyngeal candidiasis AND patient has had a documented side effect, allergy, or treatment failure/ inadequate response to both nystatin suspension and clotrimazole troche.</p> <p>Vivjoa: the patient is not of reproductive potential AND the patient has recurrent yeast infections despite a treatment course of 7-14 days with a preferred vaginal azole, a longer course of oral fluconazole (e.g. one dose every 3 days for a total of 3 doses), and Brexafemme.</p>
TRITERPENOIDS		
All products require PA	Brexafemme® (ibrexafungerp) tablets	<p>Brexafemme: The patient is not pregnant and has been counseled to use effective contraception during treatment and for 4 days after the last dose (if applicable) AND the patient has recurrent yeast infections despite a treatment course of 7-14 days with a preferred vaginal azole AND a longer course of oral fluconazole (e.g. one dose every 3 days for a total of 3 doses)</p>
ANTI-INFECTIVES ANTIMALARIALS		
<p>ATOVAQUONE/PROGUANIL (compare to Malarone®)</p> <p>CHLOROQUINE</p> <p>COARTEM® (artemether/lumefantrine)</p> <p>DARAPRIM® (pyrimethamine)</p> <p>HYDROXYCHLOROQUINE SULFATE</p> <p>MEFLOQUINE</p> <p>PRIMAQUINE</p> <p>QUINIDINE SULFATE</p> <p><u><i>Preferred After Clinical Criteria Are Met</i></u></p> <p>KRINTAFEL® (tafenoquine succinate)</p>	<p>Malarone® (atovaquone/proguanil)</p> <p>Pyrimethamine (compare to Daraprim®)</p> <p>Quinine Sulfate (compare to Qualquin®)</p> <p>Qualaquin® (quinine sulfate)</p>	<p>Krintafel: the patient is ≥ 16 years of age AND is receiving concurrent antimalarial therapy</p> <p>Malarone: patient has a documented intolerance to the generic equivalent</p> <p>Pyrimethamine: patient has a documented intolerance to brand Daraprim</p> <p>Quinine sulfate, Qualaquin: diagnosis or indication is for the treatment of malaria. (Use for leg cramps not permitted.) AND If the request is for brand Qualaquin, the patient has a documented intolerance to the generic equivalent.</p>
ANTI-PARASITICS		
<p>ALBENDAZOLE (compare to Albenza®)</p> <p>BILTRICIDE® (praziquantel)</p> <p>IVERMECTIN (compare to Stromectol®)</p>	<p>Benznidazole</p> <p>Emverm® (mebendazole)</p> <p>Lampit (nifurtimox)</p> <p>Stromectol® (ivermectin)</p>	<p>Benznidazole, Lampit: patient must be between 2-12 years of age (Benznidazole) or ≤ 18 years (Lampit) AND patient has a diagnosis of Chagas Disease (American trypanosomiasis) AND length of therapy does not exceed 60 days.</p> <p>Emverm: patient has a documented side effect, allergy, treatment failure, or contraindication to albendazole OR indication for use is hookworm infection (e.g. ancylostomiasis, necatoriasis, uninariasis).</p> <p>Stromectol: patient has a documented intolerance to the generic product.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

ANTI-INFECTIVES ANTI-VIRALS

HERPES SIMPLEX VIRUS MEDICATIONS (ORAL)		
<p>ACYCLOVIR (compare to Zovirax®) tablets, capsules ACYCLOVIR suspension (age ≤ 12 yrs) VALACYCLOVIR (compare to Valtrex®)</p>	<p>Famciclovir Sitavig[®] (acyclovir) Buccal Tablet <i>QTY LIMIT: 2 tablets/30 days</i> Valtrex[®] (valacyclovir)</p>	<p>Acyclovir suspension (age > 12 yrs): patient has a medical necessity for a non-solid oral dosage form Famciclovir: patient has a documented side effect, allergy, or treatment failure (at least one course of seven or more days) with acyclovir or valacyclovir. Sitavig: patient has a diagnosis of recurrent herpes labialis (cold sores), having at least 4 episodes in the previous year AND patient has a documented side effect or treatment failure with acyclovir AND valacyclovir. Valtrex: patient has a documented intolerance to the generic equivalent.</p>
INFLUENZA MEDICATIONS		
<p>OSELTAMIVIR (compare to Tamiflu®) <i>QTY LIMIT: 45 and 75 mg caps = 10 caps/30 days, 30 mg caps = 20 caps/30 days, 6 mg/ml suspension = 180ml/30 days</i></p>	<p>Relenza® (zanamivir) <i>QTY LIMIT: 20 blisters/30 days</i> Tamiflu® (oseltamivir) <i>QTY LIMIT: 45 and 75 mg caps = 10 caps/30 days, 30 mg caps = 20 capsule /30 days, 6 mg/ml suspension = 180 ml/30 days</i> Xofluza™ (baloxavir marboxil)</p>	<p>Relenza: There is a clinical, patient-specific reason the patient cannot use oseltamivir Tamiflu: Patient has a documented intolerance to generic Oseltamivir Xofluza: there is a clinical, patient-specific reason the patient cannot use oseltamivir. Note: A maximum of one single dose per 30 days will be approved based on the patient's body weight: 40mg (2 x 20mg tablets) for patients weighing between 40kg and 80kg or 80mg for patients weighing at least 80kg. Limitations: Amantadine and rimantadine are not CDC recommended for use in influenza treatment or chemoprophylaxis at this time and are not covered for this indication. For information regarding amantadine see "Parkinson's Medications"</p>
COVID MEDICATIONS		
<p>Paxlovid®(nirmatrelvir/ritonavir) tablets</p>		

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

CYTOMEGALOVIRUS (CMV) INFECTION MEDICATIONS		
---	--	--

<p>VALGANCICLOVIR (compare to Valcyte®) tablet</p>	<p>Livtency™ (maribavir) tablets Prevymis® (letermovir) Valcyte® tablets, solution Valganciclovir (compare to Valcyte®) solution</p>	<p>Livtency: Indication is for the treatment of CMV infection in a recipient of a hematopoietic stem cell or solid organ transplant AND infection is refractory to ganciclovir, valganciclovir, cidofovir, or foscarnet (as defined by >1 log₁₀ increase in CMV DNA levels in blood or serum after at least 14 days of therapy) AND medication will not be administered with ganciclovir or valganciclovir. For re-approval beyond 12 weeks, documentation must be submitted detailing continued medical necessity.</p> <p>Prevymis: <i>Prophylaxis of cytomegalovirus (CMV) infection in adult CMV-seropositive recipients [R+] of an allogenic hematopoietic stem cell transplant:</i> Therapy is initiated between day 0 and day 28 post-transplantation AND therapy will continue through day 100 post-transplantation (In patients at risk for late CMV infection and disease, Prevymis may be continued through Day 200 post-HSCT) AND for approval of injection, the patient must be unable to take oral medications.</p> <p><i>Prophylaxis of CMV disease in adult kidney transplant recipients at high risk (Donor CMV seropositive/Recipient CMV seronegative [D+/R-]):</i> Therapy is initiated between day 0 and day 7 post-transplantation AND therapy will continue through day 200 post-transplantation AND for approval of injection, the patient must be unable to take oral medications. AND the patient has a documented side effect, allergy, or contraindication to valganciclovir</p> <p>Valcyte: the patient has a documented intolerance to generic valganciclovir AND for approval of solution, the patient has a medical necessity for a non-solid oral dosage form.</p> <p>Valganciclovir solution: the patient has a medical necessity for a non-solid oral dosage form.</p>
--	---	--

INFLUENZA VACCINES		
--------------------	--	--

<p><u>INACTIVATED INFLUENZA VACCINE, (IIV), STANDARD DOSE (EGG BASED)</u> AFLURIA® Injection FLUARIX® Injection FLULAVAL® Injection FLUZONE® Injection</p>	<p><u>ADJUVANTED INACTIVATED INFLUENZA VACCINE, (IIV), STANDARD DOSE (EGG BASED)</u> Flud™ Injection</p> <p><u>INACTIVATED INFLUENZA VACCINE, (IIV), HIGH DOSE (EGG BASED)</u> Fluzone High-Dose® Injection</p> <p><u>RECOMBINANT INFLUENZA VACCINE, (RIV) (EGG FREE)</u> Flublok® Injection</p> <p><u>INACTIVATED INFLUENZA VACCINE, (ccIIV), STANDARD DOSE (CELL CULTURE BASED)</u> Flucelvax® Injection</p>	<p>Flucelvax: Prescriber provides clinical rationale why one of the preferred influenza vaccines cannot be used.</p> <p>Flublok: Patient is ≥ 65 years old OR Patient must have a documented severe reaction to egg based influenza vaccine AND the patient is unable to use Flucelvax.</p> <p>Flumist: Flumist is being requested for influenza prophylaxis during flu season AND The patient is between the ages of 19 and 49 years old, AND Prescriber provides documentation of a contraindication to an intramuscular injection or other compelling information to support the use of this dosage form.</p> <p>Fluzone High Dose, Flud: Patient is ≥ 65 years old OR Prescriber provides clinical rationale why one of the preferred influenza vaccines cannot be used. Note: the CDC and its Advisory Committee on Immunization Practices (ACIP) have not expressed a preference for any flu vaccine formulation for this age group.</p>
---	--	--

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

	<p><u>LIVE ATTENUATED INFLUENZA VACCINE, (LAIV4) (EGG BASED)</u> Flumist® Quadrivalent Intranasal</p>	
--	--	--

VACCINES - OTHER

<p><u>Preferred After Age Limit Is Met</u> ABRYSVO AREXVY GARDASIL SHINGRIX</p>		<p>Abrysvo: Covered if ≥ 60 years of age OR the vaccine will be administered during weeks 32 through 36 of pregnancy during September through January. Arexvy: Covered if ≥ 60 years of age. Gardasil: Covered for 19 years old to 45 years old (those under 19 should be referred to their pediatrician or PCP for state-supplied vaccine) Shingrix: Covered if ≥ 50 years of age</p> <p>Vaccines on the Advisory Committee on Immunization Practices (ACIP) list of recommended vaccines for children ≤ 18 years of age are supplied through the Vaccines for Children program administered by the Vermont Department of Health, and are not available through DVHA’s pharmacy Programs.</p> <ul style="list-style-type: none"> • Vaccines on the ACIP list of recommended vaccines for adults ≥ 19 years of age are available at many primary care provider offices and through the pharmacy programs. Vaccines are subject to the same limitations as the ACIP guideline recommendations. Providers who participate in the Blueprint for Health initiative must enroll in the Vaccines for Adults program administered by the Vermont Department of Health. The ACIP guidelines and information about enrollment in these programs can be found at http://healthvermont.gov/hc/imm/provider.aspx • Vaccines not on the recommended list may require Prior Authorization.
--	--	---

MIGRAINE THERAPY: PREVENTATIVE TREATMENTS

Calcitonin gene-related peptide (CGRP) Inhibitors: Initial approval is 6 months; renewals are 1 year

<p><u>Preferred After Clinical Criteria Are Met</u> AIMOVIG® (erenumab-aooe) QTY LIMIT: 1 injection (1mL) per 30 days AJOVY® (fremanezumab-vfrm) QTY LIMIT: 225 mg (1 injection) per 30 days or 675 mg (3 injections) every 90 days EMGALITY® (galcanezumab-gnlm) 120 mg/mL</p>	<p>Emgality ® (galcanezumab-gnlm) 100 mg/mL QTY LIMIT: 300 mg (3 injections) per 30 days, maximum of 6 months per year approved Nurtec® ODT (rimegepant) QTY LIMIT: 16 tablets/30 days Qulipta™ (atogepant)</p>	<p>Aimovig, Ajovy, Emgality 120mg/mL: The patient is 18 years of age or older AND patient has a diagnosis of episodic migraine (4-14 headache days per month with migraine lasting 4 hours or more) or chronic migraine (≥ 15 headache days per month, of which ≥ 8 are migraine days, for at least 3 months) AND patient has failed or has a contraindication to an adequate trial (≥ 60 days) of at least TWO medications for migraine prophylaxis from at least 2 different classes (tricyclic antidepressants, SNRI’s, beta-blockers, or anticonvulsants). Initial approval will be granted for 6 months. For re- approval after 6 months, the patient</p>
--	---	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

QTY LIMIT: 240 mg (2 injections) for the first 30 days followed by 120 mg (1 injection) per 30 days

Note: Please refer to “Botulinum Toxins” for Botox

QTY LIMIT: 30 tablets/30 days

Vyepti® (eptinezumab-jjmr)

must have documentation of a decrease in the number of headache days per month or decreased use of acute migraine medications such as triptans. Pharmacy claims will also be evaluated to assess compliance with the medication. Clinical justification must be provided if there is an increase in triptan use noted in the patient’s profile.

Nurtec ODT, Qulipta, Vyepti: The patient is 18 years of age or older AND The patient must have a documented side effect, allergy, or treatment failure to two preferred CGRP Inhibitors. Initial approval will be granted for 6 months. For reapproval after 6 months, the patient must have documentation of a decrease in the number of headache days per month or decreased use of acute migraine medications such as triptans. Pharmacy claims will also be evaluated to assess compliance with the medication. Clinical justification must be provided if there is an increase in triptan use noted in the patient’s profile.

Emgality 100mg/mL:

- Patient is 18 years of age or older AND
- Patient has a diagnosis of episodic cluster headache as defined by the following:
 - Severe to very severe unilateral pain felt in the orbital, supraorbital, and/or temporal regions lasting 15-180 minutes (when untreated)
 - Pain is accompanied by a sense of restlessness or agitation OR at least one of the following signs or symptoms, ipsilateral to the headache:
 - Conjunctival injection and/or lacrimation
 - Eyelid edema
 - Miosis and/or ptosis
 - Nasal congestion and/or rhinorrhea
 - Forehead and facial sweating
- Patient has ≥ 2 active cluster periods lasting 7 days to 1 year, separated by remission for periods lasting ≥ 3 months AND
- Patient has not achieved satisfactory response to adequate doses of corticosteroids (≥ 30mg prednisone or ≥ 16mg dexamethasone daily) started promptly at the start of the cluster period (Failure is defined as the need to use acute/abortive medications (oxygen, triptans, ergotamines, lidocaine) at least once daily for at least 2 days/week after the first full week of steroid therapy) AND

Patient has not achieved satisfactory response to adequate doses of verapamil (480mg/day, titrated up as needed to a max of 960mg/day) given for at least 3 weeks (Failure is defined as the need to use acute/abortive medications (oxygen, triptans, ergotamines, lidocaine) at least once daily for at least 2 days/week after 3 weeks of adequately dosed verapamil) **Note:** this requirement will be waived if the patient’s 2 most recent active cluster periods were less than 3 weeks in duration.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

MIGRAINE THERAPY: ACUTE TREATMENTS

GEPANTS		
<p>ORAL <i>Preferred After Clinical Criteria Are Met</i> NURTEC® ODT (rimegepant) <i>QTY LIMIT: 8 tablets/30 days</i></p> <p>NASAL SPRAY All products require PA</p>	<p>Ubrelyvy® (ubrogepant) <i>QTY LIMIT: 10 tablets/30 days</i></p> <p>Zavzpret™ (zavegepant) <i>QTY LIMIT: 8 units/30 days</i></p>	<p>Nurtec ODT: Patient has a documented side effect, allergy, or treatment failure with 2 preferred triptans, unless contraindicated.</p> <p>Ubrelyvy: Patient has a documented side effect, allergy, or treatment failure with 2 preferred triptans, unless contraindicated AND patient has a documented side effect, allergy, or treatment failure with Nurtec ODT.</p> <p>Zavzpret: Patient has a documented side effect, allergy, or treatment failure with 2 preferred triptans, one of which must be sumatriptan nasal spray, unless contraindicated AND patient has a documented side effect, allergy, or treatment failure with Nurtec ODT.</p>
DIHYDROERGOTAMINES		
<p>MIGRANAL® (dihydroergotamine mesylate) nasal spray <i>QTY LIMIT: 8 units/30 days</i></p>	<p>Dihydroergotamine mesylate nasal spray (compare to Migranal®) <i>QTY LIMIT: 8 units/30 days</i></p> <p>Trudhesa™ (dihydroergotamine mesylate) nasal spray <i>QTY LIMIT: 8 units/30 days</i></p>	<p>Dihydroergotamine, Trudhesa: The patient has a documented intolerance to Migranal nasal spray.</p>
DITANS		
<p>All products require PA</p>	<p>Reyvow® (lasmiditan) <i>QTY LIMIT: 8 tablets/30 days</i></p>	<p>Reyvow: Patient has a documented side effect, allergy, or treatment failure with 2 preferred triptans, unless contraindicated AND patient has a documented side effect, allergy, or treatment failure with Nurtec ODT AND counseling has been documented regarding the risks of driving impairment</p>
TRIPTANS		
<p>SINGLE AGENT ORAL ELETRIPTAN (compare to Relpax®) <i>QTY LIMIT: 12 tablets/30 days</i> FROVATRIPTAN (compare to Frova®) 2.5 mg <i>QTY LIMIT: 9 tablets/30 days</i> NARATRIPTAN <i>QTY LIMIT: 9 tablets/30 days</i> SUMATRIPTAN (compare to Imitrex®) <i>QTY LIMIT: 25 mg = 18 tablets/30 days, 50 and 100 mg = 9 tablets/30 days</i> RIZATRIPTAN (compare to Maxalt®) <i>QTY LIMIT: 12 tablets/30 days</i> RIZATRIPTAN ODT (compare to Maxalt-MLT®)</p>	<p>Almotriptan 6.25 mg, 12.5 mg <i>QTY LIMIT: 12 tablets/30 days</i> Frova® (frovatriptan) 2.5 mg <i>QTY LIMIT: 9 tablets/30 days</i> Imitrex® (sumatriptan) <i>QTY LIMIT: 25 mg = 18 tablets/30 days, 50 and 100 mg = 9 tablets/30 days</i> Maxalt® (rizatriptan) 5 mg, 10 mg tablet <i>QTY LIMIT: 12 tablets/30 days</i> Maxalt-MLT® (rizatriptan ODT) <i>QTY LIMIT: 12 tablets/30 days</i> Relpax® (eletriptan) 20 mg, 40 mg</p>	<p>Non-preferred single agents: The patient has had a documented side effect, allergy, or treatment failure with at least two preferred triptans. If a product has an AB rated generic, there must have also been a trial of the generic formulation.</p> <p>Sumatriptan/naproxen, Treximet: patient has had a documented side effect, allergy or treatment failure with 2 preferred Triptans, AND patient is unable to take the individual components separately.</p> <p>Zolmitriptan Nasal Spray, Zomig Nasal Spray, Onzetra Xsail, Tosymra: patient has had a documented side effect, allergy, or treatment failure with Sumatriptan Nasal Spray. For Zolmitriptan Nasal Spray, the patient must also have a documented intolerance to the brand Zomig Nasal Spray.</p> <p>Imitrex Injection, Zembrace: patient has had a documented intolerance to generic sumatriptan injection.</p> <p>To exceed quantity limits: patient is taking a medication for migraine prophylaxis.</p>

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

ANTI-PSYCHOTIC ATYPICAL & COMBINATIONS (CHILDREN < 18 YEARS OLD)

Preferred After Clinical Criteria Are Met TABLETS/CAPSULES

ARIPIRAZOLE (compare to Abilify®)

FDA maximum recommended dose = 30 mg/day

LURASIDONE (compare to Latuda®)

FDA maximum recommended dose =
80 mg/day

OLANZAPINE (compare to Zyprexa®)

FDA maximum recommended dose = 20 mg/day

RISPERIDONE (compare to Risperdal®)

FDA maximum recommended dose = 16 mg/day

PALIPERIDONE (compare to Invega®)

FDA maximum recommended dose =
12 mg/day

QUETIAPINE (compare to Seroquel®)

FDA maximum recommended dose = 800 mg/day

QUETIAPINE ER (compare to Seroquel® XR)

FDA maximum recommended dose =
800 mg/day

ZIPRASIDONE (compare to Geodon®)

FDA maximum recommended dose = 160 mg/day

Abilify® (aripiprazole)

FDA maximum recommended dose = 30 mg/day

Asenapine (compare to Saphris®)

QTY LIMIT: 2 tabs/day

FDA maximum recommended dose =
20 mg/day

Clozapine (compare to Clozaril®)

FDA maximum recommended dose = 900 mg/day

Clozaril® (clozapine)

FDA maximum recommended dose = 900 mg/day

Geodon® (ziprasidone)

FDA maximum recommended dose = 160 mg/day

Invega® (paliperidone)

QTY LIMIT: 3 and 9 mg = 1 tab/day, 6 mg = 2
tabs/day

FDA maximum recommended dose = 12 mg/day

Latuda® (lurasidone)

FDA maximum recommended dose = 80 mg/day

Risperdal® (risperidone)

FDA maximum recommended dose = 16 mg/day

Seroquel® (quetiapine)

FDA maximum recommended dose = 800 mg/day

Saphris® (asenapine)

QTY LIMIT: 2 tabs/day

FDA maximum recommended dose = 20 mg/day

Seroquel XR® (quetiapine XR)

FDA maximum recommended dose = 800 mg/day

Zyprexa® (olanzapine)

FDA maximum recommended dose = 20 mg/day

Target symptoms or Diagnosis that will be accepted for approval: Target Symptoms - Grandiosity/euphoria/mania; Obsessions/compulsions; Psychotic symptoms; Tics (motor or vocal). Diagnosis- Autism with Aggression and/or irritability; Disruptive Mood Dysregulation Disorder; Bipolar Disorder; Intellectual Disability with Aggression and/or Irritability; Major Depressive Disorder with psychotic features; Obsessive Compulsive Disorder; Schizophrenia/Schizoaffective Disorder; Tourette's Syndrome.

Criteria for approval of ALL drugs: Medication is being requested for one of the target symptoms or diagnoses listed above AND the patient is started and stabilized on the requested medication (Note: samples are not considered adequate justification for stabilization) OR Baseline labs including CBC, fasting glucose or HbA1C, and lipid profile have been completed AND patient meets additional criteria outlined below. Note: all requests for patients < 5 years will be reviewed by the DVHA medical director.

Asenapine, Saphris: patient has had a documented side effect, allergy or treatment failure with at least two preferred products (typical or atypical antipsychotics) one of which is risperidone or olanzapine ODT. For approval of Saphris, patient must have a documented intolerance to asenapine.

Abilify, Clozaril, Geodon, Invega, Latuda, Risperdal, Seroquel, Seroquel XR, Zyprexa: patient has a documented intolerance to the generic equivalent.

Clozapine: patient has had a documented side effect, allergy or treatment failure with at least three other antipsychotic medications (typical or atypical antipsychotics), two of which must be preferred agents.

Aripiprazole Oral Solution: patient has had a documented side effect, allergy or treatment failure with risperidone oral solution OR prescriber feels that risperidone would not be an appropriate alternative for the patient because of pre-existing medical conditions such as obesity or diabetes.

Versacloz Oral Solution: patient has had a documented side effect, allergy or treatment failure with at least three other antipsychotic medications (typical or atypical antipsychotics). AND patient is unable to use clozapine orally disintegrating tablets.

Aripiprazole ODT, Risperidone ODT, Zyprexa Zydis: Medical necessity for a specialty dosage form has been provided AND if the request is for Zyprexa Zydis, the patient has a documented intolerance to the generic equivalent.

Clozapine ODT: Medical necessity for a specialty dosage form has been provided AND patient has had a documented side effect, allergy or treatment

PREFERRED AGENTS

(No PA required unless otherwise noted)

NON-PREFERRED AGENTS

(PA required)

PA CRITERIA**Preferred After Clinical Criteria Are Met****ORAL SOLUTIONS**

RISPERIDONE (compare to Risperdal[®]) oral solution
 FDA maximum recommended dose = 16 mg/day

ORALLY DISINTEGRATING TABLETS**OLANZAPINE ODT (orally disintegrating tablets)**(compare to Zyprexa Zydys[®])*QTY LIMIT*: 5 and 10 mg = 1.5 tabs/day

FDA maximum recommended dose = 20 mg/day

Aripiprazole oral solution

FDA maximum recommended dose = 25 mg/day

Risperdal[®] (risperidone) oral solution

FDA maximum recommended dose = 16 mg/day

Versacloz[®] (clozapine) Oral Suspension*QTY LIMIT*: 18ml/day

FDA maximum recommended dose = 900 mg/day

Aripiprazole orally disintegrating tablets

QTY LIMIT: 10 and 15 mg = 2 tabs/day

FDA maximum recommended dose = 30 mg/day

Clozapine orally disintegrating tablets

FDA maximum recommended dose = 900 mg/day

Risperidone ODT

FDA maximum recommended dose = 16 mg/day

Zyprexa Zydys[®] (olanzapine orally disintegrating tablets)*QTY LIMIT*: 5 and 10 mg = 1.5 tabs/day

FDA maximum recommended dose = 20 mg/day

failure with at least three other antipsychotic medications (typical or atypical antipsychotics)

PREFERRED AGENTS

(No PA required unless otherwise noted)

NON-PREFERRED AGENTS

(PA required)

PA CRITERIA

ANTI-PSYCHOTIC ATYPICAL & COMBINATIONS (ADULTS ≥ 18 YEARS OLD)

TABLETS/CAPSULES

ARIPRAZOLE (compare to Abilify®)
 FDA maximum recommended dose = 30 mg/day

CLOZAPINE (compare to Clozaril®)
 FDA maximum recommended dose = 900 mg/day

LURASIDONE (compare to Latuda®)
 FDA maximum recommended dose = 160 mg/day

OLANZAPINE (compare to Zyprexa®)
 FDA maximum recommended dose = 20 mg/day

PALIPERIDONE (compare to Invega®)
 FDA maximum recommended dose = 12 mg/day

RISPERIDONE (compare to Risperdal®)
 FDA maximum recommended dose = 16 mg/day

QUETIAPINE (compare to Seroquel®)
 FDA maximum recommended dose = 800 mg/day

QUETIAPINE ER (compare to Seroquel® XR)
 FDA maximum recommended dose = 800 mg/day

ZIPRASIDONE (compare to Geodon®)
 FDA maximum recommended dose = 160 mg/day

Abilify® (aripiprazole)
 FDA maximum recommended dose = 30 mg/day

Abilify® Mycite (aripiprazole tablets with sensor)
QTY LIMIT: 1 tab/day
 FDA maximum recommended dose=30mg/day

Asenapine sublingual tablet (compare to Saphris®)
 FDA maximum recommended dose = 20 mg/day

Clozaril® (clozapine)
 FDA maximum recommended dose = 900 mg/day

Caplyta® (lumateperone)
QTY LIMIT: 1 capsule/day
 FDA maximum recommended dose = 42 mg/day

Fanapt® (iloperidone)
QTY LIMIT: 2 tablets/day
 FDA maximum recommended dose = 24 mg/day

Geodon® (ziprasidone)
 FDA maximum recommended dose = 160 mg/day

Invega® (paliperidone)
QTY LIMIT: 3 and 9 mg = 1 tab/day, 6 mg = 2 tabs/day
 FDA maximum recommended dose = 12 mg

Latuda® (lurasidone)
 FDA maximum recommended dose = 160 mg/day

Nuplazid™ (primavaserin)
QTY LIMIT: 2 tablets/day
 FDA maximum recommended dose = 34 mg

Rexulti® (brexpiprazole)
 FDA maximum recommended dose = 3 mg (adjunct of MDD) or 5 mg (schizophrenia)

Risperdal® (risperidone)
 FDA maximum recommended dose = 16 mg/day

Saphris® (asenapine) sublingual tablet
 FDA maximum recommended dose = 20 mg/day

Seroquel® (quetiapine)
 FDA maximum recommended dose = 800 mg/day

Seroquel XR® (quetiapine XR)
 FDA maximum recommended dose = 800 mg/day

Vraylar® (cariprazine)
QTY LIMIT: 1 capsule/day

Criteria for approval of ALL non-preferred drugs: patient has been started and stabilized on the requested medication (Note: samples are not considered adequate justification for stabilization.) OR patient meets additional criteria outlined below.

Caplyta:

Indication for use is schizophrenia/schizoaffective disorder: The patient has had a documented side effect, allergy or treatment failure with at least three preferred products (typical or atypical antipsychotics).

Indication for use is Bipolar Depression: the patient has had a documented side effect, allergy, or treatment failure with two preferred products (typical or atypical antipsychotics). If the prescriber feels that neither quetiapine or olanzapine/fluoxetine combination would be appropriate alternatives for the patient because of pre-existing conditions such as obesity or diabetes, the patient must have a documented side effect, allergy, or treatment failure with lurasidone.

Fanapt: The indication for use is the treatment of schizophrenia/schizoaffective disorder or bipolar disorder. AND The patient has had a documented side effect, allergy, or treatment failure with at least three preferred products (typical or atypical antipsychotics).

Asenapine, Saphris: The indication for use is the treatment of schizophrenia/schizoaffective disorder or bipolar disorder AND The patient has had a documented side effect, allergy, or treatment failure with at least two preferred products (typical or atypical antipsychotics), one of which is risperidone or olanzapine ODT. For approval of Saphris, patient must have a documented intolerance to asenapine.

Abilify, Clozaril, Geodon, Invega, Latuda, Risperdal, Seroquel, Seroquel XR and Zyprexa: patient has a documented intolerance to the generic equivalent.

Abilify Mycite: The patient has not been able to be adherent to aripiprazole tablets resulting in significant clinical impact (documentation of measures aimed at improving compliance is required) AND there is a clinically compelling reason why Abilify Asimulfii, Abilify Maintena or Aristada cannot be used. Initial approval will be granted for 3 months. For renewal, documentation supporting use of the tracking software must be provided and pharmacy claims will be evaluated to assess compliance with therapy.

Vraylar:

Indication for use is schizophrenia/schizoaffective disorder: the patient has had a

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>ORAL SOLUTIONS</u></p> <p>RISPERIDONE (compare to Risperdal®) oral solution FDA maximum recommended dose = 16 mg/day</p> <p><u>SHORT-ACTING INJECTABLE PRODUCTS</u></p> <p>GEODON® IM (ziprasidone intramuscular injection) FDA maximum recommended dose = 40 mg/day</p> <p>OLANZAPINE IM (compare to Zyprexa® IM) FDA maximum recommended dose = 30 mg/day</p> <p>ZYPREXA® IM (olanzapine intramuscular injection) FDA maximum recommended dose = 30 mg/day</p>	<p>FDA maximum recommended dose = 6 mg/day Zyprexa® (olanzapine) FDA maximum recommended dose = 20 mg/day</p> <p>Aripiprazole oral solution FDA maximum recommended dose = 25 mg/day</p> <p>Risperdal® (risperidone) oral solution FDA maximum recommended dose = 16 mg/day</p> <p>Versacloz® (clozapine) Oral Suspension <i>QTY LIMIT:</i> 18ml/day FDA maximum recommended dose = 900 mg/day</p>	<p>documented side effect, allergy or treatment failure with three preferred products (typical or atypical antipsychotics) OR</p> <p><i>Indication for use is Bipolar I depression:</i> the patient has had a documented side effect, allergy, or treatment failure with two preferred products (typical or atypical antipsychotics). If the prescriber feels that neither quetiapine or olanzapine/fluoxetine combination would be appropriate alternatives for the patient because of pre-existing conditions such as obesity or diabetes, the patient must have a documented side effect, allergy, or treatment failure with lurasidone.</p> <p><i>Indication for use is adjunct treatment of Major Depressive Disorder (MDD):</i> the patient has had a documented inadequate response to at least 3 different antidepressants from two different classes AND the patient has had a documented side effect, allergy, or treatment failure with two preferred atypical antipsychotic products being used as adjunctive therapy.</p> <p>Lybalvi: The patient has a documented side effect, allergy, or treatment failure with at least three antipsychotics, one of which must be aripiprazole or lurasidone AND There has been at least a 7-day opioid free interval from last use of short-acting opioids and at least a 14-day opioid free interval from last use of long-acting opioids.</p> <p>Nuplazid: The diagnosis or indication is the treatment of hallucinations/delusions associated with Parkinson’s Disease psychosis.</p> <p>Rexulti: <i>Indication for use is schizophrenia or agitation associated with dementia due to Alzheimer disease:</i> the patient has had a documented side effect, allergy or treatment failure with at least three preferred products (typical or atypical antipsychotics), one of which must be aripiprazole OR</p> <p><i>Indication for use is adjunct treatment of Major Depressive Disorder (MDD):</i> the patient has had a documented inadequate response to at least 3 different antidepressants from two different classes AND the patient has had a documented side effect, allergy or treatment failure with two preferred atypical antipsychotic products being used as adjunctive therapy, one of which must be aripiprazole</p> <p>Aripiprazole Oral Solution: the patient has had a documented side effect, allergy, or treatment failure with preferred risperidone oral solution.</p> <p>Risperdal Oral Solution: The patient has a documented intolerance to the generic product risperidone.</p> <p>Versacloz Oral Solution: The patient has a medical necessity for a non-solid oral dosage form and is unable to use clozapine orally disintegrating tablets.</p> <p>Invega Hafyera: The patient is started and stabilized on the medication OR The patient has been adequately treated with Invega Sustenna (paliperidone palmitate 1-month) for at least four months or Invega Trinza (paliperidone palmitate 3-</p>

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

LONG-ACTING INJECTABLE PRODUCTS

ABILIFY ASIMTUFI® (aripiprazole)

QTY LIMIT: 1 syringe/56 days

FDA maximum recommended dose = 960 mg/ 2 months

ABILIFY MAINTENA® (aripiprazole monohydrate)

QTY LIMIT: 1 vial/28 days

FDA maximum recommended dose = 400 mg/month

ARISTADA® (aripiprazole lauroxil)

QTY LIMIT: 441, 662, and 882 mg = 1 syringe/28 days, 1064 mg = 1 syringe/60 days

ARISTADA Initio™ (aripiprazole lauroxil)

INVEGA SUSTENNA® (paliperidone palmitate)

FDA maximum recommended dose = 234 mg/month

PERSERIS® (risperidone)

QTY LIMIT: 1 syringe/28 days

FDA maximum recommended dose = 120 mg/month

RISPERDAL® CONSTA (risperidone microspheres)

FDA maximum recommended dose = 50 mg/14 days

RYKINDO® (risperidone injection, extended-release)

Preferred After Clinical Criteria Are Met

INVEGA HAFYERA™ (paliperidone palmitate)

FDA maximum recommended dose = 1560 mg/6 months

INVEGA TRINZA® (paliperidone palmitate)

FDA maximum recommended dose = 819 mg/3 months

ORALLY DISINTEGRATING TABLETS

OLANZAPINE ODT (orally disintegrating tablets) (compare to Zyprexa Zydis®)

QTY LIMIT: 5 and 10 mg = 1.5 tabs/day

FDA maximum recommended dose = 20 mg/day

Uzedy™ (risperidone)

QTY LIMIT: 250 mg (0.7 ml)/2 months

Risperidone ER suspension (compare to Risperdal Consta)

Zyprexa Relprevv® (olanzapine pamoate)

QTY LIMIT: 405 mg = 1 vial/month, 210 and 300 mg = 2 vials/month

FDA maximum recommended dose = 600 mg/month

month) following at least one 3-month injection cycle.

Invega Trinza: The patient is started and stabilized on the medication OR tolerability has been established with Invega Sustenna for at least 4 months.

Note: This is processed via automated (electronic) step therapy.

Rykindo, Uzedy: Provider must submit clinical rationale detailing why the patient is unable to use Perseris or Risperdal Consta.

ORALLY DISINTEGRATING TABLETS: Medical necessity for a specialty dosage form has been provided AND If the request is Zyprexa Zydis, the patient has a documented intolerance to the generic equivalent.

Olanzapine/fluoxetine: The patient has had a documented side effect, allergy, or treatment failure with two preferred products OR The prescriber provides a clinically valid reason for the use of the requested medication.

Risperidone ER Injection: The patient has had a documented side effect, allergy, or treatment failure with Risperdal Consta

Secuado: The indication for use is the treatment of schizophrenia/schizoaffective disorder AND The patient has had a documented side effect, allergy or treatment failure with at least three preferred products (typical or atypical antipsychotics) and Saphris OR The indication for use is the treatment of schizophrenia/schizoaffective disorder AND the patient is unable to take oral medications AND the patient has had a documented side effect, allergy or treatment failure with a preferred long-acting injectable.

Zyorexa Relprevv: The indication for use is the treatment of schizophrenia/schizoaffective disorder AND The patient has had a documented side effect, allergy or treatment failure with a preferred product.

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

ANTI-PSYCHOTIC: TYPICALS

ORAL
HALOPERIDOL
LOXAPINE
PERPHENAZINE
PIMOZIDE
TRIFLUOPERAZINE

LONG ACTING INJECTABLE PRODUCTS
FLUPHENAZINE DECANOATE
HALOPERIDOL DECANOATE (compare to Haldol® decanoate)

Chlorpromazine
Fluphenazine
Molindone
Thioridazine
Thiothixene

Haldol® decanoate (haloperidol decanoate)

Chlorpromazine: patient has a diagnosis of acute intermittent porphyria or intractable hiccups OR patient has had a documented side effect, allergy or treatment failure with at least three preferred products (may be typical or atypical anti-psychotics).

Fluphenazine Oral Solution: patient has a requirement for an oral liquid dosage form (i.e. swallowing disorder, inability to take oral medications)

Fluphenazine tablets: patient is transitioning to the decanoate formulation or requires supplemental oral dosing in addition to decanoate OR patient has had a documented side effect, allergy or treatment failure with at least three preferred products (may be typical or atypical anti-psychotics).

All other oral medications: patient has had a documented side effect, allergy or treatment failure with at least three preferred products (may be typical or atypical anti-psychotics). If a product has an AB rated generic, one trial must be the generic.

Long Acting Injectable Products: for approval of Haldol decanoate, the patient has a documented intolerance to the generic product.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

ANTIRETROVIRAL THERAPY HUMAN IMMUNODEFICIENCY VIRUS (HIV)

SINGLE PRODUCT REGIMENS

<p><u>Tablets (STRs)</u> BIKTARVY® (bictegravir/emtricitabine/tenofovir AF) COMPLERA® (emtricitabine/rilpivirine/tenofovir) DELSTRIGO® (doravirine/lamivudine/tenofovir) DOVATO® (dolutegravir/lamivudine) EFAVIRENZ/EMTRICITABINE/TENOFOVIR GENVOYA® (elvitegravir/cobicistat/emtricitabine/tenofovir AF) ODEFSEY® (emtricitabine/rilpivirine/tenofovir AF) TRIUMEQ® (abacavir/lamivudine/dolutegravir) TRIUMEQ® PD tablets for oral suspension (abacavir/lamivudine/dolutegravir)</p> <p><u>Long-Acting Injectables</u> Cabenuva® (cabotegravir/rilpivirine) Kit</p>	<p>Juluca® (dolutegravir/rilpivirine) Symfi™ (efavirenz/lamivudine/tenofovir) Symfi™ Lo (efavirenz/lamivudine/tenofovir) Stribild® (elvitegravir/cobicistat/emtricitabine/tenofovir) Symtuza® (darunavir/cobicistat/emtricitabine/tenofovir AF)</p>	<p>Juluca: The patient has been started and stabilized on the requested medication (Note: samples are not considered adequate justification for stabilization.) OR patient is virologically suppressed (HIV-1 RNA < 50 copies per mL) on a stable oral antiretroviral regimen for at least 6 months AND the prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred products would not be suitable alternatives.</p> <p>Stribild:</p> <ul style="list-style-type: none"> • The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR • Genotype testing supporting resistance to other regimens OR • Intolerance or contraindication to preferred combination of drugs AND • CrCl > 70mL/min to initiate therapy OR CrCl > 50mL/min to continue therapy <p>Symfi, Symfi Lo: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred products would not be suitable alternatives</p> <p>Symtuza: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR Medical reasoning beyond convenience or enhanced compliance over preferred agents (Prezcobix & Descovy)</p>
COMBINATION PRODUCTS - NRTIs		
ABACAVIR/LAMIVUDINE ABACAVIR/LAMIVUDINE/ZIDOVUDINE LAMIVUDINE/ZIDOVUDINE		
COMBINATION PRODUCTS – NUCLEOSIDE & NUCLEOTIDE ANALOG RTIs		
DESCOVY® (emtricitabine/tenofovir AF) EMTRICITABINE/TENOFOVIR (compare to Truvada®)	Cimduo™ (lamivudine/tenofovir) Truvada® (emtricitabine/tenofovir)	<p>Cimduo: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred products would not be suitable alternatives.</p> <p>Truvada: patient must have a documented intolerance to the generic equivalent</p>
COMBINATION PRODUCTS – PROTEASE INHIBITORS		

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
LOPINAVIR/RITONAVIR (compare to Kaletra®)	Kaletra® (lopinavir/ritonavir)	Kaletra: patient must have a documented intolerance to generic lopinavir/ritonavir
ENTRY INHIBITORS-CCR5 CO-RECEPTOR ANTAGONISTS		
All products require PA	Maraviroc (compare to Selzentry®) Selzentry® (maraviroc)	Maraviroc, Selzentry: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred products would not be suitable alternatives. AND for approval of Maraviroc, the patient must have had a documented intolerance to Selzentry.
ENTRY INHIBITORS-FUSION INHIBITORS		
All products require PA	Fuzeon® (enfuvirtide)	Fuzeon: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred products would not be suitable alternatives.
INTEGRASE STRAND TRANSFER INHIBITORS		
ISENTRESS® (raltegravir potassium) ISENTRESS HD (raltegravir potassium) TIVICAY® (dolutegravir sodium) TIVICAY® PD (dolutegravir sodium)		
NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS (NRTI)		
ABACAIVR SULFATE (compare to Ziagen®) solution, tablet EMTRIVA® (emtricitabine) LAMIVUDINE (compare to Epivir®)	Epivir® (lamivudine) Retrovir® (zidovudine) Viread® (tenofovir disoproxil fumarate) 300mg tablet	Epivir, Retrovir, Viread 300mg, Ziagen: patient must have a documented intolerance to the generic equivalent
TENOFIVIR DISOPROXIL FUMARATE (compare to Viread®) 300mg VIREAD® (tenofovir disoproxil fumarate) 150mg, 200mg, 250mg tablet, 40mg/gm powder ZIDOVUDINE (compare to Retrovir®)	Ziagen® (abacavir sulfate) solution	
NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS (NNRTI)		
EDURANT® (rilpivirine) EFAVIRENZ (compare to Sustiva®) ETRAVIRINE (compare to Intelence®) PIFELTRO (doravirine)	Intelence® (etravirine) Nevirapine Nevirapine ER	Intelence: Patient must have a documented intolerance to Etravirine. Nevirapine, Nevirapine ER: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred products would not be suitable alternatives.
PHARMACOENHANCER-CYTOCHROME P450 INHIBITOR		

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
All products require PA	Tybost® (cobicistat)	Tybost: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR a clinically valid reason beyond compliance or convenience is given for not using a preferred combination drug or a ritonavir- based regimen with similar components
PRE-EXPOSURE PROPHYLAXIS (PrEP) AGENTS		
Apretude® (cabotegravir extended-release) 600mg/3mL IM injection Descovy® (emtricitabine/tenofovir AF) 200mg/25mg tablet Emtricitabine/Tenofovir DF (compare to Truvada®) 200mg/300mg tablet	Truvada® (Emtricitabine/Tenofovir DF) 200mg/300 mg tablet	Truvada: The patient has a documented intolerance to the generic equivalent.
PROTEASE INHIBITORS (PEPTICIC)		
ATAZANAVIR (compare to Reyataz®) EVOTAZ® (atazanavir/cobicistat) RITONAVIR (compare to Norvir®)	Fosemprenavir (compare to Lexiva®) Norvir® (ritonavir) Reyataz® (atazanavir) Viracept® (nelfinavir)	Fosemprenavir, Viracept: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred products would not be suitable alternatives. Norvir, Reyataz: patient must have a documented intolerance to the generic equivalent.
PROTEASE INHIBITORS (NON-PEPTIDIC)		
PREZCOBIX® (darunavir/cobicistat)	Aptivus® (tipranavir) Darunavir (compare to Prezista®) Prezista® (darunavir ethanolate)	Aptivus: The patient has been started and stabilized on the requested medication. (Note: samples are not considered adequate justification for stabilization.) OR The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred products would not be suitable alternatives. Darunavir, Prezista: The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why the combination product Prezcobix cannot be used AND for approval of darunavir, the patient must have a documented intolerance to brand Prezista.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

TREATMENT RESISTANT THERAPIES		
--------------------------------------	--	--

<p>All Products Require PA</p>	<p>Rukobia® (fostemsavir) <i>QTY LIMIT</i> = 2 tablets per day</p> <p>Sunlenca® (lenacapavir sodium)</p> <p>Trogarzo™ (ibalizumab-uiyk) <i>QTY LIMIT</i>: 10 vials (2000 mg) x 1 dose then 4 vials (800 mg) every 14 days thereafter</p>	<p>Sunlenca, Rukobia, Trogarzo: The patient must meet ALL of the following criteria:</p> <ul style="list-style-type: none"> • ≥ 18 years of age • Prescription is written by or in consultation with an infectious disease specialist. • Viral Load is ≥ 1,000 copies/mL (results must be submitted) • Patient has been compliant but has had an inadequate response to at least 6 months of treatment with anti-retroviral therapy (ART) • Patient has multi-drug resistant HIV-1 infection including documented resistance to at least one medication from each of the following classes: <ul style="list-style-type: none"> o Protease Inhibitor (PI) o Nucleoside Reverse Transcriptase Inhibitor (NRTI) o Non-nucleoside Reverse Transcriptase Inhibitor (NNRTI) • Medication will be used in combination with ART that includes at least one drug to which the individual’s virus is susceptible. <p>Initial approval will be granted for 6 months. For continuation of therapy, there must be a decrease in viral load from baseline AND the patient must continue to be compliant with the optimized background regimen of ART.</p>
--------------------------------	--	--

BILE SALTS AND BILIARY AGENTS		
--------------------------------------	--	--

<p>URSODIOL capsules</p>	<p>Bylvay™ (odevixibat)</p> <p>Chenodal® (chendiol)</p> <p>Cholbam® (cholic acid)</p> <p>Iqirvo® (elafibranor)</p> <p>Livmarli® (maralixibat)</p> <p>Ocaliva® (obeticholic acid)</p> <p>Rezdiffra™ (resmetirom)</p> <p>Urso® (Urosiol)</p> <p>Ursodiol tablets</p> <p>Urso® Forte (ursodiol)</p>	<p>Bylvay: The patient is experiencing moderate to severe pruritis associated with a diagnosis of progressive familial intrahepatic cholestasis (PFIC) confirmed by molecular genetic testing AND the patient does not have a ABCB11 variant resulting in non-functional or complete absence of the bile salt export pump protein (BSEP-3) AND the patient does not have a history of liver transplant or clinical evidence of decompensated cirrhosis AND baseline liver function tests and fat-soluble vitamin (A, D, E, and K) levels have been completed and will be monitored periodically during treatment AND patient has had an inadequate</p>
--------------------------	--	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>response or contraindication to cholestyramine and ursodiol. For re-approval, there must be documented clinical improvement (e.g. reduced serum bile acid or decreased pruritis).</p> <p>Chenodal: The indication for use is with radiolucent stones in well-opacifying gallbladders, in whom selective surgery would be undertaken except for the presence of increased surgical risk due to systemic disease or age AND the patient does not have any of the following contraindications to therapy: women who are pregnant or may become pregnant, known hepatocyte dysfunction or bile ductal abnormalities such as intrahepatic cholestasis, primary biliary cirrhosis or sclerosing cholangitis.</p> <p>Cholbam: The indication for use is the treatment of bile acid synthesis disorders due to single enzyme defects OR for the adjunctive treatment of peroxisomal disorders, including Zellweger spectrum disorders, AND the patient exhibits manifestations of liver disease, steatorrhea, or complications from decreased fat-soluble vitamin absorption AND the prescriber is a hepatologist or gastroenterologist. Initial approval will be granted for 3 months. For re-approval after 3 months, there must be documented clinical benefit.</p> <p>Livmarli: The patient is experiencing moderate to severe pruritis associated with a diagnosis of Alagille Syndrome (ALGS) AND baseline liver function tests and fat-soluble vitamin (A, D, E, and K) levels have been completed and will be monitored periodically during treatment AND patient has had an inadequate response or contraindication to cholestyramine and ursodiol. For re-approval, there must be documented clinical improvement (e.g. reduced serum bile acid or decreased pruritis).</p> <p>Iqirvo, Ocaliva: The indication for use is the treatment of primary biliary cholangitis (PBC) AND the patient has had an inadequate response or is unable to tolerate ursodiol. For approval of Iqirvo: Patient must have documented intolerance to Ocaliva.</p> <p>Rezdifra: The patient must have a diagnosis of nonalcoholic steatohepatitis (NASH) with fibrosis stage of F2 or F3 (clinical documentation provided) and NAFLD Activity Score (NAS) of at least 4 AND the patient does not have evidence of decompensated cirrhosis</p> <ul style="list-style-type: none"> • Prescribed by or in consultation with Gastroenterologist or Hepatologist • For reauthorization: Documentation provided indicated positive clinical response to therapy (improvement in or stabilization of fibrosis or resolution of NASH) AND the patient has not progressed to stage F4 (cirrhosis) <p>Urso, Ursodiol tablets, Urso Forte: The patient must have a documented treatment limiting side effect to generic ursodiol capsules.</p>

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

BONE RESORPTION INHIBITORS

ORAL BISPHOSPHONATES

TABLETS/CAPSULES

ALENDRONATE (compare to Fosamax[®]) tablets

IBANDRONATE

QTY LIMIT: 150 mg = 1 tablet/28 days

Actonel[®] (risedronate)
Alendronate oral solution
Atelvia (risedronate) Delayed Release Tablet
QTY LIMIT: 4 tablets/28 days
Fosamax[®] (alendronate)

Fosamax Plus D[®] (alendronate/vitamin D)
Risedronate (compare to Actonel[®])

INJECTABLE BISPHOSPHONATES

ZOLEDRONIC ACID Injection (compare to Reclast[®]) 5 mg/100mL

QTY LIMIT: 5 mg (one dose)/year

ZOLEDRONIC ACID Injection 4mg/5mL concentrate and 4 mg/100mL IV solution

Ibandronate Injection (compare to Boniva[®])
QTY LIMIT: 3 mg/3 months (four doses)/year

Reclast[®] Injection (zoledronic acid)
QTY LIMIT: 5 mg (one dose)/year

ESTROGEN AGONIST/ANTAGONIST

RALOXIFENE (compare to Evista[®]) Tablet

QTY LIMIT: 1 tablet/day

Evista[®] (raloxifene) Tablet
QTY LIMIT: 1 tablet/day

INJECTABLE RANKL INHIBITOR

All products require PA

Prolia[®] Injection (denosumab)
QTY LIMIT: 60 mg/6 months (two doses)/year
Xgeva[®] (denosumab)
QTY LIMIT: 120 mg/28 days

INJECTABLE SCLEROSTIN INHIBITOR

All products require PA

Evenity[®] (romosozumab-aqqg) injection
QTY LIMIT: 210 mg (2 syringes)/month
(Lifetime max duration = 12 months)

CALCITONIN NASAL SPRAY

All products require PA

Calcitonin Nasal Spray (compare to Miacalcin[®])

CALCITONIN INJECTION

All products require PA

Miacalcin[®] (calcitonin) Injection

Actonel, Atelvia, Risedronate: patient has had a documented side effect, allergy, or treatment failure (at least a six-month trial) to generic alendronate tablets and ibandronate AND if the request is for brand, the patient has also had a documented intolerance to generic equivalent.

Alendronate Oral Solution: prescriber provides documentation of medical necessity for the specialty dosage form (i.e. inability to swallow tablets, dysphagia).

Evista, Fosamax, Reclast: patient has a documented intolerance to the generic formulation.

Calcitonin Nasal: patient is started and stabilized on the requested medication.
Note: Calcitonin Nasal Spray (brand and generic) no longer recommended for osteoporosis.

Miacalcin Injection: patient has a diagnosis/indication of Paget's Disease

Fosamax Plus D: there is a clinical reason why the patient is unable to take generic alendronate tablets and vitamin D separately.

Forteo, Teriparatide patient has had a documented side effect, allergy, or treatment failure** to a bisphosphonate AND for approval for Forteo the patient has had a documented intolerance to generic Teriparatide.

Tymlos: patient has had a documented side effect, allergy, or treatment failure ** to a bisphosphonate and teriparatide AND prescriber has verified that the patient has been counseled about osteosarcoma risk.

Ibandronate Injection: patient has a diagnosis/indication of postmenopausal osteoporosis AND patient has had a documented side effect or treatment failure** to a preferred bisphosphonate.

Prolia Injection: patient has had a documented side effect, allergy, or treatment failure** to a preferred bisphosphonate OR medication is being used for osteopenia in women with breast cancer receiving adjuvant aromatase inhibitor therapy OR medication is being used for osteopenia in men receiving androgen deprivation therapy.

Xgeva Injection: diagnosis or indication is bone metastases from solid tumors (e.g. prostate, breast, thyroid, non-small lung cancer), multiple myeloma, hypercalcemia of malignancy, or giant cell tumor of bone.

Evenity Injection: diagnosis or indication is postmenopausal osteoporosis AND patient has no history of stroke or MI within the previous year AND patient has had a documented side effect or treatment failure** to a preferred bisphosphonate and Teriparatide.

**Treatment failure is defined as documented continued bone loss or fracture after one or more years despite treatment with a bisphosphonate.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>PARATHYROID HORMONE INJECTION</u> All products require PA</p>	<p>Forteo® (teriparatide) <i>QTY LIMIT</i>: 1 pen (2.4ml/30 days) Teriparatide (compare to Forteo®) <i>QTY LIMIT</i>: 1 pen/30 days Tymlos™ (abaloparatide) injection <i>QTY LIMIT</i>: 1 pen (1.56ml)/30 days (Lifetime max duration of treatment = 2 years)</p>	

BOTULINUM TOXINS

<p><u>Preferred After Clinical Criteria Are Met</u></p> <p>BOTOX® (onabotulinumtoxinA)</p> <p>DYSPORT® (abobotulinumtoxinA)</p>	<p>Myobloc® (rimabotulinumtoxinB) Xeomin® (incobotulinumtoxinA)</p>	<p>Criteria for approval of ALL drugs: The medication is being prescribed for an FDA approved indication AND the patient's age is FDA approved for the given indication AND the patient meets the following additional criteria (if applicable). Initial approval will be granted for 3 months unless otherwise noted. For re-approval, the patient must have documented improvement in symptoms.</p> <p><i>Additional criteria for Severe Axillary Hyperhidrosis (Botox only):</i> the patient failed an adequate trial of topical therapy.</p> <p><i>Additional criteria for Overactive bladder or detrusor overactivity (Botox only):</i> the patient failed an adequate trial of at least TWO urinary antispasmodics (either short- or long-acting formulations)</p> <p><i>Additional criteria for Chronic migraine (Botox only):</i> the patient has ≥ 15 headache days per month, of which ≥ 8 are migraine days, for at least 3 months AND the member has failed or has a contraindication to an adequate trial (≥ 60 days) of at least TWO medications for migraine prophylaxis from at least two different classes (tricyclic antidepressants, SNRI's, beta-blockers, or anticonvulsants). Initial approval will be granted for 6 months. For re-approval after 6 months, the patient must have documentation of a decrease in the number of headache days per month or decreased use of acute migraine medications such as triptans.</p> <p><i>Additional criteria for chronic sialorrhea (Myobloc and Xeomin):</i> the patient has a documented side effect, allergy, treatment failure, or contraindication to at least two anticholinergic agents (e.g. scopolamine, glycopyrrolate).</p> <p>LIMITATIONS: Coverage of botulinum toxins will not be approved for cosmetic use (e.g., glabellar lines, vertical glabellar eyebrow furrows, facial rhytides, horizontal neck rhytides, etc.). (BOTOX Cosmetic (onabotulinumtoxinA) is not covered)</p>
--	---	---

BPH AGENTS

ALPHA BLOCKERS

ALFUZOSIN ER
QTY LIMIT: 1 tablet/day
DOXAZOSIN (compare to Cardura®)
TAMSULOSIN (compare to Flomax®)
QTY LIMIT: 2 capsules/day
TERAZOSIN

ANDROGEN HORMONE INHIBITORS

DUTASTERIDE (compare to Avodart®)
QTY LIMIT: 1 capsule/day
FINASTERIDE (compare to Proscar®)
QTY LIMIT: 1 tablet/day

PDE-5 INHIBITORS

All products require PA

COMBINATION PRODUCT

All products require PA

Cardura® (doxazosin)
Cardura XL® (doxazosin)
QTY LIMIT: 1 tablet/day
Flomax® (tamsulosin)
QTY LIMIT: 2 capsules/day
Rapaflo® (silodosin)
QTY LIMIT: 1 capsule/day
Silodosin (compare to Rapaflo®)
QTY LIMIT: 1 tablet/day

Proscar® (finasteride)
QTY LIMIT: 1 tablet/day

Cialis® (tadalafil)
QTY LIMIT: 1 tablet/day
Tadalafil (compare to Cialis®)
QTY LIMIT: 1 tablet/day

Dutasteride/tamsulosin (compare to Jalyn®)
QTY LIMIT: 1 capsule/day
Entadfi™ (finasteride/tadalafil)
QTY LIMIT: 1 capsule/day
Jalyn® (dutasteride/tamsulosin)
QTY LIMIT: 1 capsule/day

Cardura, Cardura XL: The patient has had a documented side effect, allergy or treatment failure with two alpha blockers, one of which must be generic doxazosin.

Cialis, Tadalafil: The patient has a diagnosis of BPH (benign prostatic hypertrophy) AND the patient has a documented treatment failure/inadequate response to a preferred alpha blocker AND the patient has a documented treatment failure/inadequate response to a preferred 5-alpha reductase inhibitor AND for approval of Cialis, the patient must have a documented intolerance to the generic equivalent. Approval will be limited to 5mg daily for a maximum of 26 weeks.

Entadfi: The patient has a diagnosis of BPH (benign prostatic hypertrophy) AND the patient has a documented treatment failure/inadequate response to a preferred alpha blocker AND the patient has a documented treatment failure/inadequate response to a preferred 5-alpha reductase inhibitor AND the patient has a documented treatment failure/inadequate response to tadalafil. Approval will be limited to a maximum of 26 weeks.

Flomax: The patient has had a documented side effect, allergy or treatment failure with two preferred alpha blockers, one of which must be generic tamsulosin.

Rapaflo, Silodosin: The patient has had a documented side effect, allergy or treatment failure with two preferred alpha blockers

Proscar: The patient has a documented intolerance to the generic equivalent.

Dutasteride/tamsulosin, Jalyn: The patient has a diagnosis of BPH (benign prostatic hypertrophy) AND the patient has a documented treatment failure/inadequate response to combination therapy with generic tamsulosin and finasteride AND is unable to take tamsulosin and dutasteride as the individual separate agents AND for approval of Jalyn, the patient must have a documented intolerance to generic dutasteride/tamsulosin.

LIMITATIONS: Coverage of androgen hormone inhibitors will not be approved for cosmetic use in men or women (male-pattern baldness/alopecia or hirsutism). (This includes Propecia (finasteride) 1mg and its generic equivalent whose only FDA approved indication is for treatment of male pattern hair loss.).

BULK POWDERS

<https://dvha.vermont.gov/sites/dvha/files/documents/Covered%20Compounding%20Products%2011.22.23.pdf>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

CARDIAC GLYCOSIDES

DIGOXIN DIGOXIN Oral Solution		
----------------------------------	--	--

CLOSTRIDIUM DIFFICILE (C.diff) AGENTS

<p>FIRVANQ™ (vancomycin HCl) powder for oral solution <i>QTY LIMIT:</i> 1 bottle (150ml) per course of therapy. If more than 150ml is required, use of 300ml bottle is required.</p> <p>VANCOMYCIN (compare to Vancocin®) capsules</p>	<p>Dificid® (fidaxomicin) tablet <i>QTY LIMIT:</i> 20 tablets per 30 days</p> <p>Rebyota™ (fecal microbiota, live-jslm) suspension <i>QTY LIMIT:</i> 150 ml as a one time dose</p> <p>Vancocin® Vancomycin (compare to Vancocin®) oral solution <i>QTY LIMIT:</i> 1 bottle (150ml) per course of therapy. If more than 150ml is required, use of 300ml bottle is required.</p> <p>Vowst™ (fecal microbiota spores, live-brpk) capsule <i>QTY LIMIT:</i> 12 capsules/3 day supply</p> <p>Zinplava™ (bezlotoxumab) injection</p>	<p>Dificid: The patient’s diagnosis or indication is Clostridium difficile associated diarrhea (CDAD) AND for first time infection, the patient has had a side-effect, allergy, treatment failure or contraindication to oral vancomycin. OR patient is at high risk for relapse (age ≥ 65, immunocompromised, severe disease or Zar score ≥ 2).</p> <p>Vancomycin oral solution: The patient has a documented intolerance to Firvanq.</p> <p>Rebyota:</p> <ul style="list-style-type: none"> • The patient is 18 years of age or older AND • The patient has a diagnosis of Clostridium difficile infection (CDI) confirmed by a positive stool test AND • The patient has had at least 2 episodes of CDI recurrence after a primary episode (i.e., 3 episodes of CDI) or CDI recurrence after pulse dosed fidaxomicin (200 mg orally twice daily for 5 days, followed by once every other day for 20 days) AND • The patient has received at least 10 consecutive days of antibiotic therapy for the current CDI AND • Rebyota will be administered within 24 to 72 hours of completion of the current antibiotic regimen AND • The current CDI is controlled (i.e. <3 unformed/loose stools/day for 2 consecutive days) <p>Vancocin capsules: The patient has a documented intolerance to generic vancomycin capsules.</p> <p>Vowst:</p> <ul style="list-style-type: none"> • The patient is 18 years of age or older AND • The patient has a diagnosis of Clostridium difficile infection (CDI) confirmed by a positive stool test AND • The patient has a confirmed diagnosis of at least 2 recurrent episodes of
--	--	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>Clostridium difficile infection within 12 months (total of ≥ 3 episodes of CDI within 12 months) AND</p> <ul style="list-style-type: none"> • The patient has had a treatment failure (CDI recurrence) with pulse dose fidaxomicin, Zinplava AND either Rebyota or fecal transplant AND • The patient has received at least 10 consecutive days of antibiotic therapy for the current CDI AND • Vowst will be administered within 2 to 4 days of completion of the current antibiotic regimen AND • The current CDI is controlled (i.e. <3 unformed/loose stools/day for 2 consecutive days) <p>Zinplava:</p> <ul style="list-style-type: none"> • The patient is 18 years of age or older AND • The patient has a diagnosis of Clostridium difficile infection (CDI) confirmed by a positive stool test collected within the past 7 days AND • The patient is or will receive concomitant Standard of Care antibacterial therapy for CDI (e.g. vancomycin or fidaxomicin) AND • The patient is at high risk for recurrence based on at least one of the following: <ul style="list-style-type: none"> o Age ≥ 65 years o Two or more episodes of CDI within the past 6 months o The patient is immunocompromised o The patient has clinically severe CDI (e.g. fever, abdominal tenderness, WBC $\geq 15,000$ cells/mm³, albumin <30g/L, or renal failure)
CUSHING'S DISEASE		
All products require PA	<p>Isturisa® (osilodrostat) tablets Korlym® tablets (mifepristone) <i>QTY LIMIT: 4 tablets/day</i> Signifor® (pasireotide) Ampules <i>QTY LIMIT: all strengths = 2 ml (2 amps)/day</i> Maximum day supply = 30 days</p>	<p>Korlym: Patient is ≥ 18 years of age AND Patient has a diagnosis of endogenous Cushing's syndrome AND Patient is diagnosed with type 2 diabetes mellitus or glucose intolerance AND Patient has hyperglycemia secondary to hypercortisolism AND Patient has failed or is not a candidate for surgery AND Patient has a documented side effect, allergy, treatment failure or contraindication to at least 2 adrenolytic medications (e.g. ketoconazole, etomidate) AND Patient does not have any of the following contraindications to Korlym: Pregnancy (pregnancy must be excluded before the initiation of therapy or if treatment is interrupted for >14 days in females of reproductive potential. Nonhormonal contraceptives should be used during and one month after stopping treatment in all women of reproductive potential) OR Patient requires concomitant treatment with systemic corticosteroids for serious medical conditions/illnesses (immunosuppression for organ transplant) OR Patient has a history of unexplained vaginal bleeding OR Patient has endometrial hyperplasia with atypia or endometrial carcinoma OR Patient is concomitantly taking simvastatin, lovastatin, or a CYP3A substrate with a narrow therapeutic index (e.g., cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozone,</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

quinidine, sirolimus, or tacrolimus).

Isturisa, Signifor: Patient has a diagnosis of (pituitary) Cushing’s disease AND Patient is 18 years of age or older AND Pituitary surgery is not an option or has not been curative Note: Re-approval requires confirmation that the patient has experienced an objective response to therapy (i.e., clinically meaningful reduction in 24-hour urinary free cortisol levels and/or improvement in signs or symptoms of the disease).

GASTROINTESTINAL AGENTS: BOWEL PREP AGENTS, CONSTIPATION/DIARRHEA, IRRITABLE BOWEL SYNDROME-CONSTIPATION (IBS-C), IRRITABLE BOWEL SYNDROME-DIARRHEA (IBS-D), SHORT BOWEL SYNDROME, OPIOID INDUCED CONSTIPATION

Constipation: Chronic, IBS-C, or Opioid-Induced: Length of approval for non-preferred agents: Initial PA of 3 months and & 12 months thereafter

BULK-PRODUCING LAXATIVES

PSYLLIUM

OSMOTIC LAXATIVES

LACTULOSE

POLYETHYLENE GLYCOL 3350 (PEG)

STIMULANT LAXATIVE

BISACODYL

SENNA

STOOL SOFTENER

DOCUSATE

MISCELLANEOUS

DICYCLOMINE

GUANYLATE CYCLASE-C AGONIST

LINZESS® (linaclotide (age ≥ 6 years)

QTY LIMIT: 1 capsule/day

TRULANCE® (plecanatide) (age ≥ 6 years)

QTY LIMIT: 1 tablet/day

Note: Linzess® and Trulance® are contraindicated in patients less than 6 years of age due to the risk of serious dehydration.

Enemeez enema

Relistor® (methylnaltrexone) tablets
QTY LIMIT: 3 tablets/day

Enemeez enema: Patient had a trial and failure of contraindication to a preferred generic

Relistor Tablets, Symproic: The patient is current using an opiate for at least 4 weeks AND has documented opioid-induced constipation AND has had a documented side effect, allergy, or treatment failure to Amitiza and Movantik.

Relistor Injection: The patient must have documented opioid-induced constipation and be receiving palliative care AND the patient must have had documented treatment failure to a 1 week trial of 2 preferred laxatives from 2 different laxative classes used in combination.

Ibsrela, Motegrity: The patient is 18 years of age or older. AND the patient has had a documented side effect, allergy, or treatment failure to lubiprostone and either Linzess or Trulance.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>CIC-2 CHLORIDE CHANNEL ACTIVATORS</u> LUBIPROSTONE <i>QTY LIMIT: 2 capsules/day</i></p> <p><u>OPIOID ANTAGONISTS</u> MOVANTIK® (naloxegol) <i>QTY LIMIT: 1 tablet/day</i></p> <p><u>5-HT4 RECEPTOR ANTAGONISTS</u> All products require PA</p> <p><u>NHE3 INHIBITORS</u> All products require PA</p>	<p>Relistor® (methylnatrexone) injection Symproic® (naldemedine) <i>QTY LIMIT: 1 tablet/day</i></p> <p>Motegrity® (prucalopride) <i>QTY LIMIT: 1 tablet/day</i></p> <p>Ibsrela® (tenapanor) <i>QTY LIMIT: 2 tablets/day</i></p>	
Short Bowel Syndrome (SBS): Length of approval: 6 Months		
<p>All products require PA</p>	<p>Gattex® (teduglutide) Vials Maximum day supply = 30 days</p>	<p>Gattex: Patient has a diagnosis of short bowel syndrome AND Patient is receiving specialized nutritional support administered intravenously (i.e. parenteral nutrition) AND Patient does not have an active gastrointestinal malignancy (gastrointestinal tract, hepatobiliary, pancreatic), colorectal cancer, or small bowel cancer. Note: Re-approval requires evidence of decreased parenteral nutrition support from baseline.</p>
Antidiarrheal: HIV/AIDS: Length of approval: Initial approval 3 months, subsequent 1 year		
<p>DIPHENOXYLATE/ATROPINE LOPERAMIDE</p>	<p>Mytesi® (crofelemer) 125 mg DR Tablets <i>QTY LIMIT: 2 tablets/day</i></p>	<p>Mytesi: Patient has HIV/AIDS and is receiving anti-retroviral therapy AND Patient is at least 18 years of age AND Patient requires symptomatic relief of noninfectious diarrhea AND Infectious diarrhea (e.g. cryptosporidiosis, c. difficile, etc.) has been ruled out AND Patient has tried and failed at least one anti-diarrheal medication (i.e. loperamide or atropine/diphenoxylate)</p>
Antidiarrheal: IBS-D: Length of approval: Initial approval 3 months; subsequent 1 year		
<p>All products require PA</p>	<p>Alosetron (compare to Lotronex®) Lotronex® (alosetron) Viberzi® (eluxadoline) Xermelo™ (telotristat ethyl) <i>QTY LIMIT: 3 tablets/day</i></p>	<p>Lotronex/alosetron: The patient is a woman and has a diagnosis of severe diarrhea-predominant irritable bowel syndrome (IBS) with symptoms lasting 6 months or longer AND has had anatomic or biochemical abnormalities of the GI tract excluded AND has not responded adequately to conventional therapies such as loperamide and TCA's. For approval of generic alosetron, the patient must have documented intolerance to brand Lotronex.</p> <p>Viberzi: The patient has a diagnosis of IBS-D AND does not have any of the following contraindications to therapy A) known or suspected biliary duct</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>obstruction, or sphincter of Oddi disease or dysfunction B) alcoholism, alcohol abuse, alcohol addiction, or drink more than 3 alcoholic beverages/day C) a history of pancreatitis; structural diseases of the pancreas D) severe hepatic impairment (Child-Pugh Class C) AND has not responded adequately to conventional therapies such as loperamide and TCA's.</p> <p>Xermelo: The patient has a diagnosis of carcinoid syndrome diarrhea AND had an inadequate treatment response (defined as 4 or more bowel movements per day) despite use of a long-acting somatostatin analog for at least 3 consecutive months AND the medication will be used in combination with a long-acting somatostatin analog therapy. For reauthorization, documentation showing a decrease in the number of bowel movements per day is required. Note: Xermelo will not be approved in treatment naïve patients or as monotherapy.</p>

BOWEL PREP AGENTS

<p>CLENPIQ® GAVILYTE-C, GAVILTYE-G, GAVILYTE-H, GAVILYTE-N GOLYTELY MOVIPREP PEG-3350</p>	<p>Nulytely Plenvu® Sodium sulfate/Potassium sulfate/Magnesium sulfate (compare to Suprep®) Suflave™ Suprep® (sodium sulfate/potassium sulfate/magnesium sulfate) Sutab®</p>	<p>Non-preferred agents: The patient has a documented intolerance or treatment failure of at least one preferred agent (defined by failure to complete cleansing of the colon as a preparation for colonoscopy) AND if the product has an AB rated generic, there must have been a trial with the generic formulation.</p>
--	---	---

CONTINUOUS GLUCOSE MONITORS

Initial approval will be granted for 6 months; renewals up to 1 year thereafter

<p><u><i>Preferred After Clinical Criteria Are Met</i></u> DEXCOM G6 Initial prescription: 1 receiver, 1 wireless transmitter, and 9 sensors Refill Quantity Limits: 1 transmitter every 3 months, 1 sensor every 10 days (maximum of 9 sensors every 90 days) DEXCOM G7 Initial prescription: 1 receiver, 9 sensors Refill Quantity Limits: 1 sensor every 10 days (maximum of 9 sensors every 90 days)</p>	<p>Medtronic Guardian™ Connect Initial Prescription: 1 transmitter, 5 sensors Refill Quantity Limits: 1 transmitter every year, 1 sensor every 7 days (maximum of 5 sensors every 35 days) Medtronic 670G Guardian Link 3 Initial Prescription: 1 transmitter, 5 sensors Refill Quantity Limits: 1 transmitter every year, 1 sensor every 7 days (maximum of 5 sensors every 35 days) Medtronic 770G Guardian Link 3 Initial Prescription: 1 transmitter, 5 sensors</p>	<ul style="list-style-type: none"> • Patient has a diagnosis of Diabetes Mellitus AND patient age is FDA approved for the requested product AND one of the following criteria are met: <ul style="list-style-type: none"> ○ The patient requires treatment with insulin OR ○ The patient has a history of problematic hypoglycemia AND medications that could contribute to hypoglycemia (e.g. sulfonylureas, meglitinides) have been discontinued AND there is documentation of at least one of the following: Recurrent level 2 hypoglycemic events (glucose <54mg/dL (3.0mmol/L)) that persist despite multiple attempts to adjust medication(s) and/or modify the diabetes treatment plan OR a history of one level 3 hypoglycemic event (glucose <54mg/dL (3.0mmol/L))
--	---	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>FREESTYLE LIBRE 14 DAY (14-DAY SENSORS) Initial Prescription: 1 reader, 6 sensors Refill Quantity Limits: 1 sensor every 14 days (maximum of 6 sensors every 84 days)</p> <p>FREESTYLE LIBRE 2 PLUS (15-DAY SENSORS) Initial Prescription: 1 reader, 6 sensors Refill Quantity Limits: 1 sensor every 15 days (maximum of 6 sensors every 90 days)</p> <p>FREESTYLE LIBRE 3 PLUS (15-DAY SENSORS) Initial Prescription: 1 reader, 6 sensors Refill Quantity Limits: 1 sensor every 15 days (maximum of 6 sensors every 90 days)</p>	<p>Refill Quantity Limits: 1 transmitter every year, 1 sensor every 7 days (maximum of 5 sensors every 35 days)</p> <p>Medtronic 780G Guardian 4 Initial Prescription: 1 transmitter, 5 sensors Refill Quantity Limits: 1 transmitter every year, 1 sensor every 7 days (maximum of 5 sensors every 35 days)</p> <p>Medtronic MiniLink (includes Enlite Serter) Initial Prescription: 1 transmitter, 5 sensors Refill Quantity Limits: 1 transmitter every year, 1 sensor every 7 days (maximum of 5 sensors every 35 days)</p>	<p>characterized by altered mental and/or physical state requiring third party assistance for treatment of hypoglycemia</p> <ul style="list-style-type: none"> Approval of non-preferred products will be limited to cases where the CGM is directly integrated with the patient’s insulin pump. The make and model of pump must be documented on the prior authorization. <p>Re-authorization:</p> <ul style="list-style-type: none"> There is documented evidence of compliance to CGM (log data and/or office visit notes required). Replacement will be considered when medically necessary and not for recent technology upgrades (device must be malfunctioning and out of warranty). Initial Renewal Only: claims history shows a reduction in test strip utilization; for those using the same number of test strips after initiating a CGM, clinical justification needs to be provided for the continued use of a CGM.

CONTRACEPTIVES

SELECT PRODUCTS: Length of approval: 1 year
MONOPHASIC AGENTS:

Due to the extensive list of products, any monophasic BCP not listed as non-preferred is considered preferred.

Blisovi FE 24 (norethindrone/ethinyl estradiol/FE)
Drospirenone/ethinyl estradiol/levomefol
Kaitlib (norethindrone/ethinyl estradiol/FE)
Layolis FE (norethindrone/ethinyl estradiol/FE)
Lo-Estrin (norethindrone/ethinyl estradiol)
Lo-Estrin FE (norethindrone/ ethinyl estradiol/FE)
Melodetta FE (drospirenone/ethinyl estradiol/levomefol)
Mibelis FE (norethindrone/ethinyl estradiol/FE)
Nexstellis (drospirenone/estetrol)
Noretin-Eth Estra-Ferros Fum Tab Chew 0.8-25(24)
(norethindrone/ethinyl estradiol/FE)
Noretin-Eth Estra-Ferros Fum Tab Chew 1MG-20(24)
(norethindrone/ethinyl estradiol/FE)
Ogestrel (norgestrel/ethinyl estradiol)

Non-preferred agents: Trial with at least three preferred contraceptive products including the preferred formulation of the requested non-preferred agent

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
	Taytulla (norethindrone/ethinyl estradiol/FE) Wymza FE (norethindrone/ethinyl estradiol/FE)	
BIPHASIC AGENTS		
AZURETTE (desogestrel/ ethinyl estradiol) BEKYREE (desogestrel/ethinyl estradiol) DESOGESTREL/ETHINYL ESTRADIOL KARIVA (desogestrel/ ethinyl estradiol) KIMIDESS (desogestrel/ethinyl estradiol) LO LOESTRIN FE (norethindrone/ ethinyl estradiol/FE) NORETHIDRONE/ETHINYL ESTRADIOL 0.5/1-35 PIMTREA (desogestrel/ ethinyl estradiol) SIMLIYA (desogestrel/ethinyl estradiol) VIORELE (desogestrel/ ethinyl estradiol) VOLNEA (desogestrel/ethinyl estradiol)	Mircette (desogestrel/ ethinyl estradiol)	Non-preferred agents: Trial with at least three preferred contraceptive products including the preferred formulation of the requested non-preferred agent
TRIPHASIC AGENTS		
ALYACEN (norethindrone ethinyl estradiol) ARANELLE (norethindrone/ethinyl estradiol) CAZIAN (desogestrel/ ethinyl estradiol) CYCLAFEM (norethindrone/ethinyl estradiol) DASETTA (norethindrone/ethinyl estradiol) ENPRESSE (levonorgestrel/ ethinyl estradiol) LEENA (norethindrone/ethinyl estradiol) LEVONEST (levonorgestrel/ ethinyl estradiol) NATAZIA (dienogest/estradiol valerate) NORGESTIMATE/ETHINYL ESTRADIOL NORTREL 7/7/7 (norethindrone/ethinyl estradiol) PIRMELLA (norethindrone/ethinyl estradiol) TRI-ESTARYLLA (norgestimate/ ethinyl estradiol) TRI-FEMYNOR (norgestimate/ ethinyl estradiol) TRI-LINYAH (norgestimate/ ethinyl estradiol) TRI-LO-ESTARYLLA (norgestimate/ethinyl estradiol) TRI-LO-MARZIA (norgestimate/ethinyl estradiol) TRI-LO-SPRINTEC (norgestimate/ethinyl estradiol) TRI-PREVIFEM (norgestimate/ ethinyl estradiol) TRI-SPRINTEC (norgestimate/ ethinyl estradiol)	Ectopic FE (norethindrone/ethinyl estradiol/FE) Tilia FE (norethindrone/ethinyl estradiol/FE) Tri-Legest FE (norethindrone/ethinyl estradiol/FE)	Non-preferred agents: Trial with at least three preferred contraceptive products including the preferred formulation of the requested non-preferred agent

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
TRI-VYLIBRA (norgestimate/ ethinyl estradiol) TRI-VYLIBRA LO (norgestimate/ ethinyl estradiol) TRIVORA (levonorgestrel/ ethinyl estradiol) VELIVET (desogestrel/ ethinyl estradiol)		
EXTENDED CYCLE		
AMETHIA (levonorgestrel/ ethinyl estradiol) AMETHIA LO (levonorgestrel/ ethinyl estradiol) AMETHYST (levonorgestrel/ ethinyl estradiol) ASHLYNA (levonorgestrel/ ethinyl estradiol) CAMRESE (levonorgestrel/ ethinyl estradiol) CAMRESE LO (levonorgestrel/ ethinyl estradiol) DAYSEE (levonorgestrel/ ethinyl estradiol) INTROVALE (levonorgestrel/ ethinyl estradiol 3MTH) JAIMIESS (levonorgestrel/ ethinyl estradiol) JOLESSA (levonorgestrel/ ethinyl estradiol 3MTH) LEVONORGESTREL/ETHINYL ESTRADIOL TBDSPK 3 month LO-SEASONIQUE (levonorgestrel/ ethinyl estradiol) SIMPESE (levonorgestrel/ ethinyl estradiol) SEASONIQUE (levonorgestrel/ ethinyl estradiol) SETLAKIN (levonorgestrel/ethinyl estradiol)	Fayosim (levonorgestrel/ ethinyl estradiol) Quartette (levonorgestrel/ ethinyl estradiol) Rivelsa (levonorgestrel/ ethinyl estradiol)	Non-preferred agents: Trial with at least three preferred contraceptive products including the preferred formulation of the requested non-preferred agent
PROGESTIN ONLY CONTRACEPTIVES		
CAMILA (norethindrone) DEBLITANE (norethindrone) ERRIN (norethindrone) HEATHER (norethindrone) INCASSIA (norethindrone) JENCYCLA (norethindrone) JOLIVETTE (norethindrone) LYZA (norethindrone) NORA-BE (norethindrone) NORETHINDRONE 0.35MG NORLYNDA (norethindrone) SHAROBEL (norethindrone) TULANA (norethindrone)	Slynd® (drospirenone)	Non-preferred agents: Trial with at least three preferred contraceptive products including the preferred formulation of the requested non-preferred agent.
INJECTABLE CONTRACEPTIVES		

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
MEDROXYPROGESTERONE ACETATE 150MG (IM) VIAL/SYRINGE DEPO-PROVERA 104 (SUB-Q) SYRINGE (medroxyprogesterone acetate)	Depo-Provera (IM) (medroxyprogesterone acetate) 150 mg Susp vial/syringe	Depo-Provera IM: Patient must have a documented intolerance to medroxyprogesterone acetate 150mg.
VAGINAL RING		
NUVARING® (etonogestrel/ethinyl estradiol vaginal ring)	Annovera® (segesterone acetate/ethinyl estradiol vaginal ring) <i>QTY LIMIT:</i> 1 ring/year Eluryng (etonogestrel/ethinyl estradiol vaginal ring) Enilloring (etonogestrel/ethinyl estradiol vaginal ring) Etonogestrel/ethinyl estradiol vaginal ring Haloette (etonogestrel/ethinyl estradiol vaginal ring)	Non-preferred agents: Trial with at least three preferred contraceptive products including the preferred formulation of the requested non-preferred agent.
LONG ACTING REVERSIBLE CONTRACEPTIVES (LARCs)		
KYLEENA (levonorgestrel) IUD LILETTA (levonorgestrel) IUD MIRENA (levonorgestrel) IUD PARAGARD (copper) IUD SKYLA (levonorgestrel) IUD NEXPLANON (etonogestrel) Implant		
TOPICAL CONTRACEPTIVES		
TWIRLA® (levonorgestrel/ethinyl estradiol) patch XULANE PATCH (norelgestromin/ ethinyl estradiol)	Zafemy (norelgestromin/ ethinyl estradiol) patch	Zafemy: Trial with at least three preferred contraceptive products including the preferred formulation of the requested non-preferred agent.
VAGINAL CONTRACEPTIVES		
Please refer to the DVHA website for covered OTC spermicidal gels https://dvha.vermont.gov/sites/dvha/files/documents/OTCWebList_0.pdf	Phexxi™ (lactic acid, citric acid, and potassium bitartrate) vaginal gel	Phexxi: Use of hormonal contraceptives is contraindicated AND the patient has a documented side effect or allergy to nonoxynol-9
EMERGENCY CONTRACEPTIVES		
AFTERA (levonorgestrel) ECONTRA EZ (levonorgestrel) LEVONORGESTREL MY CHOICE (levonorgestrel)		

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
MY WAY (levonorgestrel) NEW DAY (levonorgestrel) OPCICON ONE-STEP (levonorgestrel) OPTION 2 (levonorgestrel)		
CORONARY VASODILATORS/ANTIANGINALS/SINUS NODE INHIBITORS		
ORAL		
ISOSORBIDE DINITRATE tablet (compare to Isordil®) ISOSORBIDE DINITRATE ER tablet ISOSORBIDE MONONITRATE tablet ISOSORBIDE MONONITRATE ER tablet NITROGLYCERIN SPRAY LINGUAL (compare to Nitrolingual Pump Spray®) NITROSTAT® (nitroglycerin SL tablet) RANOLAZINE SR 12 HR (compare to Ranexa®) <i>QTY LIMIT:</i> 500 mg = 3 tablets/day, 1000 mg = 2 tablets/day	Aspruzyo Sprinkle™ (ranolazine) granule <i>QTY LIMIT:</i> 500 mg = 3 packets/day, 1000 mg = 2 packets/day BiDil® (isosorbide dinitrate/hydralazine) Dilatrate-SR® (isosorbide dinitrate SR capsule) Isosorbide dinitrate SL tablet Isordil® (isosorbide dinitrate tablet) Nitrolingual Pump Spray® Ranexa® (ranolazine) <i>QTY LIMIT:</i> 500 mg = 3 tablets/day, 1000 mg = 2 tablets/day	<p>Aspruzyo: the patient has medical necessity for a non-solid oral dosage form.</p> <p>Dilatrate-SR, Isosorbide dinitrate SL tablet, Isordil: the patient has had a side effect, allergy, or treatment failure to at least two preferred agents.</p> <p>Nitrolingual Pump Spray: the patient has had a side effect, allergy, or treatment failure to Nitroglycerin spray lingual.</p> <p>Bidil: The prescriber provides a clinically valid reason why the patient cannot use isosorbide dinitrate and hydralazine as separate agents.</p> <p>Ranexa: the patient has a documented intolerance to the generic equivalent.</p>
TOPICAL		
NITRO-BID® (nitroglycerin ointment) NITROGLYCERIN TRANSDERMAL PATCHES (compare to Nitro-Dur®)	Nitro-Dur® (nitroglycerin transdermal patch)	<p>Nitro-Dur: patient has had a side effect, allergy, or treatment failure to generic nitroglycerin transdermal patches.</p>
SINUS NODE INHIBITORS		
All products require a PA	Corlanor® (ivabradine) <i>QTY LIMIT:</i> 60 tabs/30 days	<p>Corlanor Clinical Criteria:</p> <p><i>Diagnosis of stable, symptomatic heart failure:</i></p> <ul style="list-style-type: none"> • Left ventricular ejection fraction of $\leq 35\%$ AND • Resting heart rate ≥ 70 bpm AND • In sinus rhythm AND • Patient has persisting symptoms despite maximally tolerated doses of beta blockers or who have contraindication to beta blocker therapy <p><i>Diagnosis of Inappropriate Sinus Tachycardia:</i></p> <ul style="list-style-type: none"> • Patient has persisting symptoms despite maximally tolerated doses of beta blockers or there is a contraindication to beta blocker therapy.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

Diagnosis of Postural Orthostatic Tachycardia Syndrome (POTS)

- The patient has a documented side effect, allergy, or treatment failure with at least 2 of the following medications: fludrocortisone, midodrine, beta blocker (metoprolol or propranolol), or pyridostigmine.

CORTICOSTEROIDS: ORAL

DEXAMETHASONE tablets, elixir, intensol, solution
 DEXPAK[®] tabs (dexamethasone taper pack)
 HYDROCORTISONE tab (compare to Cortef[®])
 MEDROL[®] (methylprednisolone) 2mg tablets
 METHYLPREDNISOLONE (compare to Medrol[®]) tabs
 METHYLPREDNISOLONE DOSE PACK (compare to Medrol Dose Pack[®]) tabs
 PREDNISOLONE 3 mg/ml oral solution, syrup
 PREDNISOLONE SODIUM PHOSPHATE 3 mg/ml oral solution (compare to Orapred[®])
 PREDNISOLONE SOD PHOSPHATE ORAL SOLUTION 6.7mg/5ml (5mg/5ml base) (compare to Pediapred[®])
 PREDNISONE intensol, solution, tablets

Alkindi[®] Sprinkle (hydrocortisone) granule
 Cortef[®] (hydrocortisone) tablets
 Hemady[®] (dexamethasone) tablets
 Medrol[®] (methylprednisolone) tablets
 Medrol Dose Pak[®] (methylprednisolone) tabs
 Prednisolone sodium phosphate oral solution 25 mg/5ml
 Rayos[®] (prednisone) Delayed Release Tablet
QTY LIMIT: 1 tablet/day

Rayos: The patient has had a trial of generic immediate release prednisone and has documented side effects that are associated with the later onset of activity of immediate release prednisone taken in the morning.
All Others: The patient has a documented side effect, allergy, or treatment failure to at least two preferred medications. If a product has an AB rated generic, one trial must be the generic formulation.

COUGH AND COLD PREPARATIONS

Please refer to the DVHA website for covered OTC cough & cold products
https://dvha.vermont.gov/sites/dvha/files/documents/OTCWebList_0.pdf
 All RX generics

Note: The FDA restricts the use of prescription codeine pain and cough medicines in children. Prior authorization is required for patients <12 years of age.

Hydrocodone/chlorpheniramine (compare to Tussionex[®])
QTY LIMIT: 60 ml/RX
 Tussionex[®] (hydrocodone/chlorpheniramine)
QTY LIMIT: 60 ml/RX
 TussiCaps[®] (hydrocodone/chlorpheniramine)
QTY LIMIT: 12 capsules/RX
 All other brands

Tussionex, TussiCaps, Hydrocodone/chlorpheniramine suspension (generic): The patient has had a documented side effect, allergy, or treatment failure to two of the following generically available cough or cough/cold products: hydrocodone/homatropine (compare to Hycodan), promethazine/codeine (previously Phenergan with Codeine), guaifenesin/codeine (Cheratussin AC) or benzonatate. AND patient is 6 years old of age or greater. AND The quantity requested does not exceed 60 ml (Tussionex) or 12 capsules (TussiCaps). AND If the request is for Tussionex, the patient has a documented intolerance to generic hydrocodone/chlorpheniramine suspension.
All Other Brands: The prescriber must provide a clinically valid reason for the use of the requested medication including reasons why any of the generically available preparations would not be a suitable alternative.

CYSTIC FIBROSIS MEDICATIONS

Preferred After Clinical Criteria Are Met

KITABIS[®] (tobramycin sol)

QTY LIMIT: 56 vials/56 days; maximum day supply = 56 days (2 vials/day for 28 days, then 28 days off)

TOBI[®] Podhaler (tobramycin capsules for inhalation)

QTY LIMIT: 224 capsules/56 days; maximum day supply = 56 days (4 capsules twice daily for 28 days, then 28 days off)

TOBRAMYCIN inhalation solution (compare to Tobi[®]) 300mg/5mL

QTY LIMIT: 56 vials/56 days; maximum day supply = 56 days (2 vials/day for 28 days, then 28 days off)

Bethkis[®] (tobramycin) inhalation solution

QTY LIMIT: 56 vials/56 days; maximum day supply = 56 days (2 vials/day for 28 days, then 28 days off)

Bronchitol[®] (mannitol) capsules for inhalation

QTY LIMIT: 560 capsules/28 days; maximum day supply = 28 days

Cayston[®] (aztreonam) inhalation solution

QTY LIMIT: 84 vials/56 days; maximum day supply = 56 days (3 vials/day for 28 days, then 28 days off)

Kalydeco[®] (ivacaftor) tablets

QTY LIMIT: 2 tablets/day, maximum day supply = 30 days

Kalydeco[®] (ivacaftor) packets

QTY LIMIT: 2 packets/day; maximum day supply = 30 days

Orkambi[®] (lumacaftor/ivacaftor)

QTY LIMIT: 120/30 days; maximum day supply=30 days

Pulmozyme[®] (dornase alfa) inhalation solution

QTY LIMIT: 60/30 days; maximum day supply=30 days

Symdeko[®] (tezacaftor/ivacaftor and ivacaftor)

QTY LIMIT: 56/28 days; maximum day supply = 28 days

Tobi[®] (tobramycin) inhalation solution

QTY LIMIT: 56 vials/56 days; maximum day supply = 56 days (2 vials/day for 28 days, then 28 days off)

Tobramycin inhalation solution 300mg/4mL

QTY LIMIT: 56 vials/56 days; maximum day supply = 56 days (2 vials/day for 28 days, then 28 days off)

Trikafta[®] (elexacaftor/tezacaftor/ivacaftor)

QTY LIMIT: 84/28 days; maximum day supply = 28 days

Kitabis, Tobramycin inhalation solution (300mg/5mL), Pulmozyme: diagnosis or indication is cystic fibrosis

Bethkis, TOBI, tobramycin inhalation solutions (300mg/4mL): Diagnosis or indication is cystic fibrosis and the patient has a documented failure or intolerance to two preferred formulations of tobramycin inhalation solution.

Bronchitol: Diagnosis or indication is cystic fibrosis AND the patient is 18 years of age or older AND the patient has a documented inadequate response or contraindication to hypertonic saline and Pulmozyme AND the patient has passed the Bronchitol Tolerance Test (BTT) AND the patient has been counseled to use a short-acting beta agonist (SABA) 5-15 minutes prior to each dose.

Cayston: diagnosis or indication is cystic fibrosis and the patient has had a documented failure, intolerance or inadequate response to inhaled tobramycin therapy alone

Kalydeco: The patient has a diagnosis of Cystic Fibrosis AND Patient has a mutation on at least one allele in the cystic fibrosis transmembrane conductance regulator gene (CFTR gene) shown to be responsive to Kalydeco per FDA approval (documentation provided). AND The patient is ≥ 1 month old. Note: Renewal of Prior Authorization will require documentation of member response.

TOBI Podhaler: allowed after a trial of another form of inhaled tobramycin

Orkambi/Symdeko/Trikafta: The patient has a diagnosis of Cystic Fibrosis AND

Initial Criteria

- Patient age is FDA approved for the requested medication AND
- Patient must have a confirmed mutation in the CFTR gene shown to be responsive to the requested medication per FDA approval (documentation provided) AND
- If the patient is under the age of 18, they must have undergone a baseline ophthalmic examination to monitor for lens opacities/cataracts AND
- Prescriber is a CF specialist or pulmonologist

Ongoing Approval Criteria

- Patient has clinically documented improvement in lung function (will be applied to the first renewal request only; requirement waived on subsequent renewals)

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<ul style="list-style-type: none"> • Patient has LFTs/bilirubin monitored every 3 months for the first year of therapy and annually after the first year • ALT or AST ≤ 5 X the upper limit of normal or ALT/AST ≤ 3 X the upper limits of normal and bilirubin is ≤ 2 X the upper limit of normal • For patients under the age of 18, have follow up ophthalmic exam at least annually
DERMATOLOGICAL AGENTS		
ACTINIC KERATOSIS THERAPY		
<p>CARAC[®] (fluorouracil) 0.5% cream FLUOROURACIL (compare to Efudex[®]) 5% cream IMIQUIMOD 5% Cream</p> <p><i>C=cream, F=foam, G=gel, L=lotion, O=ointment, S=solution</i></p>	<p>Aldara[®] (imiquimod) 5 % Cream Diclofenac Sodium 3 % Gel (compare to Solaraze[®]) <i>QTY LIMIT: 1 tube/30 days</i> Efudex[®] (fluorouracil) 5% cream Fluorouracil 5%, 2% solution Fluorouracil (compare to CARAC[®]) 0.5% cream Zyclara (imiquimod) 3.75 % Cream <i>QTY LIMIT: 56 packets/6 weeks</i> Zyclara (imiquimod) 2.5%, 3.75 % Cream Pump <i>QTY LIMIT: 2 pumps/8 weeks</i></p>	<p>Aldara: the patient has a documented intolerance to generic imiquimod 5% cream Efudex cream, Fluorouracil solution: The patient has a documented intolerance to fluorouracil 5% cream. Fluorouracil 0.5% cream: The patient has a documented intolerance to brand Carac. Diclofenac Gel: The diagnosis or indication is actinic keratosis AND The patient has had a documented side effect, allergy, contraindication or treatment failure with a preferred topical fluorouracil product. Zyclara Cream: The diagnosis or indication is actinic keratosis on the face or scalp AND The patient has had a documented side effect, allergy, or treatment failure with 5-fluorouracil and imiquimod 5% cream. OR The treatment area is greater than 25 cm² on the face or scalp. AND The patient has had a documented side effect, allergy, or treatment failure with 5-fluorouracil.</p>
ANTIBIOTICS TOPICAL		
<p><u>SINGLE AGENT</u> BACITRACIN MUPIROCIN OINTMENT (compare to Bactroban[®])</p> <p><u>COMBINATION PRODUCTS</u> BACITRACIN-POLYMYXIN NEOMYCIN-BACITRACIN-POLYMYXIN</p> <p><i>C=cream, F=foam, G=gel, L=lotion, O=ointment, S=solution</i></p>	<p>Centany[®] Ointment (mupirocin) Gentamicin Cream or Ointment Mupirocin cream (compare to Bactroban[®]) Xepi cream (ozenoxacin)</p>	<p>Mupirocin cream, Centany Ointment, Xepi cream: The patient has had a documented intolerance with generic mupirocin ointment Gentamicin Cream or Ointment: The patient has had a documented side-effect, allergy, or treatment failure with at least one preferred generic topical antibiotic</p>
ANTIFUNGALS: ONYCHOMYCOSIS		

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>CICLOPIROX 8% solution <i>QTY LIMIT</i>: 6.6 ml/90 days</p> <p>JUBLIA® (efinaconazole 10% solution) <i>QTY LIMIT</i>: 48 weeks treatment</p> <p>TAVABOROLE 5% solution <i>QTY LIMIT</i>: 48 weeks treatment</p>	<p>Ciclodan® (ciclopirox 8% solution)</p> <p>Kerydin® (tavaborole 5% solution) <i>QTY LIMIT</i>: 48 weeks treatment</p>	<p>Kerydin: Patient has a documented side effect, allergy, or treatment failure to two preferred topical onychomycosis agents, one of which must be tavaborole.</p> <p>Ciclodan: Patient has a documented intolerance to generic ciclopirox 8% solution.</p> <p>LIMITATIONS: Coverage of Onychomycosis agents will NOT be approved solely for cosmetic purposes. Kits with multiple drug products or non-drug items not covered.</p>
ANTIFUNGALS: TOPICAL		
<p><u>SINGLE AGENT</u></p> <p>BUTENAFINE (compare to Mentax®) 1% C</p> <p>CICLOPIROX 0.77% C, Sus, G; 1% Sh</p> <p>CLOTRIMAZOLE 1% C, S</p> <p>ECONAZOLE 1% C</p> <p>KETOCONAZOLE 2% C, 2% Sh</p> <p>MICONAZOLE all generic/OTC products</p> <p>NYSTATIN O, C, P (compare to Mycostatin®), Nystop®, Nyamyc®)</p> <p>TOLNAFTATE 1% C, P, S</p> <p><u>COMBINATION PRODUCTS</u></p> <p>CLOTRIMAZOLE W/BETAMETHASONE C, L</p> <p>NYSTATIN W/TRIAMCINOLONE C, O</p> <p><i>C=cream, F=foam, G=gel, L=lotion, P=powder, S=solution, Sh=shampoo, Sp=spray, Sus=suspension</i></p>	<p>Ertaczo® (sertaconazole) 2% C</p> <p>Extina® (ketoconazole) 2% F</p> <p>Ketoconazole (compare to Extina®) 2% Foam</p> <p>Luliconazole 1% C</p> <p>Luzu® (luliconazole) 1% Cream</p> <p>Mentax® 1% C</p> <p>Naftifine (compare to Naftin®) 1% & 2% C, 1% G</p> <p>Naftin® (naftifine) 1% C, 1%, 2% G</p> <p>Nystop®, Nyamyc® (nystatin) P</p> <p>Oxiconazole 1% C</p> <p>Oxistat® (oxiconazole) 1% L</p> <p>Sulconazole 1% C, L</p> <p>Miconazole w/ zinc oxide (compare to Vusion®) O <i>QTY LIMIT: 50 g/30 days</i></p> <p>Vusion® (miconazole w/zinc oxide) O <i>QTY LIMIT: 50 g/30 days</i></p> <p>All other branded products</p> <p>Note: Please refer to “Dermatological: Antifungals: Onychomycosis” for ciclopirox solution</p>	<p>All Non-Preferred Agents (except Vusion): The patient has had a documented side effect, allergy, or treatment failure to at least TWO different preferred generic topical antifungal agents. (If a product has an AB rated generic, one trial must be the generic equivalent of the requested product.) OR The patient has a contraindication that supports the need for a specific product or dosage form of a brand topical antifungal.</p> <p>Miconazole w/ Zinc Oxide, Vusion: The patient has a diagnosis of diaper dermatitis complicated by documented candidiasis AND The patient is at least 4 weeks of age. AND The patient has had two trials (with two different preferred antifungal agents) used in combination with a zinc oxide diaper rash product resulting in documented side effects, allergy, or treatment failures.</p>
ANTIVIRALS: TOPICAL		
<p>ACYCLOVIR (compare to Zovirax®) 5% O</p> <p>DOCOSANOL 10% C</p> <p><i>C=cream, O=ointment, S = solution</i></p>	<p>Acyclovir (compare to Zovirax®) 5% C</p> <p>Denavir® (penciclovir) 1% C</p> <p>Penciclovir 1% C</p> <p>Xerese® (acyclovir 5%/hydrocortisone 1%) C</p> <p>Ycanth™ (cantharidin) 0.7% S <i>QTY LIMIT</i>: 8 applicators/12 weeks</p> <p>Zovirax® (acyclovir) 5% C, O</p>	<p>Acyclovir cream: The patient has a documented intolerance to acyclovir ointment</p> <p>Denavir, Penciclovir, Xerese: The patient has a treatment failure with a preferred topical acyclovir product. AND for approval of penciclovir, the patient has a documented intolerance to brand Denavir.</p> <p>Ycanth: The patient has a diagnosis of Molluscum Contagiosum AND either cryotherapy or curettage has failed to alleviate severe symptoms AND documentation must be submitted to support continued need if treatment duration exceeds 12 weeks.</p> <p>Zovirax cream, ointment: The patient has a documented intolerance to generic acyclovir ointment</p>

AXILLARY HYPERHIDROSIS THERAPY

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
Xerac-AC (aluminum chloride) 6.25% Solution		
CORTICOSTEROIDS: LOW POTENCY		
<p>ALCLOMETASONE 0.05% C, O DESONIDE 0.05% C, O FLUOCINOLONE 0.01% C, S, oil (compare to Derma-Smoothe, Synalar®) HYDROCORTISONE 0.5%, 1%, 2.5% C; 2.5% L, 0.5%, 1%, 2.5% O</p> <p><i>C=cream, F=foam, G=gel, L=lotion, O=ointment, S=solution</i></p>	<p>Derma-Smoothe® (fluocinolone 0.01%) oil Desonide 0.05% L Synalar® (fluocinolone) 0.01%</p> <p>S All other brands</p>	<p>CRITERIA FOR APPROVAL (NON-PREFERRED AGENTS): The patient has a documented side effect, allergy, or treatment failure to at least two different preferred agents of similar potency. (If a product has an AB rated generic, one trial must be the generic.)</p>
CORTICOSTEROIDS: MEDIUM POTENCY		
<p>BETAMETHASONE DIPROPIONATE 0.05% C, L, O BETAMETHASONE VALERATE 0.1% C, L, O BETAMETHASONE VALERATE 0.12% (compare to Luxiq®) F FLUOCINOLONE 0.025% C, O (compare to Synalar®) FLUTICASONE 0.05% C; 0.005% O HYDROCORTISONE VALERATE 0.2% C, O MOMETASONE FUROATE 0.1% C, L, O, S TRIAMCINOLONE ACETONIDE 0.025%, 0.1% C, L, O</p> <p><i>C=cream, F=foam, G=gel, L=lotion, O=ointment, S=solution</i></p>	<p>Clocortolone 0.1% C (compare to Cloderm®) Cloderm® (clocortolone) 0.1% C Desoximetasone 0.05% C, O (compare to Topicort®) Flurandrenolide C, L, O Fluticasone 0.05%, L Hydrocortisone Butyrate 0.1% C, O, S Kenalog® (triamcinolone) Aerosol Spray Luxiq® (betamethasone valerate) F Prednicarbate 0.1% C, O Synalar® (fluocinolone) 0.025% C, O Topicort® (desoximetasone) 0.05% C, O Triamcinolone Aerosol Spray Trianex® (triamcinolone) 0.05%</p> <p>O All other brands</p>	<p>CRITERIA FOR APPROVAL (NON-PREFERRED AGENTS): The patient has a documented side effect, allergy, or treatment failure to at least two different preferred agents of similar potency. (If a product has an AB rated generic, one trial must be the generic.)</p>
CORTICOSTEROIDS: HIGH POTENCY		
<p>AUGMENTED BETAMETHASONE 0.05% C, L (compare to Diprolene® AF) BETAMETHASONE VALERATE 0.1% C, O DESOXIMETASONE 0.25% C, O (compare to Topicort®) FLUOCINONIDE 0.05% C, G, O, TRIAMCINOLONE ACETONIDE 0.5% C, O</p> <p><i>C=cream, F=foam, G=gel, L=lotion, O=ointment, S=solution</i></p>	<p>Apexicon E® (diflorasone) 0.05% C Desoximetasone 0.05% G Diflorasone diacetate 0.05% C, O (compare to Apexicon E®) Halcinonide 0.1% C Halog® (halcinonide) all products</p> <p>Topicort® (desoximetasone) 0.05% G; 0.25% C, O, Spray</p> <p>All other brands</p>	<p>CRITERIA FOR APPROVAL (NON-PREFERRED AGENTS): The patient has a documented side effect, allergy, or treatment failure to at least two different preferred agents of similar potency. (If a product has an AB rated generic, one trial must be the generic.)</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
CORTICOSTEROIDS: VERY HIGH POTENCY		
<p>AUGMENTED BETAMETHASONE 0.05% C, L, O (compare to Diprolene®) 0.05% G CLOBETASOL PROPIONATE 0.05%, C, F, G, L, O, S, Shampoo, Spray HALOBETASOL PROPIONATE (compare to Ultravate®) 0.05% C, O <i>C=cream, F=foam, G=gel, L=lotion, O=ointment, S=solution</i></p>	<p>Bryhali® (halobetasol propionate) L Clobetasol propionate emulsion (compare to Olux E®) 0.05% F Diprolene® (augmented betamethasone) 0.05% L, O Fluocinonide (compare to Vanos®)0.1% C Halobetasol (compare to Lexette™) 0.05% F Impeklo™ (clobetasol propionate) 0.05% L Lexette™ (halobetasol) 0.05% F Olux®/Olux E® (clobetasol propionate) 0.05% F Tovet® (clobetasol propionate aerosol) 0.05% F Vanos® (fluocinonide) 0.1% C All other brands</p>	<p>CRITERIA FOR APPROVAL (NON-PREFERRED AGENTS): The patient has a documented side effect, allergy, or treatment failure to at least two different preferred agents of similar potency. (If a product has an AB rated generic, one trial must be the generic.)</p>
GENITAL WART THERAPY		
<p>IMIQUIMOD 5 % (compare to Aldara®) cream PODOFILOX SOLUTION (compare to Condylox®)</p>	<p>Aldara® (imiquimod) 5% cream Condylox® Gel (podofilox gel) Imiquimod (compare to Zyclara®) 3.75% Cream QTY Limit: 56 packets/8 weeks Imiquimod (compare to Zyclara®) 3.75% Cream Pump <i>QTY LIMIT: 2 pumps/ 8 weeks</i> Veregan® (sinecatechins ointment) <i>QTY LIMIT: 15 grams (1 tube)/30 days</i> Zyclara® (imiquimod 3.75%) Cream <i>QTY LIMIT: 56 packets/8 weeks</i> Zyclara® (imiquimod 2.5%, 3.75%) Cream Pump <i>QTY LIMIT: 2 pumps/8 weeks</i></p>	<p>Aldara cream, Zyclara cream: The patient has had a documented intolerance to generic imiquimod Condylox gel, Veregan: The patient has had a documented side effect, allergy, or treatment failure with imiquimod. Imiquimod pump, Zyclara pump: The patient has had a documented intolerance to generic imiquimod cream and Zyclara cream.</p>
IMMUNOMODULATORS		
<p>ELIDEL® (pimecrolimus) for ages ≥ 2 TACROLIMUS 0.03% Ointment for ages ≥ 2 TACROLIMUS 0.1% Ointment for ages ≥ 16</p>	<p>Cibinqo® (abrocitinib) tablets QTY LIMIT: 1 tab/day Maximum 30 days supply Eucrisa® (crisaborole) Ointment</p>	<p>Eucrisa: The patient has a diagnosis of mild-moderate atopic dermatitis (eczema) AND the patient has had a documented side effect, allergy, or treatment failure (defined as daily treatment for at least one month) with at least one preferred topical calcineurin inhibitor AND the quantity requested does not exceed 60 grams/fill and 180 grams/ 6 months. Trial of calcineurin</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>Preferred After Clinical Criteria Are Met</u></p> <p>ADBRY (tralokinumab-ldrm) subcutaneous injection <i>QTY LIMIT:</i> 6 syringes the first 28 days then 4 syringes every 28 days thereafter</p> <p>DUPIXENT® (dupilumab) subcutaneous injection <i>QTY LIMIT:</i> 4 syringes/pens the first 28 days then 2 Syringes/pens every 28 days thereafter</p> <p>OPZELURA® (ruxolitinub) cream QTY Limit: 8 tubes (60 g) for 8 weeks of therapy</p> <p><i>Note:</i> please refer to <i>Dermatological Agents: Corticosteroids</i> category for preferred topical corticosteroids.</p>	<p>Pimecrolimus cream (compare to Elidel®) Rinvoq® (upadactinib) extended-release tablet <i>QTY LIMIT:</i> 1 tablet/day Maximum 30 days supply</p>	<p>inhibitor will be waived for patients ≥ 3 months through < 2 years of age.</p> <p>Opzelura:</p> <ul style="list-style-type: none"> • The patient is ≥ 12 years of age AND • The patient has a diagnosis of mild-moderate atopic dermatitis (eczema) AND • The patient has had a documented side effect, allergy, or treatment failure (defined as daily treatment for at least one month) with at least one moderate to high potency topical corticosteroid within the last 6 months, unless contraindicated AND • The patient has had a documented side effect, allergy, or treatment failure (defined as daily treatment for at least one month) of a preferred topical calcineurin inhibitor and crisabarole ointment AND • Patient is not receiving Opzelura in combination with another biologic medication (e.g. dupilumab), oral JAK inhibitor (e.g. upadactinib), or systemic immunosuppressant (e.g. cyclosporine) AND • The quantity requested does not exceed 60 grams/fill; maximum of 8-weeks of continuous use. • Request will be approved for 8 weeks of therapy per calendar year <p>Pimecrolimus: The patient has a documented intolerance to brand Elidel.</p> <p>Adbry, Cibinqo, Dupixent, Rinvoq:</p> <ul style="list-style-type: none"> • The patient's age is FDA approved for the given indication AND • The patient has a diagnosis of moderate to severe atopic dermatitis AND • The prescription is initiated in consultation with a dermatologist, allergist, or immunologist AND • At least 10% of the body's surface area is involved AND • The patient has had a documented side effect, allergy, or treatment failure (defined as daily treatment for at least one month) with at least one moderate to high potency topical corticosteroid and one preferred topical calcineurin inhibitor within the last 6 months AND • Initial approval will be granted for 6 months. For re-approval after 6 months, the prescriber must submit documentation of clinical improvement in symptoms. Renewals may be granted for up to 1 year. <p>Cibinqo additional criteria: The patient has had a documented side effect, allergy, or treatment failure with Adbry or Dupixent AND the patient has had a documented side effect, allergy, or treatment failure with Rinvoq.</p> <p>Rinvoq additional criteria: The patient has had a documented side effect, allergy, or treatment failure with Adbry or Dupixent.</p> <p>Dupixent: Diagnosis is Prurigo Nodularis:</p> <ul style="list-style-type: none"> • The patient must be 18 years of age or older AND • Diagnosis is confirmed based on the following: chronic pruritic lasting ≥ 6 weeks, history and/or signs of repeated scratching, and multiple localized or generalized pruriginous skin lesion (e.g. whitish or pink papules, nodules and/or plaques) AND

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<ul style="list-style-type: none"> The patient has had a documented side effect, allergy, or treatment failure (defined as daily treatment for at least one month) with at least one moderate to high potency topical corticosteroid and one preferred topical calcineurin inhibitor within the last 6 months For continuation of therapy after the initial 6-month authorization, there must be documented improvement in PN symptoms
SCABICIDES AND PEDICULOCIDES		
PERMETHRIN 5 % (compare to Elimite®) C PERMETHRIN 1 % CR, L	Crotan 10% L Ivermectin 0.5% L	Non-preferred Scabicides: The patient has had a documented side effect or allergy to permethrin cream and Natroba or treatment failure with two treatments of permethrin cream and Natroba.
PIPERONYL BUTOXIDE AND PYRETHRINS G, S, Sh NATROBA® (spinosad 0.9 %) Ss <i>C=cream, CR=crème rinse, G=gel, L=lotion, S=solution, Sh=shampoo, Sp=spray, Ss=suspension</i>	Malathion L (compare to Ovide®) Ovide® (malathion) L Spinosad (compare to Natroba) Ss Vanalice® (piperonyl butoxide/pyrethrins) G	Non-Preferred Pediculicides: The patient has had a documented side effect or allergy to OTC permethrin and piperonyl butoxide and pyrethrins and one treatment of Natroba OR treatment failure with two treatments of OTC permethrin and/or piperonyl butoxide and pyrethrins and one treatment of Natroba. For approval of Ovide® Lotion, the patient must also have a documented intolerance to the generic equivalent product.
WOUND CARE		
All products require PA	Filsuvez® (birch triterpenes) gel Vyjuvek® (beremagene geperpavec-svdt)	Filsuvez; <ul style="list-style-type: none"> The patient has a diagnosis of dystrophic or junctional epidermolysis bullosa AND The patient is at least 6 months old AND The patient does not have current evidence or history of squamous cell carcinoma or active infection in the area requiring Filsuvez application. AND The patient has used standard wound care treatments, including silicone or foam dressings without wound resolution AND Initial approval will be granted for 6 months. For reapproval, the patient must have a documented reduction in the number of wounds, decrease in wound size, increase in granulation tissue, or complete wound closure. Vyjuvek: <ul style="list-style-type: none"> The patient has a diagnosis of dystrophic epidermolysis bullosa (DEB) with confirmed mutation(s) in the collagen type VII alpha 1 chain (COL7A1) gene AND The patient does not have current evidence or history of squamous cell carcinoma or active infection in the area requiring Vyjuvek application. AND

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<ul style="list-style-type: none"> • The patient has used standard wound care treatments, including silicone or foam dressings without wound resolution AND has tried and failed Filsuvez • The intended wounds cover a large area of the patient’s body OR are likely to present for an extended duration (e.g. months of healing) AND • The intended wounds have presented significant detrimental health consequences to the patient and are unlikely to be resolved with standard wound care management (e.g. hospitalizations, frequent professional wound care management, significant quality of life disruptions) AND • Approval will be limited to a maximum of 10 mL per 28 days • Initial approval will be granted for 6 months. For reapproval, the patient must have a documented reduction in the number of wounds, decrease in wound size, increase in granulation tissue, or complete wound closure
DESMOPRESSIN: INTRANASAL/ORAL		
<p><u>INTRANASAL</u> All products require PA</p> <p><u>ORAL</u> DESMOPRESSIN</p>	<p>DDAVP[®] (desmopressin) Nasal Solution or Spray 0.01% Desmopressin Nasal Solution or Spray 0.01 % (compare to DDAVP[®]) Noctiva[™] (desmopressin) Nasal Spray Stimate[®] (desmopressin) Nasal Solution 1.5 mg/ml</p> <p>Nocdurna[®] (desmopressin) SL tablets <i>QTY LIMIT:</i> 1 tablet/day DDAVP[®] (desmopressin) tablets</p>	<p>CRITERIA FOR APPROVAL:</p> <p>Intranasal (except as indicated below): The diagnosis or indication for the requested medication is (1) Diabetes Insipidus, (2) hemophilia type A, or (3) Von Willebrand disease AND If the request is for brand DDAVP, the patient has a documented intolerance to generic desmopressin spray or solution.</p> <p>Oral: The diagnosis or indication for the requested medication is (1) Diabetes Insipidus and/or (2) primary nocturnal enuresis AND The patient has had a documented intolerance to generic desmopressin tablets</p> <p>Nocdurna, Noctiva: Patient is ≥18 years of age (Nocdurna) or ≥50 years of age (Noctiva) AND the indication for use is the treatment of nocturia due to nocturnal polyuria (defined as nighttime urine production exceeding 1/3 of the 24-hour urine production) causing patient to awaken more than 2 times per night to void for at least 6 months AND patient has eGFR > 50ml/min/1.73m² AND patient does not have increased risk of severe hyponatremia (e.g. concomitant use of loop diuretics or corticosteroids, diagnosis of CHF, or uncontrolled hypertension) AND serum sodium concentrations are normal before starting therapy AND patient has had a documented intolerance to generic desmopressin tablets.</p> <p>LIMITATIONS: Desmopressin intranasal formulations will not be approved for the treatment of primary nocturnal enuresis (PNE) due to safety risks of hyponatremia. Oral tablets may be prescribed for this indication.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

DIABETIC TESTING SUPPLIES

<p>Please refer to the DVHA website for covered Diabetic testing supplies. Test strips are subject to a quantity limit of 200 strips per 30 days. https://dvha.vermont.gov/sites/dvha/files/doc_library/Vermont%20PDSL%20August%202023.pdf</p>		<p>CRITERIA FOR APPROVAL: The prescriber demonstrates that the patient has a medical necessity for clinically significant features that are not available on any of the preferred meters/test strips. CRITERIA FOR APPROVAL to Exceed QTY LIMIT: Chart notes must be provided documenting medical necessity. LIMITATIONS: Talking monitors are not covered under the pharmacy benefit.</p>
--	--	---

ENDOMETRIOSIS/UTERINE FIBROIDS AGENTS

<p>LUPRON DEPOT® (leuprolide acetate for depot suspension) <i>QTY LIMIT:</i> 3.75 mg kit/month or 11.25 mg kit/3 months SYNAREL® (nafarelin acetate) nasal solution <u>Preferred After Clinical Criteria are Met</u> MYFEMBREE® (relugolix/estradiol/norethindrone) tablet <i>QTY LIMIT:</i> 1 tab/day Orilissa® (elagolix) tablets <i>QTY LIMIT:</i> 200mg dose = 2 tabs/day; maximum of 6 months; 150mg = 1 tab/day</p>	<p>Lupaneta Pack™ (leuprolide acetate for depot suspension and norethindrone acetate tablets) <i>QTY LIMIT:</i> 3.75 mg kit/month or 11.25 mg kit/3 months Oriahnn® (elagolix and elagolix/estradiol/norethindrone) capsules <i>QTY LIMIT:</i> 2 caps/day</p>	<p>Lupaneta Pack: patient has a documented intolerance to Lupron Depot and norethindrone tablets used in combination. Myfembree, Orilissa: Patient has a documented side effect, allergy, or treatment failure to at least TWO medications from at least 2 different classes (oral contraceptives, NSAIDs, progestins). Note: Use of GnRH receptor antagonists will be limited to 2 years. Oriahnn: Patient has a documented side effect, allergy, or treatment failure to at least TWO medications from at least 2 different classes (oral contraceptives, NSAIDs, progestins) AND the patient has a documented side effect, allergy, or treatment failure with Myfembree or Orilissa. Note: Use of GnRH receptor antagonists will be limited to 2 years.</p>
---	--	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

EPINEPHRINE: SELF-ADMINISTERED

<p>EPIPEN-JR INJ 0.15mg EPIPEN INJ 0.3mg EPINEPHRINE INJ (compare to EpiPen-Jr®) (authorized generic, Mylan labeler code 49502 is the only preferred form) 0.15mg EPINEPHRINE INJ (compare to EpiPen®) (authorized generic, Mylan labeler code 49502 is the only preferred form) 0.3mg</p>	<p>Auvi-Q® Inj 0.1mg Auvi-Q® Inj 0.15mg Auvi-Q® Inj 0.3mg Epinephrine Inj 0.15 mg Epinephrine Inj 0.3 mg Symjepi® Inj 0.15mg Symjepi® Inj 0.3mg</p>	<p>Non-preferred Agents (0.15mg, 0.3mg): The patient must have a documented intolerance to a preferred epinephrine product. Auvi-Q 0.1mg: Patient weight is 7.5kg to 15kg (16.5 to 33 lbs).</p>
---	---	--

ESTROGENS: VAGINAL

<p><u>ESTRADIOL</u> ESTRACE VAGINAL® Cream ESTRING® Vaginal Ring VAGIFEM® Vaginal Tablets</p> <p><u>CONJUGATED ESTROGENS</u> PREMARIN VAGINAL® Cream</p> <p><u>ESTRADIOL ACETATE</u> FEMRING® Vaginal Ring</p>		
--	--	--

GASTROINTESTINAL

INFLAMMATORY BOWEL DISEASE BIOLOGICS: Initial approval is 3 months; renewals are 1 year

Preferred After Clinical Criteria Are Met
INJECTABLE

ADALIMUMAB-ADBM (compare to Cyltezo®) biosimilar to Humira®

AVSOLA® (infliximab-axxq) biosimilar to Remicade®

HUMIRA® (adalimumab)

QTY LIMIT: 6 syringes/28 days for the first month (Crohn's starter kit); 2 syringes/28 days subsequently

INFLECTRA® (infliximab-dyyb) biosimilar to Remicade®

SKYRIZI® (risankizumab-rzaa)

QTY LIMIT: 360 mg (2.4ml)/56 days after initial IV loading dose

Abrilada™ (adalimumab-afzb) biosimilar to Humira®
Adalimumab-adaz (compare to Hyrimoz®) biosimilar to Humira®

Adalimumab-fkjp (compare to Hulio®) biosimilar to Humira®

Amjevita™ (adalimumab-atto) biosimilar to Humira®

Cimzia® (certolizumab pegol)

QTY LIMIT: 1 kit/28 days

Cyltezo® (adalimumab-adbm) biosimilar to Humira®

Entyvio® (vedolizumab)

IV: QTY LIMIT: 300 mg X 3/42 days, 300 mg X 1 every 56 days thereafter;

Subcutaneous: QTY LIMIT: 216 mg (2 Pens) x 28 days

Hadlima™ (adalimumab-bwvd) biosimilar to Humira®

Hulio® (adalimumab-fkjp) biosimilar to Humira®

Hyrimoz® (adalimumab-adaz) biosimilar to Humira®

Idacio® (adalimumab-aacf) biosimilar to Humira®

OmvoH™ (mirikizumab-mrzk)

QTY LIMIT: 200 mg (2ml) prefilled syringe or autoinjector/28 days after initial IV loading dose

Remicade® (infliximab)

Renflexis™ (infliximab-abda) biosimilar to Remicade®

Simlandi® (adalimumab-ryvk) biosimilar to Humira®

Simponi® (golimumab) SC

QTY LIMIT: 3 of 100 mg prefilled syringe or autoinjector X 1, then 100 mg/28days

Stelara® (ustekinumab)

QTY LIMIT: 90mg (1 mL)/56 days after initial IV loading dose

Tremfya® (guselkumab)

QTY LIMIT: 200 mg IV through 8 weeks, then max dose 200 mg subcutaneous/28 days

Tysabri® (natalizumab)

Yuflyma® (adalimumab-aaty) biosimilar to Humira®

Yusimry™ (adalimumab-aqvh) biosimilar to Humira®

Clinical Criteria for approval of ALL drugs (Crohn's Disease): Patient has a diagnosis of moderate to severe Crohn's disease and has already been stabilized on the medication OR patient meets additional criteria outlined below:

Avsola, Humira, Inflectra, Skyrizi: The patient has had a treatment failure with at least one conventional agent (e.g. methotrexate, corticosteroids) OR there is evidence of severely active disease and early introduction of a biologic without prior medication trials is medically necessary.

Cimzia, Entyvio, Simponi, Stelara, Tysabri: The patient never responded to a 12-week course of anti-TNFα therapy (primary nonresponse) OR the patient previously responded to infliximab (secondary nonresponse) and has a documented side effect, allergy, or treatment failure with adalimumab.

Stelara Note: Initial IV dose for Stelara will be approved through the medical benefit. All subsequent subcutaneous doses may be approved through the pharmacy benefit with quantity limit of 90 mg every 8 weeks.

Note: For maintenance regimens outside of FDA approved dosing intervals, including monthly dosing intervals, clinical notes must include supporting evidence of drug failure at standard dosing intervals and clinical justification for shortened dosing interval. Approval will be granted for 6 months. For renewal the patient must show increased clinical benefit with shorter dosing interval.

Humira Biosimilars: The patient must be unable to use Humira.

Remicade, Renflexis, Zymfentra: The patient must be unable to use Avsola or Inflectra. **For approval of Zymfentra, the patient must have had a documented side effect, allergy, or treatment failure with Humira**

Rinvoq: The patient has had a treatment failure with at least one conventional agent (e.g. 5-ASA, corticosteroids) AND the patient has a documented side effect, allergy, or treatment failure with a preferred TNF inhibitor.

Clinical Criteria for approval of ALL drugs (Ulcerative Colitis): Patient has a diagnosis of moderate to severe Ulcerative Colitis and has already been stabilized on the medication OR patient meets additional criteria outlined below:

Avsola, Humira, Inflectra, Skyrizi: The patient has had a treatment failure with at least one conventional agent (e.g. 5-ASA, corticosteroids) OR there is evidence of severely active disease and early introduction of a biologic without prior medication trials is medically necessary.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

<p>ORAL</p> <p>XELJANZ® (tofacitinib) tablet <i>QTY LIMIT:</i> 2 tablets/day</p> <p>XELJANZ® XR (tofacitinib) tablet <i>QTY LIMIT:</i> 1 tablet/day</p> <p>XELJANZ® (tofacitinib) oral solution</p>	<p>Zymfentra™ (infliximab-dyyb) <i>QTY LIMIT:</i> 240 mg (2ml) prefilled syringe or pen/28 days</p> <p>Rinvoq® (upadactinib) extended-release tablet <i>QTY LIMIT:</i> 1 tablet/day Maximum 30 days supply</p> <p>Velsipity® (etrasimod) tablets <i>QTY LIMIT:</i> 1 tablet/day Maximum 30 days supply</p> <p>Zeposia® (ozanimod) capsule <i>QTY LIMIT:</i> 1 capsule/day</p>	<p>Entyvio, Omvoh, Simponi, Stelara, Tremfya, Velsipity, Zeposia: The patient has had a treatment failure with at least one conventional agent (e.g. 5-ASA, corticosteroids) AND the patient has a documented side effect, allergy, or treatment failure with at least one preferred biologic. Velsipity Note: for approval of Velsipity, the patient must have a documented side effect, allergy, contraindication or treatment failure with Zeposia. Stelara Note: Initial IV dose for Stelara will be approved through the medical benefit. All subsequent subcutaneous doses may be approved through the pharmacy benefit with quantity limit of 90 mg every 8 weeks.</p> <p>Note: For maintenance regimens outside of FDA approved dosing intervals, including monthly dosing intervals, clinical notes must include supporting evidence of drug failure at standard dosing intervals and clinical justification for shortened dosing interval. Approval will be granted for 6 months. For renewal, the patient must show increased clinical benefit with shorter dosing interval.</p> <p>Humira Biosimilars: The patient must be unable to use Humira.</p> <p>Remicade, Renflexis, Zymfentra: The patient must be unable to use Avsola or Inflectra. For approval of Zymfentra, the patient must have had a documented side effect, allergy, or treatment failure with Humira</p> <p>Rinvoq: The patient has had a treatment failure with at least one conventional agent (e.g. 5-ASA, corticosteroids) AND the patient has a documented side effect, allergy, or treatment failure with a preferred TNF inhibitor AND the patient has a documented side effect, allergy, or treatment failure with Xeljanz or Xeljanz XR.</p> <p>Xeljanz, Xeljanz XR: The patient has had a treatment failure with at least one conventional agent (e.g. 5-ASA, corticosteroids) AND the patient has a documented side effect, allergy, or treatment failure with at least one preferred TNF Inhibitor. Note: Induction of Xeljanz 10mg twice daily or XR 22mg once daily will be limited to 16 weeks. Treatment should be discontinued after 16 weeks if adequate therapeutic response is not achieved. For patients with loss of response during maintenance treatment with 5mg twice daily or XR 11mg once daily, approval of 10mg twice daily or XR 22mg once daily will be considered and limited to the shortest duration possible.</p>
H. PYLORI COMBINATION THERAPY		
<p>PYLERA® (bismuth subcitrate, metronidazole, tetracycline) capsules <i>QTY LIMIT:</i> 120 caps/10 days</p>	<p>Bismuth Subcitrate, Metronidazole, Tetracycline (compare to Pylera®) <i>QTY LIMIT:</i> 120 caps/10 days</p> <p>Lansoprazole, Amoxicillin, Clarithromycin <i>QTY LIMIT:</i> 112 caps & tabs/14 days</p> <p>Omeclamox-Pak® (omeprazole, clarithromycin, amoxicillin)</p>	<p>CRITERIA FOR APPROVAL: The patient has a documented treatment failure with Pylera used in combination with a PPI.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
	<p><i>QTY LIMIT:</i> 80 caps & tabs/10 days Talicia® (omeprazole, amoxicillin, rifabutin) delayed release capsules <i>QTY LIMIT:</i> 168 caps/14 days Voquezna® Dual Pack (vonoprazan, amoxicillin) <i>QTY LIMIT:</i> 112 caps & tabs/14 days Voquezna® Triple Pack (vonoprazan, amoxicillin, clarithromycin) <i>QTY LIMIT:</i> 112 caps & tabs/14 days</p>	
H-2 BLOCKERS		
<p>FAMOTIDINE (compare to Pepcid®) tablet</p> <p><u>SYRUPS AND SPECIAL DOSAGE FORMS</u> FAMOTIDINE oral suspension (compare to Pepcid®) age ≤ 12 years</p>	<p>Cimetidine (compare to Tagamet®) tablet Nizatidine capsule Pepcid® (famotidine) tablet</p> <p>Famotidine (compare to Pepcid®) oral suspension (age >12 years)</p>	<p>Cimetidine tablet, Nizatidine capsule, Pepcid tablet: The patient has had a documented side effect, allergy, or treatment failure to famotidine.</p> <p>Famotidine Oral Suspension (Age >12): Patient has a medical necessity for a liquid dosage form.</p>
INFLAMMATORY BOWEL AGENTS (ORAL & RECTAL PRODUCTS)		
<u>MESALAMINE PRODUCTS</u>		
<u>ORAL</u>		
<p>APRISO® (mesalamine capsule extended release) LIALDA® (mesalamine tablet extended release) PENTASA ER® (mesalamine cap CR)</p>	<p>Delzicol® (mesalamine capsule delayed-release) <i>QTY LIMIT:</i> 6 capsules/day Mesalamine capsule delayed release (compare to Delzicol®) <i>QTY LIMIT:</i> 6 capsules/day Mesalamine capsule extended release 0.375gm (compare to Apriso®) Mesalamine tablet delayed release (compare to Asacol® HD) Mesalamine tablet extended release 1.2 g (compare to Lialda®)</p>	<p>Azulfidine, Colazal: patient has had a documented intolerance to the generic equivalent of the requested medication. Budesonide ER 9mg, Ortikos: the patient has a documented intolerance to brand-name Uceris. Delzicol, Mesalamine capsule DR, Mesalamine tablet DR, Mesalamine tablet ER: The patient has had a documented side effect, allergy, or treatment failure to 2 preferred oral mesalamine products. sfRowasa, Uceris Rectal Foam: The patient has had a documented intolerance to mesalamine enema or suppositories. LIMITATIONS: Kits with non-drug products are not covered.</p>
<u>RECTAL</u>		
<p>MESALAMINE ENEMA (compare to Rowasa®) MESALAMINE SUPPOSITORY</p>	<p>sfRowasa® (mesalamine enema sulfite free)</p>	
<u>CORTICOSTEROIDS</u>		
<u>ORAL</u>		
<p>BUDESONIDE 24HR <i>QTY LIMIT:</i> 3 capsules/day UCERIS® (budesonide) ER Tablet <i>QTY LIMIT = 1 tablet/day</i></p>	<p>Budesonide ER 9 mg tablet (compare to Uceris®) <i>QTY LIMIT:</i> 1 tablet/day</p>	

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>RECTAL</u> All products require PA</p> <p><u>OTHER</u></p> <p>BALSALAZIDE (compare to Colazal) DIPENTUM[®] (olsalazine) SULFAZINE SULFAZINE EC SULFASALAZINE (compare to Azulfidine [®]) SULFASALAZINE DR</p>	<p>Ortikos[®] (budesonide) ER capsule QTY LIMIT: 1 capsule/day</p> <p>Uceris[®] Rectal Foam (budesonide)</p> <p>Azulfidine[®] (sulfasalazine) Colazal[®] (balsalazide)</p>	
PROKINETIC AGENTS		
<p><u>TABLETS</u> METOCLOPRAMIDE tabs (compare to Reglan[®])</p> <p><u>ORAL SOLUTION</u> METOCLOPRAMIDE oral solution</p> <p><u>NASAL SPRAY</u> All products require PA</p>	<p>Reglan[®] (metoclopramide)</p> <p>Gimoti[™] (metoclopramide) nasal spray</p>	<p>Reglan: The patient has had a documented intolerance to generic metoclopramide tablets.</p> <p>Gimoti: The patient has a documented intolerance to metoclopramide tablets and oral solution.</p>
PROTON PUMP INHIBITORS		
<p><u>ORAL CAPSULES/TABLETS</u> ESOMEPRAZOLE (compare to Nexium[®]) LANSOPRAZOLE generic RX capsules (compare to Prevacid[®]) OMEPRAZOLE RX capsules (compare to Prilosec[®]) OMEPRAZOLE/SODIUM BICARB capsules (compare to Zegerid[®]) PANTOPRAZOLE tablets (compare to Protonix[®]) ZEGERID RX [®] (omeprazole/sodium bicarb) caps</p>	<p>Aciphex[®] (rabeprazole) tablets QTY LIMIT: 1 tab/day Dexlansoprazole (compare to Dexilant[®]) capsules QTY LIMIT: 1 cap/day Dexilant[®] (dexlansoprazole) capsules QTY LIMIT: 1 cap/day Nexium[®] (esomeprazole) capsules QTY LIMIT: 1 cap/day Omeprazole generic OTC tablets QTY LIMIT: 1 tab/day Omeprazole magnesium generic OTC 20 mg capsules QTY LIMIT: 1 cap/day Prevacid[®] RX (lansoprazole) capsules QTY LIMIT: 1 cap/day Prevacid[®] 24 hr OTC (lansoprazole) capsules QTY LIMIT: 1 cap/day</p>	<p>Lansoprazole ODT, Nexium powder for suspension, Protonix packet (for patients ≥ 12 years old): The patient has a requirement for a non-solid oral dosage form (e.g. an oral liquid, dissolving tablet or sprinkle).</p> <p>Pantoprazole packet, Prevacid Solutabs, Prilosec packet,; The patient has a requirement for a non-solid oral dosage form (e.g. an oral liquid, dissolving tablet or sprinkle). AND the member has had a documented side effect, allergy or treatment failure to two preferred specialty dosage formulations.</p> <p>Dexlansoprazole: The patient has had a documented side effect, allergy, or treatment failure to three preferred PPIs AND the patient has had a documented intolerance to brand Dexilant.</p> <p>Other single-ingredient non-preferred medications: The member has had a documented side effect, allergy, or treatment failure to three preferred PPIs AND if the product has an AB rated generic, there must be a trial of the generic.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
	Protonix® (pantoprazole) tablets <i>QTY LIMIT:</i> 1 tab/day Rabeprazole (compare to Aciphex®) tablets <i>QTY LIMIT:</i> 1 tab/day Voquezna® (vonoprazan) tablets <i>QTY LIMIT:</i> 20 mg tablets = 1 tablet/day for a max of 8 weeks then 10 mg tablets = 1 tablet/day for a max of 6 months	

SUSPENSION & SPECIAL DOSAGE FORMS		
LANSOPRAZOLE ODT (compare to Prevacid Solutab®) (age < 12 years) <i>QTY LIMIT:</i> 1 tab/day NEXIUM® (esomeprazole) powder for suspension (age < 12 years) <i>QTY LIMIT:</i> 1 packet/day PROTONIX® (pantoprazole) packet (age < 12 years) <i>QTY LIMIT:</i> 1 packet/day	Konvomep® (omeprazole/sodium bicarbonate) oral suspension <i>QTY LIMIT:</i> 8 weeks of therapy Nexium® (esomeprazole) powder for suspension (age ≥ 12 years) <i>QTY LIMIT:</i> 1 packet/day Omeprazole/Sodium bicarbonate (compare to Zegerid®) packet for oral suspension <i>QTY LIMIT:</i> 1 packet/day Pantoprazole (compare to Protonix®) packet <i>QTY LIMIT:</i> 1 packet/day Prevacid Solutabs® (lansoprazole) <i>QTY LIMIT:</i> 1 tab/day Prilosec® (omeprazole magnesium) packet <i>QTY LIMIT:</i> 2 packets/day Zegerid RX® (omeprazole/sodium bicarbonate) packet for oral suspension <i>QTY LIMIT:</i> 1 packet/day	<p>Konvomep, Omeprazole/sodium bicarb packet, Zegerid packet: The patient has a documented side effect, allergy, or treatment failure to omeprazole/sodium bicarb capsules OR patient has a medical necessity for a non-solid oral dosage form and the patient has a documented side effect, allergy, or treatment failure with lansoprazole ODT or Nexium powder for suspension.</p> <p>LIMITATIONS: First-Lansoprazole® and First-Omeprazole Suspension Kits are not covered as Federal Rebate is no longer offered.</p>

GAUCHER'S DISEASE MEDICATIONS

All products require PA	Cerezyme® (imiglucerase for injection) Cerdelga® (eliglustat) <i>QTY LIMIT:</i> 2 caps/day Elelyso® (taliglucerase alfa for injection) Vpriv® (velaglucerase alfa for injection) Miglustat (compare to Zavesca®) <i>QTY LIMIT:</i> 3 caps/day Zavesca® (miglustat) <i>QTY LIMIT:</i> 3 caps/day **Maximum days supply per fill for all drugs is 14 days**	<p>CRITERIA FOR APPROVAL: The diagnosis or indication is Gaucher disease (GD) type I. AND The diagnosis has been confirmed by molecular or enzymatic testing.</p> <p>Age Limits</p> <p>Elelyso, Vpriv: for patients ≥ 4 years old Cerezyme: for patients ≥ 2 years old Cerdelga, Miglustat, Zavesca: for patients ≥ 18 years old</p> <p>Cerezyme/Vpriv additional criteria: Failure, intolerance or other contraindication to enzyme replacement therapy with Elelyso</p> <p>Cerdelga additional criteria:</p> <ul style="list-style-type: none"> Testing to verify if CYP2D6 extensive metabolizer (EM), intermediate
-------------------------	---	--

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>metabolizer (IM), poor metabolizer (PM), or if CYP2D6 genotype cannot be determined</p> <ul style="list-style-type: none"> ○ Dose max: 84mg twice/day if EM or IM ○ Dose max: 84mg/day if PM ○ Case by case determination if CYP2D6 cannot be determined <p>Miglustat, Zavesca additional criteria:</p> <ul style="list-style-type: none"> • For whom enzyme replacement therapy is not a therapeutic option (e.g. due to allergy, hypersensitivity, or poor venous access) AND for approval of miglustat, the patient must have a documented intolerance to brand Zavesca.

GOUT AGENTS

<p>ALLOPURINOL (compare to Zyloprim®) COLCHICINE tablets COLCHICINE/PROBENECID FEBUXOSTAT (compare to Uloric®) <i>QTY LIMIT:</i> 40 mg tablets = 1 tablet/day PROBENECID</p>	<p>Colchicine capsules Mitigare® (colchicine) capsule <i>QTY LIMIT:</i> 2 capsules/day Uloric® (febuxostat) <i>QTY LIMIT:</i> 40 mg tablets = 1 tablet/day Zyloprim® (allopurinol)</p>	<p>Colchicine capsules, Mitigare: the patient has a documented intolerance to generic colchicine tablets.</p> <p>Uloric: The patient has had a documented intolerance to generic febuxostat.</p> <p>Zyloprim: The patient has had a documented intolerance to generic allopurinol</p>
--	--	--

GROWTH STIMULATING AGENTS

ACHONDROPLASIA TREATMENTS		
<p>All products require PA</p>	<p>Voxzogo™ (vosoritide)</p>	<p>Voxzogo: The patient must have a diagnosis of achondroplasia confirmed with genetic testing AND the medication must be prescribed by a pediatric endocrinologist AND Confirmation of non-closure of epiphyseal plates (x-ray determining bone age) must be provided for females > age 12 and males > age 14 AND Voxzogo will not be used in combination with growth hormone (e.g. somatropin), growth hormone analogs (e.g. somapacitan), or insulin-like growth factor (IGF-1) (e.g. mecasermin) AND patient's standing height, weight, BMI, and upper to lower body ratio will be measured at baseline and monitored throughout therapy. For re-approval, the patient must have an improvement in growth velocity compared to pre-treatment baseline.</p>
GROWTH HORMONE		
<p><u><i>Preferred After Clinical Criteria Are Met</i></u></p> <p>GENOTROPIN® NORDITROPIN®</p>	<p>Humatrope® Ngenla™ (somatogon-ghla) Nutropin® AQ Omnitrope® Saizen®</p>	<p>Criteria for Approval Pediatric: 1) The patient must have one of the following indications for growth hormone: Turner syndrome confirmed by genetic testing. Prader-Willi Syndrome confirmed by genetic testing. □ Growth deficiency due to chronic renal failure. Patient who is Small for</p>

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

Skytrofa® (lonapegsomatropin-tcgd)
Sogroya® (somapacitan-beco)
Zomacton®

Specialized Indications – See Specific Criteria

Increlex® (mecasermin)
Serostim®
Zorbtive®

Gestational Age (SGA) due to Intrauterine Growth Retardation (IUGR) and catch-up growth not achieved by age 2 (Birth weight less than 2500g at gestational age of <37 weeks or a birth weight or length below the 3rd percentile for gestational age). OR Pediatric Growth Hormone Deficiency confirmed by results of two provocative growth hormone stimulation tests (insulin, arginine, levodopa, propranolol, clonidine, or glucagon) showing results (peak level) <10ng/ml. 2) The requested medication must be prescribed by a pediatric endocrinologist (or pediatric nephrologist if prescribed for growth deficiency due to chronic renal failure). 3) Confirmation of non-closure of epiphyseal plates (x-ray determining bone age) must be provided for females > age 12 and males > age 14. 4) Initial requests can be approved for 6 months. Subsequent requests can be approved for up to 1 year with documentation of positive response to treatment with growth hormone.

Criteria for Approval Adult: The patient must have one of the following indications for growth hormone: Panhypopituitarism due to surgical or radiological eradication of the pituitary. OR Adult Growth Hormone Deficiency confirmed by one growth hormone stimulation test (insulin, arginine, levodopa, propranolol, clonidine, or glucagon) showing results (peak level) <5ng/ml. Growth hormone deficient children must be retested after completion of growth.

LIMITATIONS: Coverage of Growth Hormone products will not be approved for patients who have Idiopathic Short Stature.

Humatrope, Nutropin AQ, Omnitrope, Saizen, Skytrofa, Zomacton: The patient has a documented side effect, allergy, or treatment failure to both preferred agents.

Ngenla, Sogroya: The patient has a documented side effect, allergy, or treatment failure to both preferred agents AND the patient has a documented side effect, allergy, or treatment failure to Skytrofa.

Increlex: Member has growth hormone gene deletion AND neutralizing antibodies to growth hormone, OR primary insulin-like growth factor (IGF-1) deficiency (IGFD), defined by the following: o Height standard deviation score < -3 AND Basal IGF-1 standard deviation score < -3 AND Normal or elevated growth hormone level AND Member is ≥ 2 years old (safety and efficacy has not been established in patients younger than 2), AND Member has open epiphysis, AND Member is under the care of an endocrinologist or other specialist trained to diagnose and treat growth disorders.

Serostim: A diagnosis of AIDS associated wasting/anorexia

Zorbtive: A diagnosis of short bowel syndrome. Concomitant use of specialized nutritional support (specialty TPN) Prescription must be issued by gastroenterologist (specialist)

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

hATTR TREATMENTS

<p>All products require PA</p>	<p>Amvuttra™ (vutrisiran) 25mg/0.5ml injection for subcutaneous use <i>QTY LIMIT</i>: 1 syringe (0.5ml) every 3 months</p> <p>Onpattro® (patisiran) 10 mg/5ml intravenous injection Weight < 100kg (0.3 mg/kg every 3 weeks) Weight ≥ 100kg (30 mg every 3 weeks)</p> <p>Vyndamax® (tafamidis) <i>QTY LIMIT</i>: 1 capsule/day</p> <p>Vyndaqel® (tafamidis meglumine) <i>QTY LIMIT</i>: 4 capsules/day</p> <p>Wainua™ (eplontersen)</p>	<p>Amvuttra, Onpattro, Wainua:</p> <ul style="list-style-type: none"> The patient is ≥ 18 years of age with a diagnosis of polyneuropathy of hereditary transthyretin mediated (hATTR) amyloidosis (Documentation of TTR mutation by genetic testing or the presence of amyloid deposits via tissue biopsy has been submitted) AND The medication is being prescribed by or in consultation with a neurologist AND Clinical signs and symptoms of the disease (e.g., peripheral/autonomic neuropathy, motor disability, cardiovascular dysfunction, renal dysfunction) are present and other causes of neuropathy have been excluded AND Patient is receiving vitamin A supplementation AND Initial approval will be granted for 3 months. For re-approval, the patient must have documentation of clinical improvement or slower progression of the disease than would otherwise be expected. <p>Vyndamax, Vyndaqel:</p> <ul style="list-style-type: none"> The patient is ≥ 18 years of age with a diagnosis of cardiomyopathy of wild type transthyretin-mediated amyloidosis or hereditary transthyretin mediated (hATTR) amyloidosis AND The presence of amyloid deposits showing cardiac involvement via tissue biopsy or imaging has been submitted AND The medication is being prescribed by or in consultation with a cardiologist AND Initial approval will be granted for 6 months. For re-approval, the patient must have a decrease in the frequency of cardiovascular-related hospitalizations or slower progression of the disease than would otherwise be expected.
--------------------------------	---	--

HEART FAILURE

ANGIOTENSIN RECEPTOR – NEPRILYSIN INHIBITOR (ARNI)		
ENTRESTO® (valsartan/sacubitril) <i>QTY LIMIT</i> : 2 tablets/day		
CARDIAC MYOSIN INHIBITORS		

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
All procutts require PA	Camzyos® (mavacamten) <i>QTY LIMIT</i> : 1 capsule/day	<p>Camzyos:</p> <ul style="list-style-type: none"> The diagnosis or indication is symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) AND <ul style="list-style-type: none"> LVEF \geq 55% AND Valsalva LVOT peak gradient \geq50mmHg at rest or with provocation AND The patient has a documented side effect, allergy, or treatment failure at a maximally tolerated dose to at least two of the following: Non-vasodilating beta blocker (e.g., atenolol, bisoprolol, metoprolol, nadolol, propranolol), Non-dihydropyridine calcium channel blocker (i.e., diltiazem, verapamil), and Disopyramide AND The medication will not be used concurrently with disopyramide, ranolazine, verapamil with a beta blocker, or diltiazem with a beta blocker. Approval will be granted for 12 months. For reapproval, there must be a documented positive clinical response as supported by one of the following: Stable or reduction in New York Heart Association (NYHA) class AND Patient has a left ventricular ejection fraction of greater than or equal to 50%
LOOP DIURETICS		
BUMETANIDE (compare to Bumex®) tablet FUROSEMIDE (compare to Lasix®) tablet, oral solution TORSEMIDE (compare to Demadex®) tablet	Edecrin® (ethacrynic acid) tablet Ethacrynic Acid (compare to Edecrin®) Furoscix® (furosemide) injection <i>QTY LIMIT</i> : 4 injections/30 days (maximum 30-day supply) Lasix® (furosemide) tablet	<p>Ethacrynic Acid, Edecrin: The patient has had a documented side effect, allergy or treatment failure to at least two preferred agents and for Ethacrynic acid the patient has a documented intolerance to brand Edecrin.</p> <p>Furoscix: The indication for use is the treatment of congestion due to fluid overload in adults with NYHA Class II or Class III chronic heart failure AND the medication is being prescribed by or in consultation with a cardiologist AND the patient is experiencing symptoms despite compliance with oral loop diuretic therapy AND oral loop diuretic therapy will be resumed as soon as practical AND medical reasoning beyond convenience is provided for not pursuing therapy in an outpatient infusion setting. PA approval will be authorized for 1 month. Subsequent approvals will be for 1 year.</p> <p>Lasix: The patient has a documented intolerance to generic furosemide.</p>
SODIUM-GLUCOSE CO-TRANSPORTER (SGLT) INHIBITORS		
FARXIGA® (dapagliflozin) JARDIANCE® (empagliflozin)	Inpefa® (sotagliflozin) <i>QTY LIMIT</i> : 1 tab/day	Inpefa: The patient has a documented side effect, allergy, or contraindication to Farxiga and Jardiance.
SOLUBLE GUANYLATE CYCLASE (sGC) STIMULATORS		

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

All products require PA	Verquvo® (vericiguat) tablet <i>QTY LIMIT:</i> 1 tablet/day	<p>Verquvo: The diagnosis or indication is symptomatic heart failure (HF) with ejection fraction < 45% AND the patient has been hospitalized for HF within the previous 6 months or required the use of IV diuretics within the past 3 months AND the patient is not pregnant AND the patient is concurrently receiving the maximum tolerated dose of one agent from each of the following classes, unless contraindicated:</p> <ul style="list-style-type: none"> • ARNI, ACE-I, or ARB • Beta Blocker (metoprolol, carvedilol, or bisoprolol) • Aldosterone antagonist if LVEF ≤ 35% or LVEF ≤ 40% with diabetes mellitus or post myocardial infarction (MI) with HF symptoms
-------------------------	--	--

HEMATOPOIETICS

Colony Stimulating Factors

Eflapegrastim Products

All products require PA

Filgrastim Products

NEUPOGEN® (filgrastim) Vial, Syringe
NIVESTYM™ (figrastim-aafi) Vial, Syringe

Pegfilgrastim Products

FULPHILA™ (pegfilgrastim-jmdb) Syringe
NEULASTA® (pegfilgrastim) Syringe
NEULASTA® Onpro® (pegfilgrastim) kit
ZIEXTENZO® (pegfilgrastim-bmez)

Rolvedon™ (eflapegrastim-xnst) Syringe

Granix® (tbo-filgrastim) Vial, Syringe
Leukine® (sargramostim)
Releuko™ (filgrastim-ayow)
Zarxio® (filgrastim-sndz) Syringe

Fylnetra® (pegfilgrastim-pbbk)
Nyvepria (pegfilgrastim-apgf)
Stimufend® (pegfilgrastim-fpgk)
Udenyca™ (pegfilgrastim-cbqv)

Granix, Leukine, Releuko, Zarxio: The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons Neupogen would not be a suitable alternative.

Fylnetra, Nyvepria, Rolvedon, Stimufend, Udenyca: The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred pegfilgrastim products would not be suitable alternatives.

Erythropoietic Stimulating Agents

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>Preferred After Clinical Criteria Are Met</u> EPOGEN® (epoetin alpha) MIRCERA® (methoxypolyethylene glycolepoetin beta)</p>	<p>Aranesp® (darbepoetin alfa) Procrit® (epoetin alpha) Retacrit® (epoetin alpha-epbx)</p>	<p>Aranesp, Procrit, Epogen, Retacrit: diagnosis or indication for the requested medication is anemia due to one of the following: Chronic kidney disease/renal failure, Post-renal transplant, use of zidovudine for the treatment of human immunodeficiency virus (HIV) (other causes of anemia, such as iron/folate/vitamin B12 deficiency have been eliminated), Surgery patients at high risk for perioperative blood loss, Cancer chemotherapy, Use of ribavirin or interferon therapy for Hepatitis C, Myelodysplastic syndrome. Hemoglobin level at initiation of therapy is <10 g/dL OR for patients currently maintained on therapy, hemoglobin level is < 11 g/dL in dialysis patients with chronic kidney disease, < 10 g/dL in non-dialysis patients with chronic kidney disease, or < 12 g/dL in patients treated for other indications AND for approval of Aranesp or Procrit, or Retacrit the patient has had a documented side effect, allergy, or treatment failure to Epogen.</p> <p>Mircera: The diagnosis or indication for the requested medication is anemia due to chronic kidney disease/renal failure AND Hemoglobin level at initiation of therapy is <10g/dl OR For patients currently maintained on therapy, hemoglobin level is ≤11 g/dL in dialysis patients with chronic kidney disease, ≤10 g/dL in non-dialysis patients with chronic kidney disease.</p>

HEMOPHILIA TREATMENTS

(Factor VII Deficiency)

All products require PA

Novoseven® RT
Sevenfact®

Novoseven RT: Medication is being used for the treatment of acute bleeding episodes in a patient with Hemophilia A or B with inhibitors OR Patient has congenital Factor VII deficiency.

Sevenfact: Medication is being used for the treatment of acute bleeding episodes in a patient with Hemophilia A or B with inhibitors AND there is a clinically compelling reason why Novoseven RT cannot be used.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
Hemophilia A (Factor VIII Deficiency)		
<p>ALTUVIIIIO™ (antihemophilic factor (recombinant), Fc- VWF-XTEN fusion protein-ehtl) HEMLIBRA® (emicizumab-kxwh) HEMOFIL® M JIVI® KOATE®-DVI KOVALTRY® NOVOEIGHT® NUWIQ® OBIZUR® XYNTHA®</p>	<p>Advate® Adynovate® Afstyla® Eloctate® Esperoct® Kogenate® Recombinate</p>	<p>Adynovate, Eloctate, Esperoct: Documentation must include why the member is unable to use the preferred extended half-life concentrate Jivi or Altuviiiio. Advate, Afstyla, Kogenate, Recombinate: Documentation must be provided why member is unable to use each of the preferred non-extended concentrates.</p>
Hemophilia B (Factor IX Deficiency)		
<p>ALPHANINE® SD BENEFIX® IXINITY® PROFILNINE® REBINYN® RIXUBIS®</p>	<p>Alprolix® Idelvion® Kcentra®</p>	<p>All Non-Preferred Products: The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred products would not be suitable alternatives. For approval of Alprolix or Idelvion, documentation must include why the member is unable to use the preferred extended half-life concentrate Rebinyn.</p>
Von Willebrand Factor		
<p>ALPHANATE® HUMATE-P® WILATE®</p>	<p>Vonvendi®</p>	<p>All Non-Preferred Products: The prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why any of the preferred products would not be suitable alternatives.</p>
AHF-Anti-Inhibitor Coagulation Complex		
<p>All products require PA</p>	<p>Feiba®</p>	<p>Feiba: medication is being used for the treatment of acute bleeding episodes or routine prophylaxis in a patient with Hemophilia A or B with inhibitors.</p>
Gene Therapy		

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
All products require PA	<p>Beqvez™ (fidanacogene elaparvovec-dzkt) Hemgenix® (etranacogene dezaparvovec-drlb) Roctavian™ (valoctocogene roxaparvovec-rvox)</p>	<p>Criteria for all gene therapy products: The provider, healthcare facility, and patient will attest to continued clinical information exchange with the Department of Vermont Health Access, as is necessary to meet the terms of any value-based rebate agreements that have been entered into by the Department of Vermont Health Access. (e.g. patient response to therapy, clinical lab values to support treatment success/failure, annual follow up documentation).</p> <p>Beqvez:</p> <ul style="list-style-type: none"> • Patient is ≥ 18 years of age AND • Patient has a diagnosis of severe congenital Factor IX deficiency, as evidenced by < 1% of normal circulating factor IX AND • Patient has the following: <ul style="list-style-type: none"> ○ Current and continuous use of Factor IX prophylaxis therapy for the previous 6 months as evidenced by claims history or clinical documentation, without breaks in adherence. (Continuous use is defined as routine prophylaxis with defined frequency, e.g. twice weekly, once every two weeks) AND ○ Current or historical life-threatening hemorrhage despite use of preferred prophylaxis therapy OR ○ Repeated, serious spontaneous bleeding episodes requiring hospitalization AND • Patient has been tested and found negative for Factor IX inhibitor titers and had no prior history for Factor IX inhibition AND • Patient must have a negative baseline anti-AAVRh74var antibody titer AND • The patient meets one of the following: <ul style="list-style-type: none"> ○ Patient is not HIV positive; or ○ Patient is HIV positive and is virally suppressed with anti-viral therapy (i.e., < 20 copies of HIV per mL or DC4+ cell count > 200 mm³) AND • The patient is not currently using antiviral therapy for hepatitis B or C AND does not have significant liver dysfunction/significant fibrosis. • Baseline liver function tests will be completed prior to start of therapy and continued per package insert following Beqvez administration AND • Factor IX activity will be monitored weekly for 3 months AND • Approval will be granted for a max one-time dose per lifetime and may not be renewed <p>Hemgenix:</p> <ul style="list-style-type: none"> • Patient is ≥ 18 years of age AND • Patient has a diagnosis of severe congenital Factor IX deficiency, as evidenced by < 1% of normal circulating factor IX AND • Patient has the following: <ul style="list-style-type: none"> ○ Current and continuous use of Factor IX prophylaxis therapy for the previous 6 months as evidence by claims history or clinical documentation, without breaks in adherence. (Continuous use is defined as routine prophylaxis with defined frequency, e.g. twice weekly, once every two weeks) AND ○ Current or historical life-threatening hemorrhage despite use of preferred prophylaxis therapy OR Repeated, serious spontaneous bleeding episodes requiring hospitalization AND

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

		<ul style="list-style-type: none"> • Patient has been tested and found negative for Factor IX inhibitor titers (if test result is positive, re-test within approximately 2 weeks. If re-test is also positive, Hemgenix should not be administered) AND • Patient must have a baseline anti-AAV5 antibody titer of less than or equal to 1:678 measured by ELISA AND • Baseline liver function tests will be completed prior to start of therapy and continued weekly for 3 months following Hemgenix administration AND • Factor IX activity will be monitored weekly for 3 months AND Factor IX prophylaxis therapy will be discontinued when circulating factor IX levels reach 5% • Approval will be granted for a max one-time dose per lifetime and may not be renewed <p>Roctavian:</p> <ul style="list-style-type: none"> • Patient is ≥ 18 years of age AND • Patient has a diagnosis of severe congenital Factor VIII deficiency, as evidenced by $< 1\%$ of normal circulating factor VIII AND • Patient has the following: <ul style="list-style-type: none"> ○ Current and continuous use of Factor VIII prophylaxis therapy for the previous 12 months as evidenced by claims history or clinical documentation, without breaks in adherence. (Continuous use is defined as routine prophylaxis with defined frequency, e.g. twice weekly, once every two weeks) AND ○ Current or historical life-threatening hemorrhage despite use of preferred prophylaxis therapy OR repeated, serious spontaneous bleeding episodes requiring hospitalization AND • Patient must be anti-AAV5 antibody negative, • Patient has been tested and found negative for Factor VIII inhibitor titers AND • The patient meets one of the following: <ul style="list-style-type: none"> ○ Patient is not HIV positive; or ○ Patient is HIV positive and is virally suppressed with anti-viral therapy (i.e., < 200 copies of HIV per mL) AND • The patient's hepatitis B surface antigen is negative AND • The patient meets one of the following: <ul style="list-style-type: none"> ○ Patient's hepatitis C virus (HCV) antibody is negative; or ○ Patient's HCV antibody is positive, and the patient's HCV RNA is negative AND • The patient is not currently using antiviral therapy for hepatitis B or C AND does not have significant liver dysfunction/significant fibrosis. • The patient does not have significant renal impairment (Creatinine ≥ 1.5mg/dL)
--	--	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>cirrhotic, HBV negative, no prior liver transplantation, and not pregnant.</p> <ul style="list-style-type: none"> See prior authorization form for detailed requirements and for documentation required. <p>For approval of a non-preferred agent, the provider must submit clinical documentation detailing why the patient is not a candidate for a preferred direct acting agent regimen.</p>

HEREDITARY ANGIOEDEMA MEDICATIONS

TREATMENT

Preferred After Clinical Criteria are Met

BERINERT® (human C1 inhibitor)
 ICATIBANT (compare to Firazyr®)
QTY LIMIT: 3 syringes (9 ml)/fill

Firazyr® (icatibant)
QTY LIMIT: 3 syringes (9 ml)/fill
 Kalbitor® (escallantide)
QTY LIMIT: 6 vials (2 packs) per fill
 Ruconest® (recombinant C1 esterase inhibitor)
QTY LIMIT: 4 vials/fill

Berinert, Firazyr, Icatibant: The diagnosis or indication is treatment of an acute Hereditary Angioedema (HAE) attack AND for approval of Firazyr, the patient must have a documented intolerance to generic Icatibant. (Approval may be granted so that 2 doses may be kept on hand for Berinert and 3 doses for Icatibant/Firazyr).

Kalbitor, Ruconest: The diagnosis or indication is treatment of an acute Hereditary Angioedema (HAE) attack AND the patient has a documented side effect, allergy, treatment failure or contraindication to a preferred agent (Approval may be granted so that 2 doses may be kept on hand.)

PROPHYLACTIC

Preferred After Clinical Criteria are Met

CINRYZE® (human C1 inhibitor)
QTY LIMIT: 20 vials/30days
 HAEGARDA® (human C1 inhibitor)
 ORLADEYO™ (berotralstat)
QTY LIMIT: 1 capsule/day
 TAKHZYRO™ (lanadelumab-flyo)
QTY LIMIT: 2 vials/28 days

Cinryze, Haegarda, Orladeyo, Takhzyro: The diagnosis or indication is prophylaxis of Hereditary Angioedema (HAE) attacks.

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

HIDRADENITIS SUPPURATIVA

BIOLOGICS: Initial approval is 3 months; renewals are 1 year

Preferred After Clinical Criteria Are Met

INJECTABLE

ADALIMUMAB-ADBM (compare to Cyltezo®) biosimilar to Humira®

HUMIRA® (adalimumab)

QTY LIMIT: 6 syringes/28 days for the first month (HS starter kit); 4 syringes/28 days subsequently

Abrilada™ (adalimumab-afzb) biosimilar to Humira®
Adalimumab-adaz (compare to Hyrimoz®) biosimilar to Humira®
Adalimumab-adbm (compare to Cyltezo®) biosimilar to Humira®
Adalimumab-fkjp (compare to Hulio®) biosimilar to Humira®
Amjevita™ (adalimumab-atto) biosimilar to Humira®
Cosentyx® (secukinumab)
Cyltezo® (adalimumab-adbm) biosimilar to Humira®
Hadlima™ (adalimumab-bwwd) biosimilar to Humira®
Hulio® (adalimumab-fkjp) biosimilar to Humira®
Hyrimoz® (adalimumab-adaz) biosimilar to Humira®
Idacio® (adalimumab-aacf) biosimilar to Humira®
Simlandi® (adalimumab-ryvk) biosimilar to Humira®
Yuflyma® (adalimumab-aaty) biosimilar to Humira®
Yusimry™ (adalimumab-aqvh) biosimilar to Humira®

Humira:

- The patient has a diagnosis of moderate-severe hidradenitis suppurativa (Hurley Stage II-III) AND
- The medication is being prescribed by, or in consultation with, a dermatologist AND
- The patient has not responded to a 12-week course of standard antibiotic therapy with an oral tetracycline (e.g. Doxycycline) or clindamycin plus rifampin, unless contraindicated.

Humira Biosimilars: the patient must be unable to use Humira.

Cosentyx additional criteria: the prescriber must provide evidence of a trial and failure or contraindication to Humira. **Note:** Cosentyx approvals for 300mg dose(s) must use “300DOSE” package (containing 2x150mg pens or syringes) Approval will not be granted for 2 separate 150mg packages.

HYPERKALEMIA AGENTS

Lokelma™ (sodium zirconium cyclosilicate)

SPS® (sodium polystyrene sulfonate) suspension
Veltassa® (patiromer sorbitex calcium) powder packets
QTY LIMIT: 1 packet/day

SPS: The patient has potentially life-threatening hyperkalemia AND where clinically appropriate, a loop or thiazide diuretic has failed for potassium removal AND newer cation exchangers (i.e. SZC or patiromer) are not available AND the patient does not have any high risk factors for intestinal necrosis defined as:

- Postoperative patients
- Patients with an ileus
- Patients with a large or small bowel obstruction
- Patients with constipation or at risk of becoming constipated (eg, due to opioid use)
- Patients with underlying bowel disease, eg, ulcerative colitis or Clostridioides difficile colitis

Veltassa: The patient requires therapy for the treatment of non-emergent hyperkalemia AND where clinically appropriate, medications known to cause

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		hyperkalemia (e.g. ACE inhibitors, ARBs, aldosterone antagonists, NSAIDs) have been discontinued or reduced to the lowest effective dose AND where clinically appropriate, a loop or thiazide diuretic has failed for potassium removal, AND the patient has been counseled to follow a low potassium diet (\leq 3 grams/day).
HYPOTHYROID AGENTS		
ARMOUR THYROID tablet EUTHYROX® (levothyroxine) tablet LEVOTHYROXINE tablet LEVOXYL® (levothyroxine) tablet LIOTHYRONINE (compare to Cytomel®) tablet NP THYROID® (thyroid) tablet UNITHROID® (levothyroxine) tablet <u>Preferred after clinical criteria met</u> ERMEZA™ (levothyroxine) oral solution	Cytomel® (liothyronine) tablet Levothyroxine capsule (compare to Tirosint®) Synthroid® (levothyroxine) tablet Thyquidity™ (levothyroxine) oral solution Tirosint®-Sol (levothyroxine) oral solution	Ermeza: The patient has a medical necessity for a non-solid oral dosage form (dysphagia) Thyquidity, Tirosint-Sol: The patient has a medical necessity for a non- solid oral dosage form and the medication cannot be administered by crushing oral tablets AND the patient must have documented intolerance to Ermeza. Levothyroxine capsule: patient has had a documented side effect, allergy, or treatment failure to 2 preferred hypothyroid agents. Cytomel, Synthroid: The patient has a documented intolerance to the generic equivalent.
IDIOPATHIC PULMONARY FIBROSIS (IPF)		
PIRFENIDONE (compare to Esbriet) <i>QTY LIMIT:</i> 267 mg tablets/capsules = 270 tabs/caps per month, 801mg tablets = 90 tabs/caps per month	Esbriet® (pirfenidone) <i>QTY LIMIT:</i> 267 mg tablets = 270 tabs/month, 801 mg tablets = 90 tabs/month Ofev® (nintedanib) <i>QTY LIMIT:</i> 60 tabs/month	Clinical Criteria: Esbriet, Ofev <ul style="list-style-type: none"> ○ Age \geq 18 ○ Diagnosis of idiopathic pulmonary fibrosis (pirfenidone and Ofev) OR chronic fibrosing interstitial lung disease or systemic sclerosis associated interstitial lung disease (Ofev Only) ○ May not be used in combination. ○ The prescriber is a pulmonologist. ○ Clinical documentation that the member is a non-smoker or has not smoked in 6 weeks. ○ FVC \geq 50% of predicted ○ For approval Esbriet or Ofev (idiopathic pulmonary fibrosis diagnosis only), the patient must have a documented intolerance to generic pirfenidone. Reauthorization Criteria: <ul style="list-style-type: none"> ○ Documentation the patient is receiving clinical benefit from Esbriet or Ofev therapy as evidenced by < 10% decline in percent predicted FVC or < 200mL decrease in FVC AND ○ There is clinical documentation that the member has remained tobacco-free.

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

IMMUNOLOGIC THERAPIES FOR ASTHMA

Initial 6 months, Renewal 1 year

Preferred After Clinical Criteria are Met

DUPIXENT® (dupilumab) subcutaneous injection,
pre-filled syringe, and auto-injector pen

QTY LIMIT: 4 syringes/pens the first
28 days then 2 syringes/pens every
28 days thereafter

NUCALA® (mepolizumab) subcutaneous injection,
auto-injector pen

QTY LIMIT: 1mL every 28 days

XOLAIR® (omalizumab) subcutaneous injection
vial, prefilled syringe

QTY LIMIT: 900 mg every 28 days

Cinqair® (reslizumab) Intravenous injection

Fasenra® (benralizumab) subcutaneous injection, pre-
filled syringe and auto-injector pen

QTY LIMIT: 1 mL every 28 days for 3 doses then 1
mL every 56 days

Nucala® (mepolizumab) subcutaneous
injection, vial, pre-filled syringe

QTY LIMIT: 1mL every 28 days

Tezspire™ (tezepelumab-ekko) subcutaneous injection,
pre-filled syringe and auto-injector pen

QTY LIMIT: 1.91 mL every 28 days

Xolair:

Diagnosis of moderate to severe persistent asthma:

- The patient must be 6 years of age or older AND
- The patient has a history of uncontrolled asthma symptoms (symptoms occurring almost daily or waking at night with asthma at least once a week) or 2 or more exacerbations in the previous year despite regular use of medium-high dose ICS/LABA for a minimum of 3 consecutive months, with or without oral corticosteroids. Pharmacy claims will be evaluated to assess compliance with therapy. AND
- The prescriber is a pulmonologist, allergist, or immunologist AND
- Patient has tested positive to at least one perennial aeroallergen by skin or blood test (i.e.: RAST, CAP, intracutaneous test) AND
- Patient has an IgE level ≥ 30 and ≤ 700 IU/ml (ages 12 and older) OR IgE level ≥ 30 and ≤ 1300 IU/ml (ages 6-11) prior to beginning therapy with Xolair.
- For continuation of therapy after the initial 6-month authorization, the patient must continue to receive therapy with an ICS/LABA AND have either a decreased frequency of exacerbations, decreased use of maintenance oral corticosteroids, reduction in the signs and symptoms of asthma, or an increase in predicted FEV1 from baseline.

Diagnosis of chronic idiopathic urticaria:

- The patient must be 12 years of age or older AND
- The patient has a therapeutic failure or contraindication to an H1 antihistamine (e.g. cetirizine, fexofenadine) at double the daily dose
- For continuation of therapy after the initial 6-month authorization, the patient must have documented clinical improvement in symptoms.

Diagnosis of Chronic Rhinosinusitis with Nasal Polyps:

- Patient is 18 years of age or older AND
- Prescriber is an allergist or ENT specialist AND
- Patient has had an inadequate response to at least a 3-month trial of 2 different nasal corticosteroids AND
- Patient has had an inadequate response to at least a 10-14 day course of oral corticosteroids AND
- Patient will use Xolair concurrently with an Intranasal corticosteroid.
- For continuation of therapy after the initial 6-month authorization, the patient must continue to receive therapy with an intranasal corticosteroid AND there must be documented improvement in nasal symptoms.

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

Diagnosis of IgE mediated food allergy:

- The patient must be 1 year of age or older AND
- The patient has a diagnosis of IgE-mediated food allergy to at least one of the following: cashew, egg, hazelnut, milk, peanut, walnut wheat AND
- Patient has an IgE level ≥ 30 and ≤ 1850 IU/ml AND
- Prescriber is an allergist or immunologist AND
- The patient has history of significant symptomatic allergic reaction that was demonstrated through signs and symptoms (hives, swelling, wheezing, hypotension, or gastrointestinal symptoms)
- For continuation of therapy after the initial 6 month authorization, the patient must have clinical documentation of food avoidance, a positive clinical response to therapy, and confirmed adherence to treatment.

Limitations: Xolair use will not be approved if requested for patients with a diagnosis of moderate to severe persistent asthma who are currently smoking.

Fasenra, Nucala, Cinqair:

Diagnosis of moderate to severe persistent asthma:

- The patient must be 6 years of age or older for Nucala, 12 years of age or older for Fasenra, or 18 years of age or older for Cinqair AND
- The patient must have a diagnosis of severe persistent asthma with an eosinophilic phenotype as defined by pre-treatment blood eosinophil count of ≥ 150 cells per mcL within the previous 6 weeks or ≥ 300 cells per mcL within 12 months prior to initiation of therapy AND
- The patient has a history of uncontrolled asthma symptoms (symptoms occurring almost daily or waking at night with asthma at least once a week) or 2 or more exacerbations in the previous year despite regular use of medium-high dose ICS/LABA for a minimum of 3 consecutive months, with or without oral corticosteroids. Pharmacy claims will be evaluated to assess compliance with therapy. AND
- The prescriber is an allergist, immunologist, or pulmonologist. AND
- For approval of Cinqair or Fasenra, the patient must have a documented side effect, allergy, or treatment failure with Dupixent or Nucala. For approval of Nucala vial or prefilled syringe, the patient must be unable to use the auto-injector.
- For continuation of therapy after the initial 6-month authorization, the patient must continue to receive therapy with an ICS/LABA AND have either a decreased frequency of exacerbations, decreased use of maintenance oral corticosteroids, reduction in the signs and symptoms of asthma, or an increase in predicted FEV₁ from baseline.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p><i>Diagnosis of hypereosinophilic syndrome (Nucala only):</i></p> <ul style="list-style-type: none"> • Patient must be 12 years of age or older AND • The patient must have a blood eosinophil count of $\geq 1,000$ cells per mcl AND • The patient has had at least 2 HES flares within the past 12 months AND • The patient is on a stable dose of background HES therapy (chronic or episodic corticosteroids, immunosuppressive, or cytotoxic therapy) for at least 4 weeks prior to treatment initiation AND • The prescriber is an allergist, hematologist, immunologist, or pulmonologist • For continuation of therapy after the initial 6-month authorization, the patient must continue to receive background HED therapy AND there must be documented improvement in the number or frequency of HES flares. <p><i>Diagnosis of Chronic Rhinosinusitis with Nasal Polyps (Nucala Only):</i></p> <ul style="list-style-type: none"> • Patient is 18 years of age or older AND • Prescriber is an allergist or ENT specialist AND • Patient has had an inadequate response to at least a 3-month trial of 2 different nasal corticosteroids AND • Patient has had an inadequate response to at least a 10–14-day course of oral corticosteroids AND • Patient will use Nucala concurrently with an intranasal corticosteroid • For continuation of therapy after the initial 6-month authorization, the patient must continue to receive therapy with an intranasal corticosteroid AND there must be documented improvement in nasal symptoms. <p>Dupixent:</p> <p><i>Diagnosis of moderate to severe persistent asthma:</i></p> <ul style="list-style-type: none"> • The patient must be 6 years of age or older AND • The patient must have an eosinophilic phenotype as defined by pre-treatment blood eosinophil count of ≥ 150 cells per mcL within the previous 6 weeks or ≥ 300 cells per mcL within 12 months prior to initiation of therapy OR the patient is dependent on oral corticosteroids. • The patient has a history of uncontrolled asthma symptoms (symptoms occurring almost daily or waking at night with asthma at least once a week) or 2 or more exacerbations in the previous year despite regular use of medium-high dose ICS/LABA for a minimum of 3 consecutive months, with or without oral corticosteroids. Pharmacy claims will be evaluated to assess compliance with therapy. AND • The prescriber is an allergist, immunologist, or pulmonologist AND

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<ul style="list-style-type: none"> • For continuation of therapy after the initial 6-month authorization, the patient must continue to receive therapy with an ICS/LABA AND have either a decreased frequency of exacerbations OR decreased use of maintenance oral corticosteroids OR reduction in the signs and symptoms of asthma OR an increase in predicted FEV1 from baseline. <p><i>Diagnosis of Chronic Rhinosinusitis with Nasal Polyps:</i></p> <ul style="list-style-type: none"> • Patient is 12 years of age or older AND • Prescriber is an allergist or ENT specialist AND • Patient has had an inadequate response to at least a 3-month trial of 2 different nasal corticosteroids AND • Patient has had an inadequate response to at least a 10–14-day course of oral corticosteroids AND • Patient will use Dupixent concurrently with an intranasal corticosteroid • For continuation of therapy after the initial 6-month authorization, the patient must continue to receive therapy with an intranasal corticosteroid AND there must be documented improvement in nasal symptoms. <p><i>Diagnosis of Eosinophilic Esophagitis:</i></p> <ul style="list-style-type: none"> • Patient is 1 years of age or older, weighing at least 15kg AND • Prescriber is an allergist or gastroenterologist AND • Diagnosis is confirmed by endoscopic esophageal biopsy showing ≥ 15 intraepithelial eosinophils per high-power field AND • Symptoms of esophageal dysfunction are present (e.g. pain while swallowing, sensation of food being stuck in the throat or chest) AND • The patient has had an inadequate response after a minimum trial of 8 weeks to at least one of the following: swallowed topical corticosteroids (e.g. Budesonide) or high-dose proton inhibitor. • For continuation of therapy after the initial 6-month authorization, there must be documented improvement in EoE symptoms. <p><i>Diagnosis of COPD:</i></p> <ul style="list-style-type: none"> • The patient must have a diagnosis of COPD • The patient must have an eosinophilic count of ≥ 300 cells per mL within 12 months prior to initiation of therapy • The patient must have post-bronchodilator $FEV_1/FVC < 0.7$ AND FEV_1 30-70% of predicted • The patient has a history of uncontrolled disease, as indicated by ≥ 2 moderate or ≥ 1 severe exacerbation despite being on standard of care defined as triple therapy (LAMA+LABA+ICS) for at least 3 months prior to request, and at a stable dose for at least 1 month prior. Note: LAMA-LABA allowed if ICS is contraindicated. Pharmacy claims will be evaluated to assess compliance with therapy. AND • The prescriber is a pulmonologist • If member is current tobacco user, they must receive tobacco cessation counseling

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>Limitations: Dupixent®, Fasentra®, Nucala® and Cinqair® will not be considered in patients with a diagnosis of moderate to severe persistent asthma who are currently smoking or in combination with omalizumab or Tezepelumab.</p> <p>Tezspire:</p> <ul style="list-style-type: none"> • The patient must be 12 years of age or older AND <ul style="list-style-type: none"> • The patient has a history of uncontrolled asthma symptoms (symptoms occurring almost daily or waking at night with asthma at least once a week) or 2 or more exacerbations in the previous year despite regular use of medium-high dose ICS/LABA for a minimum of 3 consecutive months, with or without oral corticosteroids. Pharmacy claims will be evaluated to assess compliance with therapy. AND • The prescriber is an allergist, immunologist, or pulmonologist AND <ul style="list-style-type: none"> • If the patient has an eosinophilic phenotype (as defined by pretreatment blood eosinophil count of ≥ 150 cells per mL within the previous 6 weeks or ≥ 300 cells per mL within 12 months prior to initiation of therapy), there must have been a documented side effect, allergy, or treatment failure with Dupixent or Nucala AND • For continuation of therapy after the initial 6-month authorization, the patient must continue to receive therapy with an ICS/LABA AND have either a decreased frequency of exacerbations OR decreased use of maintenance oral corticosteroids OR reduction in the signs and symptoms of asthma OR an increase in predicted FEV1 from baseline. <p>Limitations: Tezspire will not be considered in patients who are currently smoking or in combination with anti-IgE, anti-IL4, or anti-IL5 monoclonal antibodies.</p>

IMMUNOSUPPRESANTS, ORAL

AZATHIOPRINE 50 MG tablet CYCLOSPORINE capsule CYCLOSPORINE MODIFIED MYCOPHENOLATE MOFETIL tablet, capsule, suspension MYCOPHENOLIC ACID delayed release tablet	Astagraf® XL (tacrolimus) capsule Azasan® (azathioprine) tablet Azathioprine 75 mg and 100 mg tablet Cellcept® (mycophenolate mofetil) tablet, capsule, suspension Envarsus® XR (tacrolimus) tablet Everolimus (compare to Zortress®) tablet Gengraf® (cyclosporine modified) capsule, solution	<p>Criteria (except Lupkynis and Rezero): The patient has been started and stabilized on the requested product OR the patient has a documented side effect, allergy, or treatment failure to a preferred agent (if a product has and AB rated generic, there must be a trial of the generic formulation).</p> <p>Lupkynis:</p> <ul style="list-style-type: none"> • The patient has a diagnosis of Systemic Lupus Erythematosus (SLE) AND
---	---	--

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
SIROLIMUS tablet TACROLIMUS capsule	Imuran® (azathioprine) tablet Lupkynis™ (voclosporin) capsule Myfortic® (mycophenolic acid) delayed release tablet Neoral® (cyclosporine modified) capsule, solution Prograf® (tacrolimus) capsule, granules for suspension Rapamune® (sirolimus) tablet, solution Rezurock™ (belumosudil) tablet Sandimmune® (cyclosporine) capsule, solution Zortress® (everolimus) tablet	<ul style="list-style-type: none"> The patient has active Lupus Nephritis confirmed by urine/blood tests or kidney biopsy AND The patient is ≥ 18 years of age AND Medication is prescribed by, or in consultation with, a nephrologist or rheumatologist AND The patient has clinical progression (e.g. worsening of proteinuria or serum creatinine) after 3 months of induction therapy with corticosteroids plus cyclophosphamide or mycophenolate mofetil OR failure to respond after 6 months of induction therapy with corticosteroids plus cyclophosphamide or mycophenolate mofetil AND Medication will be used in combination with background immunosuppressive therapy (e.g. mycophenolate mofetil and systemic corticosteroids) AND The patient has a documented intolerance or treatment failure with Benlysta <p>Rezurock:</p> <ul style="list-style-type: none"> The patient is ≥ 12 years of age AND The patient has a diagnosis of Chronic Graft-versus-host disease AND The patient has had a treatment failure with at least 2 prior courses of systemic immunosuppressant therapy (e.g. Corticosteroids, rituximab) AND The prescriber attests to monthly monitoring of liver function tests (total bilirubin, AST, and ALT)

CRYOPYRIN ASSOCIATED PERIODIC SYNDROMES (CAPS) AND PERIODIC FEVER SYNDROME (PFS)

All Products Require PA	Arcalyst® (rilonacept) <i>QTY LIMIT:</i> 2 vials for loading dose, then 1 vial per Week Ilaris® (canakinumab)	<p>Arcalyst: The diagnosis is Cryopyrin-Associated Periodic Syndrome (CAPS), including Familial Cold Autoinflammatory Syndrome (FCAS) or Muckle-Wells Syndrome (MWS) OR Maintenance of remission of Deficiency of Interleukin-1 Receptor Antagonist (DIRA) OR for treatment of recurrent pericarditis (RP) AND the patient’s age is FDA approved for the given indication.</p> <p>Ilaris: The diagnosis is Cryopyrin-Associated Periodic Syndrome (CAPS), including Familial Cold Autoinflammatory Syndrome (FCAS) or Muckle-Wells Syndrome (MWS) OR Familial Mediterranean Fever (FMF) OR Hyper-IgD periodic fever syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD) OR Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS) OR Active Still’s Disease including Adult-Onset Still’s Disease (AOSD) and Systemic Juvenile Idiopathic Arthritis (SJIA) AND the patient’s age is FDA approved for the given indication</p> <p>Note: Medical Records to support the above diagnosis must accompany the Prior Authorization request. Authorization for continued use shall be reviewed at least every 12 months to confirm patient has experienced disease stability or improvement while on therapy.</p>
-------------------------	---	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

IRON CHELATING AGENTS

DEFERASIROX tablet	Deferasirox dispersible tablet, granule pack Deferiprone tablet Exjade® (deferasirox) dispersible tablet Ferriprox® (deferiprone) tablet, solution Jadenu®(deferasirox) tablet, granule pack	<p>Deferasirox dispersible tablet, Exjade dispersible tablet: The patient has a medical necessity for a non-solid oral dosage form AND for approval of Exjade, the patient has a documented intolerance to generic deferasirox dispersible tablets.</p> <p>Deferiprone tablet, Ferriprox tablet, Jadenu tablet: the patient has a documented intolerance to generic deferasirox tablets</p> <p>Deferasirox granule pack, Ferriprox solution, Jadenu granule pack: The patient has a medical necessity for a non-solid oral dosage form AND The patient has a documented intolerance to generic deferasirox dispersible tablets.</p>
--------------------	--	--

LIPOTROPICS

BILE ACID SEQUESTRANTS

CHOLESTYRAMINE powder (compare to Questran®) CHOLESTYRAMINE LIGHT powder (compare to Questran Light®) COLESTIPOL tablets (compare to Colestid®) WELCHOL® (colesevelam) tablets, powder packets	Colesevelam (compare to Welchol®) Colestid® tablets, granules (colestipol) Colestipol granules, packets Prevalite powder (cholestyramine light) Questran® powder (cholestyramine) Questran Light® powder (cholestyramine light)	<p>Colesevelam: The patient has had a documented intolerance to the brand name equivalent.</p> <p>Colestipol granules, packets: The patient has a documented side effect, allergy, or treatment failure with two preferred bile acid sequestrants.</p> <p>Prevalite, Questran, Questran Light, Colestid: The patient has had a documented intolerance to the preferred generic formulation.</p>
---	--	--

FIBRIC ACID DERIVATIVES

GEMFIBROZIL (compare to Lopid®) 600 mg FENOFIBRATE MICRONIZED CAPSULE (compare to Lofibra® capsules) 67 mg, 134 mg, 200 mg FENOFIBRATE NANOCRYSTALIZED (compare to Tricor®) 48 mg, 145 mg tablets FENOFIBRATE TABLETS (compare to Lofibra® tablets) 54 mg, 160 mg	Antara® (fenofibrate micronized) 30 mg, 43 mg, 90 mg, 130 mg F Fenofibrate capsule (compare to (Lipofen®) 50 mg, 150 mg Fenofibrate micronized (compare to Antara®) 43 mg, 130 mg Fenofibric acid (compare to Trilipix) 45 mg, 135 mg delayed release capsule Fenofibric acid 35 mg, 105 mg <i>QTY LIMIT:</i> 1 capsule/day Fenoglide® (fenofibrate MeltDose) 40 mg, 120 mg Lipofen® (fenofibrate) 50 mg, 150 mg	<p>Lopid: The patient has had a documented intolerance to generic gemfibrozil.</p> <p>All other non-preferred medications: The patient has had a documented side effect, allergy, or treatment failure with two preferred fibric acid derivatives (If a product has an AB rated generic, there must have been a trial with the generic formulation.)</p>
--	--	--

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
	Lopid [®] (gemfibrozil) 600 mg Tricor [®] (fenofibrate nanocrystallized) 48 mg, 145 mg Trilipix (fenofibric acid) 45 mg, 135 mg delayed release capsule	
MISC. HOMOZYGOUS FAMILIAL HYPERCHOLESTEROLEMA (HoFH) AGENTS		
All products require PA	Evkeeza [™] (evinacumab-dgnb) intravenous solution Juxtapid [®] (lomitapide) Capsule <i>QTY LIMIT:</i> 5 and 10 mg caps = 1/day, 20 mg cap = 3/day Maximum day supply per fill is 28 days	CRITERIA FOR APPROVAL: Total cholesterol levels > 290mg/dL or LDL-C > 190mg/dL (adults) OR Total cholesterol levels > 260mg/dL or LDL-C > 155mg/dL (children < 16 years) and TG within reference range or Confirmation of diagnosis by gene testing AND Documented adherence to prescribed lipid lowering medications for the previous 90 days AND Recommended or prescribed by a lipidologist or Cardiologist AND Inability to reach goal LDL-C despite a trial of 2 or more maximum tolerated dose of statins (one of which must be atorvastatin or rosuvastatin), ezetimibe 10mg daily, and Repatha
NICOTINIC ACID DERIVATIVES		
NIACIN NIACIN extended release		
STATINS		
ATORVASTATIN (compare to Lipitor [®]) LOVASTATIN PRAVASTATIN ROSUVASTATIN (compare to Crestor [®]) SIMVASTATIN (compare to Zocor [®]) Note: All preferred agents have a quantity limit of 1 tablet/day except Lovastatin 40mg which has a quantity limit of 2 tablets/day	Altoprev [®] (lovastatin SR) Atorvaliq [®] (atorvastatin) oral suspension Crestor [®] (rosuvastatin) Ezallor [®] (rosuvastatin) sprinkle capsule Fluvastatin Fluvastatin ER (compare to Lescol [®] XL) Lescol [®] XL (fluvastatin ER) Lipitor [®] (atorvastatin) Livalo [®] (pitavastatin) Zocor [®] (simvastatin) Zypitamag [™] (pitavastatin) Note: All non-preferred agents have a quantity limit of 1 tablet/day except fluvastatin IR which has a quantity limit of 2 tablets/day.	Non-preferred agents (except as noted below): The patient must have a documented side effect, allergy, or treatment failure to 3 preferred statins. If the product has an AB rated generic, one trial must be the generic formulation. Atorvaliq, Ezallor: medical necessity for a specialty dosage form has been provided. Zypitamag: The patient must have a documented side effect, allergy, or treatment failure to 3 preferred statins AND clinical justification is provided documenting why the patient is unable to use Livalo. LIMITATIONS: Simvastatin 80 mg: initiation of simvastatin 80 mg or titration to 80 mg is not recommended by the FDA due to the increased risk of myopathy, including rhabdomyolysis. Patients may only continue on this dose when new to Medicaid if the patient has been taking this dose for 12 or more months without evidence of muscle toxicity. If the request is for Zocor 80 mg, the patient must have met the prior treatment length requirement and have a documented intolerance to the generic equivalent.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
MISCELLANEOUS/COMBOS		
<p>Ezetimibe (compare to Zetia®) <i>QTY LIMIT: 1 tab/day</i></p> <p>Omega-3-acid ethyl esters (compare to Lovaza®)</p> <p>Ezetimibe/simvastatin (compare to Vytorin®) 10/10 mg, 10/20mg, and 10/40mg <i>QTY LIMIT: 1 tab/day</i></p>	<p>Amlodipine/atorvastatin (compare to Caduet®) <i>QTY LIMIT: 1 tab/day</i></p> <p>Caduet® (atorvastatin/amlodipine) <i>QTY LIMIT: 1 tab/day</i></p> <p>Ezetimibe/simvastatin (compare to Vytorin®) 10/80mg strength only</p> <p>Icosapent Ethyl (compare to Vascepa®) <i>QTY LIMIT: 4 caps/day</i></p> <p>Lovaza® (omega-3-acid ethyl esters)</p> <p>Omega-3-acid ethyl esters (compare to Lovaza®)</p> <p>Nexletol® (bempedoic acid) <i>QTY LIMIT: 1 tab/day</i></p> <p>Nexlizet® (bempedoic acid/ezetimibe) <i>QTY LIMIT: 1 tab/day</i></p> <p>Vascepa® (icosapent ethyl) <i>QTY LIMIT: 4 caps/day</i></p> <p>Vytorin® (ezetimibe/simvastatin) <i>QTY LIMIT: 1 tab/day</i></p> <p>Zetia® (ezetimibe) <i>QTY LIMIT: 1 tab/day</i></p>	<p>Lovaza, Vytorin, Zetia: patient must have a documented intolerance to the generic equivalent.</p> <p>Icosapent Ethyl, Vascepa: <i>Indication for use is severe hypertriglyceridemia:</i></p> <ul style="list-style-type: none"> The patient has pre-treatment triglyceride levels > 500 mg/dL AND The patient has a documented contraindication, side effect, allergy, or treatment failure to Omega-3-acid ethyl esters. <p><i>Indication for use is cardiovascular risk reduction:</i></p> <ul style="list-style-type: none"> The patient has pre-treatment triglyceride levels > 150 mg/dL AND The patient is receiving adjunct therapy with a maximally tolerated high intensity statin AND For approval of icosapent ethyl, the patient has had a documented intolerance to brand Vascepa <p>Amlodipine/atorvastatin, Caduet: The patient is unable to take the individual separate agents AND for approval of Caduet, the patient must have also had a documented intolerance to the generic equivalent.</p> <p>Nexletol, Nexlizet: The patient has had an inadequate response to a 3-month trial of atorvastatin or rosuvastatin OR Patient has demonstrated statin intolerance as defined by statin-related rhabdomyolysis or skeletal related muscle symptoms AND Patient (if eligible) will continue adjunct therapy with maximally tolerated high intensity statin. If patient is using simvastatin, dose should not exceed 20 mg/day; if patient is using pravastatin, dose should not exceed 40 mg/day</p> <p>Ezetimibe/simvastatin (10/80): the patient has been taking this dose for 12 or more months without evidence of muscle toxicity.</p>
PCSK9 INHIBITORS		
<p><u>Preferred After Clinical Criteria Are Met</u></p> <p>PRALUENT® (alirocumab) <i>QTY LIMIT: 2ml (75 mg injection every 2 weeks or 300 mg every month)/28 days</i> Max 28-day supply</p> <p>REPATHA® (evolocumab) Sureclick, prefilled syringe <i>QTY LIMIT: 2ml (2 injections)/28 days</i> Max 28-day supply</p> <p>REPATHA® (evolocumab) Pushtronix™ <i>QTY LIMIT: 3.5ml (One single-use infusor and prefilled cartridge)/28 days, Max 28-day supply</i></p>	<p>Leqvio® (inclisiran) prefilled syringe</p>	<p>Criteria for approval:</p> <ul style="list-style-type: none"> The patient's age is FDA approved for the given indication AND Concurrent use with statin therapy AND Documented adherence to prescribed lipid lowering medications for the previous 90 days AND Inability to reach goal LDL-C despite a trial of 2 or more maximum tolerated dose of statins (one of which must be atorvastatin or rosuvastatin) For approval of Leqvio, the patient must have a documented side effect, allergy, or treatment failure (defined as inability to get within 10% of stated LDL-C goal, not to exceed guideline recognized goals) with a minimum 12-week trial of both Praluent and Repatha.

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

MISCELLANEOUS

KUVAN® (sapropterin) 100mg, 500mg powder
PYRIDOSTIGMINE BROMIDE
SAPROPTERIN 100mg powder
TRANEXAMIC ACID (compare to Lysteda®)
QTY LIMIT: 30 tablets/28 days
FENSOLVI® (leuprolide acetate) subcutaneous injection
QTY LIMIT: 1 vial every 6 months

Preferred After Clinical Criteria Are Met
CARBAGLU® dispersible tablets (carglumic acid)

CRYSVITA® (burosumab-twza)
FABRAZYME (agalsidase beta) IV

Brineura™ (cerliponase alfa)
QTY LIMIT: 1 package per 14 days (Brineura Injection, 2 vials of 150mg/5ml, and Intraventricular Electrolytes Injection, 1 vial of 5ml)

Carglumic acid (compare to Carbaglu®) dispersible tablets

Daybue™ (trofinetide) solution
QTY LIMIT: 120 mL/day

Eohilia™ (budesonide oral suspension)

Elaprase® (idursulfase)
QTY LIMIT: calculated dose/week

Firdapse® (amifampridine)
QTY LIMIT: 8 tablets/day

Galafold™ (migalastat)
QTY LIMIT: 14 caps/28 days
Maximum day supply = 28 days

Gamifant® (emapalumab-lzsg)
Hyftor™ (sirolimus) topical gel

Korsuva® (difelikefalin)
Kuvan (sapropterin) tablets

Hydroxyprogesterone caproate 250 mg/ml vial (intramuscular injection)

Jylamvo® (methotrexate) oral solution
Luxturna® (voretigine neparvovec-rzyl) suspension for subretinal injection
QTY LIMIT: one injection per eye per lifetime

Lysteda® tablets (tranexamic acid)
QTY LIMIT: 30 tablets/28 days

Mestinon®

Myalept® (metreleptin) vial for subcutaneous injection
QTY LIMIT: one vial/day
Maximum day supply per fill = 30 days

Oxlumo™ (lumasiran)

Palynziq™ (pegvaliase-pqpz)

Rivfloza® (nedosiran) injection

Ruzurgi® (amifampridine)
QTY LIMIT: 10 tablets/day

Brineura:

- Patient is 3 years of age or older AND
- The diagnosis or indication is late infantile neuronal ceroid lipofuscinosis type 2 (CLN2), confirmed by deficiency of the lysosomal enzyme tripeptidyl peptidase-1 (TPP1) (results of genetic testing must be submitted AND
- The prescriber is a neurologist or other physician specializing in intraventricular administration

Note: Brineura will be approved as a medical benefit ONLY and will NOT be approved if billed through pharmacy point of sale. Initial approval will be granted for 3 months. Renewal may be granted for up to 12 months. For therapy continuation, clinical documentation must be submitted documenting improvement or maintenance of motor ability OR slower progression of disease than would otherwise be expected AND a 12-lead ECG evaluation is performed every 6 months.

Carbaglu, Carglumic Acid: The diagnosis or indication for the requested medication is hyperammonemia due to N-acetylglutamate synthetase (NAGS) deficiency, propionic acidemia, or methylmalonic acidemia AND The prescriber is a specialist in metabolic disorders (e.g., medical geneticist) or prescriber is in consultation with a specialist AND **for approval of generic product**, the patient has had a documented intolerance to the generic equivalent of the requested medication.

Crysvita:

- Patient is ≥ 1 year of age AND
- Patient has a diagnosis of X-linked hypophosphatemia AND
- Medication is prescribed by or in consultation with an endocrinologist or nephrologist AND
- Patient has not received oral phosphate or vitamin D analogs within 1 week prior to starting therapy AND
- Baseline fasting serum phosphorous level is below the lower limit of the laboratory normal reference range AND Patient does not have severe renal impairment, defined as a GFR of <30 mL/min AND

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
	<p>Sapropterin (compare to Kuvan®) tablets, 500mg powder</p> <p>Skyclarys® (omaveloxolone) <i>QTY LIMIT: 3 capsules/day</i></p> <p>Sohonos™ (palovarotene)</p> <p>Tepezza® (teprotumumab-trbw) vial for IV infusion</p> <p>Veozah™ (fezolinetant) tablet <i>QTY LIMIT: 1 tablet/day</i></p> <p>Vyvgart® (efgartigimod alfa-fcab) IV solution</p> <p>Vyvgart® Hytrulo (efgartigimod alfa and hyaluronidase-qvfc human recombinant injection) SC solution</p> <p>Xolremdi™ (mavorixafor) capsule</p> <p>Xatmep™ (methotrexate) oral solution</p> <p>Zokinvy® (lonafarnib) capsule</p>	<ul style="list-style-type: none"> Dose does not exceed 90mg every 14 days (pediatrics) or 90mg every 28 days (adults) <p>Note: Initial approval will be granted for 6 months. Renewal may be granted for up to 1 year. For therapy continuation, patient must have disease response as indicated by one of the following:</p> <ul style="list-style-type: none"> Increased serum phosphate levels, not exceeding the upper limit of the laboratory normal range. A reduction in serum total alkaline phosphatase activity. Improvement in symptoms (e.g. skeletal pain, linear growth, etc.). Improvement in radiographic imaging of Rickets/osteomalacia. <p>Daybue:</p> <ul style="list-style-type: none"> The patient is ≥ 2 years of age. The prescription is initiated by or in consultation with a neurologist or other developmental specialist. The patient has a diagnosis of typical Rett syndrome per the Rett Syndrome Diagnostic Criteria (must meet ALL): <ul style="list-style-type: none"> Partial or complete loss of acquired purposeful hand skills. Partial or complete loss of acquired spoken language. Gait abnormalities: Impaired (dyspraxia) or absence of ability. Stereotypic hand movements such as hand wringing/squeezing, clapping/tapping, mouthing, and washing/rubbing automatisms. The patient does not have any of the Exclusion Criteria: <ul style="list-style-type: none"> Brain injury secondary to trauma (peri- or postnatally), neurometabolic disease, or severe infection that causes neurological problems. Grossly abnormal psychomotor development in first 6 months of life The patient has a documented disease-causing mutation in the <i>MECP2</i> gene. The patient is not using any insulin. Detailed clinical baseline has been provided using an objective measure or tool (Rett Syndrome Behavior Questionnaire (RSBQ)). Initial approval will be granted for 3 months. For reapproval, the patient must have a documented clinical improvement in disease as evidenced by ≥ 10% reduction in the RSBQ questionnaire score. Patients with a baseline RSBQ score of ≤ 30 must have at least a ≥ 3-point reduction AND The patient has not experienced significant weight loss (>5% from baseline). <p>Elaprased (Hunter's Syndrome Injectable): The diagnosis or indication for the requested medication is Hunter's Syndrome</p> <p>Eohilia: The patient has had a documented side effect, allergy, treatment failure with swallowed budesonide nebulizer solution or fluticasone. Note: Approval will be granted for 12 weeks of therapy.</p> <p>Fabrazyme: Diagnosis or indication is Fabry Disease.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>Firdapse, Ruzurgi: patient has a diagnosis of Lambert-Eaton myasthenic syndrome (LEMS) AND prescription is initiated by or in consultation with a neurologist AND patient does not have a history of seizures AND for approval of Firdapse, the patient must have a documented intolerance to Ruzurgi. Initial approval will be granted for 3 months with documentation of the patient's baseline clinical muscle strength assessment using a standardized rating scale. For re-approval after 3 months, the patient must have improved, or stable symptoms documented with the appropriate standardized rating scale</p> <p>Galafold: Patient is ≥ 18 years of age AND Diagnosis or indication is Fabry Disease with an amenable galactosidase alpha (GLA) gene variant for treatment (results must be submitted) AND enzyme replacement therapy is not a therapeutic option (e.g. due to allergy, hypersensitivity, or poor venous access).</p> <p>Gamifant: the patient has a diagnosis of primary hemophagocytic lymphohistiocytosis (HLH) with refractory, recurrent, or progressive disease or intolerance with conventional HLH therapy (e.g. etoposide + dexamethasone) AND the patient is a candidate for a stem cell transplant AND Gamifant will be administered in combination with dexamethasone</p> <p>Hyftor: The patient has 3 or more angiofibromas (≥ 2mm in diameter with redness in each) on the face, associated with tuberous sclerosis AND the patient has completed all ACIP recommended age-appropriate vaccinations prior to starting therapy. Initial approval will be granted for 3 months. For re-approval, there must be documented reduction in the size and redness of angiofibromas from baseline.</p> <p>Korsuva: The patient has a diagnosis of moderate-to-severe pruritis associated with chronic kidney disease AND the patient is receiving hemodialysis AND the patient has a documented side effect, allergy, or treatment failure with at least 1 topical and 1 systemic pruritis treatment (e.g. antihistamines, corticosteroids, gabapentin, pregabalin, capsaicin)</p> <p>Kuvan tabs, Sapropterin tabs: patient has a documented intolerance to the powder formulation.</p> <p>Luxturna: patient must have inherited retinal dystrophy due to mutations in both copies of the RPE65 gene (results of genetic testing must be submitted) AND patient has sufficient viable retinal cells as determined by the treating physician(s) AND Luxturna will be administered by a retinal specialist/surgeon experienced in performing intraocular surgery and associated with an Ocular Gene Therapy Treatment Center.</p> <p>Lysteda the patient has had a documented intolerance to the generic product.</p> <p>Myalept: Patient has a diagnosis of congenital or acquired generalized lipodystrophy AND Patient has one or more of the following metabolic</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>abnormalities AND is refractory to current standards of care for lipid and diabetic management: Insulin resistance (defined as requiring > 200 units per day), Hypertriglyceridemia, Diabetes AND Prescription is written by or in consultation with an endocrinologist AND The prescriber is registered in the MYALEPT REMS program. Reauthorization for continued use criteria: Patient has experienced an objective response to therapy • Sustained reduction in hemoglobin A1c (HbA1c) level from baseline OR • Sustained reduction in triglyceride (TG) levels from baseline.</p> <p>Oxlumo, Rivfloza: The patient has a diagnosis of Primary Hyperoxaluria Type I (PH1) confirmed via genetic testing (identification of alanine: glyoxylate aminotransferase gene (AGXT) mutation) AND urinary oxalate excretion > 0.5mmol/1.73 m² or urinary oxalate: creatinine ratio is above the upper limit of normal for age AND medication is being prescribed by, or in consultation, with a nephrologist or urologist AND patient has not previously received a liver transplant. For approval of Rivfloza: patient must be at least 9 years of age and have relatively preserved kidney function (eGFR ≥30ml/min/1.73m²).</p> <p>Palynziq: Patient is 18 years of age or older AND has a diagnosis of phenylketonuria AND has uncontrolled blood phenylalanine (PHE) concentrations (> 600 micromol/L) on existing management, including restricting dietary phenylalanine and protein intake and treatment with sapropterin. For re-approval, the patient must have achieved at least a 20% reduction in PHE concentration from pre-treatment baseline or a PHE ≤ 600 micromol/L after 16 weeks of continuous treatment with the maximum dosage of 40mg daily. Note: Palynziq has a black box warning for anaphylaxis which can occur at any time during treatment. Patients, pharmacies, and physicians must be enrolled in the Palynziq REMS program AND concurrent auto-injectable epinephrine must be prescribed.</p> <p>Note: Initial approval will be granted for 4 loading doses (the first 3 loading doses should be administered at 14-day intervals; the 4th loading dose should be administered 30 days after the 3rd dose). Renewal may be granted for up to 12 months with a maximum of 3 doses approved per year (12mg(5ml) every 4 months). For therapy continuation, clinical documentation must be submitted documenting improvement or maintenance of motor ability OR slower progression of disease than would otherwise be expected.</p> <p>Sapropterin 500mg powder: patient has a documented intolerance to brand Kuvan</p> <p>Skyclarys:</p> <ul style="list-style-type: none"> • The patient is ≥ 16 years of age • The patient has a diagnosis of Friedreich’s Ataxia with a confirmed mutation in the frataxin (FXN) gene AND • The patient must have a stable modified Friedreich’s Ataxia Rating Scale (mFARS) score between 20 and 80, be able to complete maximal exercise testing, and have a left ventricular ejection fraction of at least 40%. • Baseline liver function tests will be completed prior to start of therapy and

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

		<p>continued monthly for 3 months following Skyclarys administration AND</p> <ul style="list-style-type: none"> Baseline B-type natriuretic peptide (BNP) will be obtained, and level does not exceed 200pg/mL AND the patient has no history of clinically significant cardiac disease AND <p>Reauthorization Criteria:</p> <ul style="list-style-type: none"> For continuation, there must be stable or slower progression of the disease than would otherwise be expected <p>Sohonos: The patient has a diagnosis of fibrodysplasia ossificans progressive (FOP) AND has a confirmed R206H mutation in the activin receptor IA (ACVR1) gene AND patient is a female ≥ 8 years of age or a male ≥ 10 years of age. Initial approval will be authorized for 1 year. For re-approval, the patient must have documentation of a positive response to therapy defined as a reduction in new heterotopic ossification (HO) symptoms.</p> <p>Tepezza:</p> <ul style="list-style-type: none"> Patient has a diagnosis of Thyroid Eye Disease (TED) related to Graves' Disease AND Patient has a baseline Clinical Activity Score (CAS) ≥ 4 in the most severely affected eye AND Patient has active TED associated with at least one of the following: <ul style="list-style-type: none"> Lid retraction ≥ 2 mm Moderate or severe soft tissue involvement Exophthalmos ≥ 3 mm above normal for race and gender Diplopia (double vision) Patient is euthyroid, defined as free triiodothyronine (T3) and thyroxine (T4) levels within the normal limits, OR Patient has free T3 and T4 levels less than 50% above or below the normal limits and is undergoing treatment to correct the hypo- or hyperthyroidism to maintain a euthyroid state AND Patient has had an inadequate response or contraindication to high-dose intravenous glucocorticoid therapy. <p>Veozah: The indication for use is moderate to severe vasomotor symptoms (VMS) associated with menopause AND documentation has been provided detailing the frequency and severity of these symptoms AND the patient has had a documented side effect, allergy, contraindication, or treatment failure, defined by at least 4 weeks of therapy, to one preferred Hormone Replacement Therapy (HRT) and two preferred non-hormonal therapies (i.e., SSRIs, SNRIs, gabapentin, pregabalin, clonidine). For re-approval, there must be a documented improvement in the frequency or severity of VMS.</p> <p>Vyvgart, Vyvgart Hytrulo:</p> <ul style="list-style-type: none"> Patient is ≥ 18 years of age AND Patient has a diagnosis of generalized Myasthenia Gravis with Myasthenia Gravis Foundation of America (MGFA) clinical classification class II to IV AND Patient is anti-acetylcholine receptor (AChR) positive AND MG-Activities of Daily Living (MG-ADL) total score of ≥ 5 at baseline AND
--	--	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<ul style="list-style-type: none"> • Patient has IgG levels of at least 6g/L AND • Patient has had an inadequate response with at least 2 immunosuppressive therapies (e.g. corticosteroids, azathioprine, cyclosporine, mycophenolate) over the course of at least 12 months AND • Maximum of four doses per 50 days AND • For approval of Vyvgart Hytrulo, the prescriber must provide a clinically compelling reason for the use of the requested medication including reasons why Vyvgart IV would not be a suitable alternative. • For re-approval, the patient must have had a positive response to therapy as evidenced by a 2-point reduction in the MG-ADL score. <p>Jylamvo, Xatmep: The patient has a diagnosis consistent with the FDA indication of the requested drug AND patient has a requirement for an oral liquid dosage form (i.e. swallowing disorder, inability to take oral medication)</p> <p>Xolremdi: The patient meets the FDA approved age AND has a diagnosis of WHIM syndrome confirmed by genetic confirmation of a CXCR4 variant AND has a baseline absolute neutrophil count (ANC) ≥ 400 cells/uL</p> <p>Zokinvy: The patient meets FDA approved age and BSA AND the patient has a diagnosis of Hutchinson-Gilford Progeria Syndrome (HGPS) OR the patient has a diagnosis of processing-deficient Progeroid Laminopathies with documentation of either Heterozygous LMNA mutation with progerin-like protein accumulation or Homozygous or compound heterozygous ZMPSTE24 mutations.</p> <p>Note: A single-dose of 10mg/kg will be approved per active CDI. A repeat dose will not be approved for recurrence of the same active infection</p>

AMYOTROPHIC LATERAL SCLEROSIS (ALS)		
RILUZOLE (Compare to Rilutek®)	Exservan™ (riluzole) film Qalsody®(tofersen) injection <i>QTY LIMIT:</i> 100 mg (15 ml) every 14 days x 3 doses and 100 mg (15 ml)/28 days thereafter Radicava® (edaravone) IV injection Rilutek® (riluzole) Tiglutik™ (riluzole) suspension	<p>Exservan, Tiglutik: patient must be unable to take whole or crushed Riluzole tablets</p> <p>Qalsody:</p> <ul style="list-style-type: none"> • The diagnosis is amyotrophic lateral sclerosis (ALS) AND • Documentation has been provided indicating the presence of a mutation in the superoxide dismutase 1 (SOD1) gene AND • The patient is ≥ 18 years old AND • Patient has a slow vital capacity (%SVC) spirometry test $\geq 50\%$ of predicted as adjusted for sex, age, and height at screening. AND • Patient is not dependent on invasive ventilation or tracheostomy AND • Baseline ALS Functional Rating Scale-Revised (ALSFRS-R) total score has been completed AND • Initial approval will be granted for 6 months. For re-approval there must be documented response to therapy compared to baseline as evidenced by either stable or slowing decline on ALSFRS-R rating scale (patient has not experienced a rapid disease progression while on therapy). <p>Radicava:</p> <ul style="list-style-type: none"> ○ The diagnosis is amyotrophic lateral sclerosis (ALS) AND

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<ul style="list-style-type: none"> ○ Disease duration is ≤ 2 years AND ○ Patient has functionally retained most activities of daily living AND ○ Patient has normal respiratory function (defined as a % predicted forced vital capacity of ≥ 80%) AND ○ Patient does not have a sulfite allergy AND ○ Initial approval will be granted for 14 doses/28 days and all subsequent approvals will be for 10 doses/28 days <p>Rilutek: Patient must have a documented intolerance with riluzole.</p>

COMPLEMENT INHIBITORS		
<p>All products require PA</p>	<p>Enjaymo™ (sutimlimab-jome) Empaveli™ (pegcetacoplan) subcutaneous solution <i>QTY LIMIT: 8 vials/28 days</i> Soliris® (eculizumab) vial Ultomiris® (ravulizumab-cwvz) Voydeya™ (danicopan) Zilbrysq® (zilucoplan)</p>	<p>Enjaymo: The patient has a diagnosis of cold agglutinin syndrome (CAD) AND the patient does not have an active chronic systemic infection (e.g. Hepatitis B, Hepatitis C, HIV) AND the medication is prescribed by, or in consultation with, a hematologist AND the patient has had at least one blood transfusion in the 6 months prior to starting Enjaymo AND the patient has received the pneumococcal, Haemophilus influenzae, and meningococcal vaccines at least 2 weeks prior to therapy initiation. Authorization for continued use shall be reviewed to confirm that the patient has experienced an objective response to the therapy (e.g. stabilization of hemoglobin levels, reductions in transfusions, improvement in hemolysis, etc.)</p> <p>Empaveli: The patient has a diagnosis of paroxysmal nocturnal hemoglobinuria (PNH) documented by flow cytometry AND The patient has received the meningococcal vaccine at least 2 weeks prior to therapy initiation. Authorization for continued use shall be reviewed to confirm that the patient has experienced an objective response to the therapy (e.g. stabilization of hemoglobin levels, reductions in transfusions, improvement in hemolysis, etc.). Note: For patients switching from eculizumab, an additional 4 weeks of eculizumab will be approved before continuing monotherapy with Empaveli. For patients switching from ravulizumab, Empaveli will be initiated no more than 4 weeks after the last dose of ravulizumab. Ongoing combination therapy of complement inhibitors will not be approved.</p> <p>Soliris: <i>Indication for use is Atypical Hemolytic Uremic Syndrome:</i> Dose requested must be within the FDA parameters for loading and maintenance dose <i>Indication for use is paroxysmal nocturnal hemoglobinuria (PNH):</i> Diagnosis is documented by flow cytometry AND The patient has received the meningococcal vaccine at least 2 weeks prior to therapy initiation. Authorization for continued use shall be reviewed to confirm that the patient has experienced an objective response to the therapy (e.g. stabilization of hemoglobin levels, reductions in transfusions, improvement in hemolysis, etc.) <i>Indication for use is Myasthenia Gravis:</i> The patient is anti-acetylcholine receptor</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

		<p>(AChR) antibody positive AND the patient has a documented side effect, allergy, or treatment failure with at least 2 immunosuppressive therapies (e.g. corticosteroids, azathioprine, cyclosporine, mycophenolate, etc.).</p> <p>Ultomiris: The patient has a diagnosis of Atypical Hemolytic Uremic Syndrome or a diagnosis of paroxysmal nocturnal hemoglobinuria (PNH) documented by flow cytometry AND The patient has received the meningococcal vaccine at least 2 weeks prior to therapy initiation. Authorization for continued use shall be reviewed to confirm that the patient has experienced an objective response to the therapy (e.g. stabilization of hemoglobin levels, reductions in transfusions, improvement in hemolysis, etc.) Note: Dose requested must be within the weight-based parameters for loading and maintenance dose</p> <p>Voydeya: The patient has a diagnosis of extravascular hemolysis (EVH) with paroxysmal nocturnal hemoglobinuria (PNH) AND there has been disease progression despite being on a stable dose of ravulizumab or eculizumab AND the patient will remain on ravulizumab or eculizumab (only approved as add-on therapy) Note: requires provider to be enrolled in REMS program.</p> <p>Zilbrysq; The patient is ≥ 18 years of age AND</p> <ul style="list-style-type: none"> • Patient has a diagnosis of generalized Myasthenia Gravis with Myasthenia Gravis Foundation of America (MGFA) clinical classification class II and IV AND • Patient is anti-acetylcholine receptor (AChR) positive AND • MG-Activities of Daily Living (MG-ADL) total score of ≥ 6 at baseline AND • Patient has had an inadequate response with at least 2 immunosuppressive therapies (e.g. corticosteroids, azathioprine, cyclosporine, mycophenolate) over the course of at least 12 months. • For re-approval, the patient must have had a positive response to therapy as evidenced by a 2-point reduction in the MG-ADL score or QMG score.
--	--	--

GLYCOPYRROLATE		
GLYCOPYRROLATE 1 mg, 2 mg tablets (compare to Robinul [®] , Robinul Forte [®])	Cuvposa [®] oral solution (glycopyrrolate) Maximum days supply per fill is 30 days Dartisla ODT [™] (glycopyrrolate) <i>QTY LIMIT = 4 tabs/day</i> Glycopyrrolate 1mg/5ml oral solution (compare to Cuvposa) Robinul [®] (glycopyrrolate) 1mg Robinul [®] Forte (glycopyrrolate) 2mg	<p>Cuvposa, Glycopyrrolate oral solution: The patient has medical necessity for a non-solid oral dosage form OR the dose cannot be obtained from the tablet formulation.</p> <p>Dartisla ODT: The patient has been established on the 2mg dosage strength of another form of glycopyrrolate AND the patient has a documented intolerance to glycopyrrolate tablets and solution.</p> <p>Robinul, Robinul Forte: The patient has a documented intolerance to glycopyrrolate tablets.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
INJECTABLE METHOTREXATE		
METHOTREXATE 25 MG/ML solution for injection	Otrexup® or Rasuvo® Single-dose auto-injector for subcutaneous use (methotrexate) QTY LIMIT: 4 syringes/28 days RediTrex® Prefilled syringe for subcutaneous use (methotrexate) QTY LIMIT: 4 syringes/28 days	Otrexup, Rasuvo, Reditrex: The patient has a diagnosis of rheumatoid arthritis (RA), polyarticular juvenile idiopathic arthritis (pJIA) or psoriasis. AND The patient has been intolerant to oral methotrexate AND The patient has been unable to be compliant with a preferred form of injectable methotrexate (includes difficulty with manual dexterity)
Immunoglobulin A Nephropathy (IgAN) Agents		
All products require PA	Filspari™ (sparsentan) tablet QTY LIMIT: 1 tablet/day Tarpeyo™ (budesonide) delayed release capsule	Filspari, Tarpeyo: <ul style="list-style-type: none"> • The patient has a diagnosis of Immunoglobulin A Nephropathy (IgAN) confirmed by biopsy AND • eGFR \geq is 30ml/min/1.73m² (Filspari) or eGFR \geq is 35ml/min/1.73m² (Tarpeyo) AND • The patient meets one of the following: Proteinuria \geq 1g/day or Urine protein-to-creatinine ratio (UPCR) \geq 1.5 g/g AND • The patient is on a stable dose of maximally tolerated ACE-I or ARB therapy for a minimum of 3 months AND • The patient's kidney function has continued to decline despite treatment with a preferred oral corticosteroid AND • Duration of therapy does not exceed 9 months (Tarpeyo only) • The prescriber, patient, and pharmacy are enrolled in the REMS program (Filspari only)
MINERALOCORTICOID RECEPTOR ANTAGONISTS		
EPLERENONE SPIRONOLACTONE	Aldactone® (spironolactone) CaroSpir® (spironolactone) oral suspension Inspra® (eplerenone) Kerendia® (finerenone)	Aldactone, Inspra: The patient has a documented intolerance to the generic formulation Carospir: patient has a medical necessity for a specialty dosage form (i.e. dysphagia, swallowing disorder). Kerendia: The patient has a diagnosis of chronic kidney disease (CKD) associated with Type II Diabetes AND the estimated glomerular filtration rate at baseline is \geq 25 mL/min/1.73m ² AND the urine albumin-to-creatinine ratio is \geq 30mg/g AND the patient is currently receiving, or has a contraindication to, an ACE inhibitor or angiotensin receptor blocker (ARB)
NEUROMYELITIS OPTICA SPECTRUM DISORDERS (NMOSD)		
All Products Require PA	Enspryng® (satralizumab-mwge) prefilled syringe QTY LIMIT = 3/28 days for the first month then 1/28 days thereafter Soliris® (eculizumab) vial Uplizna® (inebilizumab-cdon) vial QTY LIMIT = 300mg x 2 doses for the first 2 weeks then 300mg every 6 months thereafter	Enspryng, Soliris, Uplizna: <ul style="list-style-type: none"> • The patient is \geq 18 years AND • Diagnosis or indication is the treatment of neuromyelitis optica spectrum disorder (NMOSD) AND • Patient is anti aquaporin-4 (AQP4) antibody positive AND • Patient has a history of one or more relapses that required rescue therapy within the year prior to screening, or 2 or more relapses that required rescue therapy in 2 years prior to screening AND • Patient must have a documented side effect, allergy, treatment failure, or contraindication to rituximab. • Initial approval will be granted for 6 months. Renewal requires documentation of improvement or stabilization of neurologic symptoms such as a decrease in acute relapses, reduced hospitalization, or reduction

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>in plasma exchange treatments. Soliris, Uplizna additional criteria: The patient must have a documented side effect, allergy, treatment failure or contraindication to Enspryng.</p>
SOMATOSTATIN ANALOGS		
<p>OCTREOTIDE ACETATE solution for injection SANDOSTATIN® (octreotide acetate) LAR Depot</p>	<p>Bynfezia® (octreotide) pen Mycapssa® (octreotide) capsule QTY LIMIT: 4 caps/day Sandostatin® (octreotide) solution for injection Somatuline® Depot Injection (lanreotide) QTY LIMIT: 60 mg syringe = 0.2 ml/28 days, 90 mg syringe = 0.3 ml/28 days, 120 mg = 0.5 ml/28 days</p>	<p>Bynfezia, Sandostatin: the patient has a documented intolerance to Octreotide injection. Mycapssa: the diagnosis or indication is long-term maintenance treatment of acromegaly AND the patient has already responded to and tolerated treatment with an injectable somatostatin analog AND there is a clinically valid reason why the patient is unable to use Sandostatin LAR Depot. Somatuline: the patient has a documented side effect, allergy, treatment failure, or contraindication to Sandostatin LAR Depot.</p>
SPINAL MUSCULAR ATROPHY		
<p><u>Preferred After Clinical Criteria Are Met</u> ZOLGENSMA® (onasemnogene abeparvovec-xioi) intravenous suspension</p>	<p>Evrysdi® (risdiplam) oral solution Spinraza (nusinersen) injection 12mg/5ml single-dose vial</p>	<p>Evrysdi:</p> <ul style="list-style-type: none"> • The diagnosis is spinal muscular atrophy (SMA) AND • Patient is 2 months of age or older AND • Medication is prescribed per the dosing guidelines in the package insert AND • A negative pregnancy test is obtained for females of reproductive potential prior to initiating therapy and patient has been advised to use effective contraception during treatment and for at least 1 month after her last dose AND <p>A patient who has been started on Spinraza will not be approved for Evrysdi until at least 3 months after the fifth dose (i.e. nine months after the first loading dose, three months after the fifth dose). Concurrent use will not be approved.</p> <p>Note: For therapy continuation, clinical documentation must be submitted documenting improvement or maintenance of motor ability OR slower disease progression than would otherwise be expected.</p> <p>Spinraza:</p> <ul style="list-style-type: none"> • The diagnosis is spinal muscular atrophy (SMA) type 1,2, or 3 (results of genetic testing must be submitted) AND • The patient has at least 2 copies of the SMN2 gene AND • The need for invasive or noninvasive ventilation (if applicable) does not exceed more than 16 hours per 24 hour period AND • Baseline motor ability has been established using one of the following exams: <ul style="list-style-type: none"> ○ Hammersmith Infant Neurological Exam (HINE) ○ Hammersmith Functional Motor Scale Expanded (HFMSSE) ○ Upper Limb Module Test (non-ambulatory) ○ Children’s Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND) AND • Prior to starting therapy, and prior to each dose, the following laboratory tests will be conducted: Platelet count, prothrombin time (PT), activated partial thromboplastin time (aPTT), and quantitative spot urine protein • Concurrent use with Evrysdi will not be approved.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>Note: Initial approval will be granted for 4 loading doses (the first 3 loading doses should be administered at 14-day intervals; the 4th loading dose should be administered 30 days after the 3rd dose). Renewal may be granted for up to 12 months with a maximum of 3 doses approved per year (12mg(5ml) every 4 months). For therapy continuation, clinical documentation must be submitted documenting improvement or maintenance of motor ability OR slower progression of disease than would otherwise be expected.</p> <p>Zolgensma:</p> <ul style="list-style-type: none"> • The patient is less than 2 years of age AND • The diagnosis is spinal muscular atrophy (SMA) AND • The patient has bi-allelic mutations of the SMN1 gene AND • The patient does not have advanced SMA (e.g. complete paralysis of limbs or permanent ventilator dependence) AND • Medication is prescribed per the dosing guidelines in the package insert (recommended dose is 1.1 x 10⁴ vector genomes per kilogram) AND • Baseline anti-AAV9 antibodies are less than 1:50 AND • Prior to starting therapy and periodically for at least 3 months, the following laboratory tests will be conducted: Liver function (AST, ALT, total bilirubin, prothrombin time), platelet counts, and troponin-I <p>Note: The safety and effectiveness of repeat administration has not been evaluated. Approval is limited to a single intravenous infusion.</p> <ul style="list-style-type: none"> •

SYSTEMIC LUPUS ERYTHEMATOSUS (SLE)

	<p>Benlysta® (belimumab) Maximum days supply per fill = 28 days</p> <p>Saphnelo™ (anifrolumab-fnia)</p>	<p>Benlysta: <i>Indication for use is Systemic Lupus Erythematosus (SLE):</i></p> <ul style="list-style-type: none"> • The patient is positive for autoantibodies (anti-nuclear antibody (ANA) and/or anti-double-stranded DNA (anti-dsDNA) AND • The patient has had a documented inadequate response or intolerance to at least TWO of the following agents: NSAIDs, hydroxychloroquine, corticosteroids, azathioprine, methotrexate, mycophenolate mofetil AND • Initial approval will be granted for 3 months. For therapy continuation, clinical documentation must be submitted documenting stable disease activity OR reduction in disease activity or corticosteroid dose. Note: The efficacy of Benlysta® has not been evaluated in patients with severe active central nervous system lupus. Benlysta has not been studied in combination with other biologics or intravenous cyclophosphamide. Use of Benlysta is not recommended in these situations. <p><i>Indication for use is Active Lupus Nephritis:</i></p> <ul style="list-style-type: none"> • Diagnosis has been confirmed by urine/blood tests or kidney biopsy AND • The patient is ≥ 18 years of age AND • Medication is prescribed by, or in consultation with, a nephrologist or rheumatologist AND • The patient has clinical progression (e.g. worsening of proteinuria or serum creatinine) after 3 months of induction therapy with corticosteroids plus cyclophosphamide or mycophenolate mofetil OR failure to respond after 6
--	---	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>months of induction therapy with corticosteroids plus cyclophosphamide or mycophenolate mofetil AND</p> <ul style="list-style-type: none"> Medication will be used in combination with background immunosuppressive therapy (e.g. mycophenolate mofetil and systemic corticosteroids) AND Initial approval will be granted for 3 months. For therapy continuation, clinical documentation must be submitted documenting stable disease activity OR reduction in disease activity. <p>Saphnelo:</p> <ul style="list-style-type: none"> The patient has a diagnosis of moderate-severe Systemic Lupus Erythematosus AND The patient is ≥ 18 years of age AND Medication is prescribed by, or in consultation with, a nephrologist or rheumatologist AND The patient has had a documented inadequate response or intolerance to at least TWO of the following agents: hydroxychloroquine, corticosteroids, azathioprine, methotrexate, mycophenolate mofetil AND The patient has had a documented intolerance or treatment failure with Benlysta <ul style="list-style-type: none"> Initial approval will be granted for 3 months. For therapy continuation, clinical documentation must be submitted documenting stable disease activity OR reduction in disease activity or corticosteroid dose. Note: The efficacy of Saphnelo has not been evaluated in patients with severe active lupus nephritis or severe active central nervous system lupus. Saphnelo has not been studied in combination with other biologics or intravenous cyclophosphamide. Use of Saphnelo is not recommended in these situations.

MOOD STABILIZERS

LITHIUM CARBONATE (formerly Eskalith®) LITHIUM CARBONATE SR (compare to Lithobid®, formerly Eskalith CR®) LITHIUM CITRATE SYRUP	Equetro® (carbamazepine SR) Lithobid® (lithium carbonate SR)	<p>Lithobid: The patient has had a documented side effect, allergy, or treatment failure with the generic equivalent of the requested medication.</p> <p>Equetro: The patient has had a documented side effect, allergy, or treatment failure with a carbamazepine product from the anticonvulsant therapeutic drug category</p>
--	---	--

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

MOVEMENT DISORDERS

TETRABENAZINE

QTY LIMIT: 50 mg/day at initial approval (12.5 mg tablets ONLY), up to 100 mg/day at subsequent approvals (12.5 mg pr 25 mg tablets)

Preferred After Clinical Criteria Are Met

AUSTEDO® (deutetrabenazine) tablets

QTY LIMIT: 48 mg/day

Maximum 1-month supply per fill

AUSTEDO XR® (deutetrabenazine) extended-release tablets

QTY LIMIT: 6 mg and 12 mg = 1 tablet/day; 24 mg = 2 tablets/day; Starter pack = 42 tablets/28 days

Maximum 1-month supply per fill

INGREZZA® (valbenazine tosylate) capsules

QTY LIMIT: 80 mg/day

Maximum 1-month supply per fill

Xenazine® (tetrabenazine) tablets

QTY LIMIT: 50 mg/day at initial approval (12.5 mg tablets ONLY), up to 100 mg/day at subsequent approvals (12.5 mg or 25 mg tablets)

Maximum 1-month supply per fill

Austedo, Austedo XR, Ingrezza: The diagnosis or indication for the requested medication is Huntington's Disease (HD) with chorea or Tardive Dyskinesia (TD) AND the results of an Abnormal Involuntary Movement Scale (AIMS) exam have been submitted AND the patient is ≥ 18 years of age. **If the request is for Huntington's Disease (HD) with chorea, patient has documented side effect, allergy, or treatment failure with Tetrabenazine.** For re-approval, there must be documented clinical improvement.

Xenazine: The diagnosis or indication for use is Tourette Syndrome OR the diagnosis or indication for use is Huntington's Disease (HD) with Chorea or Tardive Dyskinesia (TD) AND the patient is ≥ 18 years of age AND for approval of Xenazine, the patient must have a documented intolerance to tetrabenazine.

Note: Austedo, Tetrabenazine, and Xenazine are contraindicated in patients with Huntington's Disease who are suicidal or with untreated/inadequately treated depression.

MULTIPLE SCLEROSIS MEDICATIONS

INJECTABLES

INTERFERONS

AVONEX® (interferon B-1a)

BETASERON® (interferon B-1b)

REBIF® (interferon B-1a)

REBIF® REBIDOSE (interferon B-1a)

OTHER

COPAXONE® 20 mg (glatiramer acetate)

QTY LIMIT: 1 kit/30 days

Preferred After Clinical Criteria are Met

Briumvi™ (ublituximab-xiiy)

Extavia® (interferon beta-1b)

Copaxone® 40 mg (glatiramer)

QTY LIMIT: 12 syringes (12 ml)/28 days

Glatiramer Acetate (compare to Copaxone®) 20 mg

QTY LIMIT: 1 kit/30days

Glatiramer Acetate (compare to Copaxone®) 40 mg

QTY LIMIT: 12 syringes (12 ml)/28 days

Glatopa® 20 mg (glatiramer acetate)

QTY LIMIT: 1 carton (30 syringes/30 days)

Ampyra, Aubagio, Gilenya, Tecfidera: patient must have a documented intolerance to the generic equivalent.

Bafiertam, Vumerity: Patient is ≥ 18 years AND has a diagnosis of relapsing forms of Multiple Sclerosis AND the patient has a documented side effect, allergy, treatment failure, or contraindication to at least two preferred drugs, one of which must be Dimethyl fumarate.

Copaxone 40 mg Syringe: The patient is unable to tolerate or be compliant with Copaxone 20 mg daily dosing.

Extavia: Patient has a diagnosis of multiple sclerosis. AND The provider provides a clinical reason why Betaseron cannot be prescribed.

Glatiramer, Glatopa: Patient is ≥ 18 years AND diagnosis of relapsing forms of Multiple Sclerosis AND the provider provides a clinical reason why

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>KESIMPTA® (ofatumumab) TYSABRI® (natalizumab)</p> <p>ORAL DALFAMPRIDINE ER tablet (compare to Ampyra®) QTY LIMIT: 2 tablets/day Maximum 30-day supply per fill DIMETHYL FUMARATE QTY LIMIT: 2 capsules/day Maximum 30-day supply per fill FINGOLIMOD capsule (compare to Gilenya®) QTY LIMIT: 1 capsule/day Maximum 30-day supply per fill TERIFLUNOMIDE (compare to Aubagio®) tablet QTY LIMIT: 1 tablet/day Maximum 30-day supply per fill</p>	<p>Glatopa® 40 mg (glatiramer) QTY LIMIT: 12 syringes (12 ml)/28 days Lemtrada® (alemtuzumab) intravenous Ocrevus® (ocrelizumab) QTY LIMIT: 300 mg X 2 doses, then 600 mg every 6 months thereafter Plegridy® (peginterferon beta-1a)</p> <p>Ampyra® (dalfampridine ER) tablet QTY LIMIT: 2 tablets/day Maximum 30-day supply per fill Aubagio® (teriflunamide) tablet QTY LIMIT: 1 tablet/day Maximum 30-day supply per fill Bafiertam® (monomethyl fumarate) capsule QTY LIMIT: 4 capsules/day Maximum 30-day supply per fill Gilenya® (fingolimod) capsule QTY LIMIT: 1 capsule/day Maximum 30-day supply per fill Mavenclad® (cladribine) tablet Mayzent® (siponimod) tablet Ponvory™ (ponesimod) tablet QTY LIMIT: 1 tablet/day Maximum 30-day supply per fill Tascenso ODT® (fingolimod) QTY LIMIT: 1 capsule/day Maximum 30-day supply per fill Tecfidera® (dimethyl fumarate) QTY LIMIT: 2 capsules/day Maximum 30-day supply per fill Vumerity® (diroximel fumarate) capsule QTY LIMIT: 4 capsules/day Zeposia® (ozanimod) capsule QTY LIMIT: 1 capsule/day</p>	<p>Copaxone cannot be prescribed. For Glatopa: Clinical reason why Glatiramer cannot be used</p> <p>Kesimpta: Patient is ≥18 years AND has a diagnosis of relapsing multiple sclerosis AND has a documented side effect, allergy, or treatment failure to one preferred drug.</p> <p>Mavenclad: Patient is ≥ 18 years AND has a diagnosis of relapsing-remitting MS (RRMS) or active secondary progressive MS (SPMS) AND Documentation is provided showing ≥ 1 relapse within the past year AND baseline CBC w/ diff (including lymphocyte count), liver function tests, and MRI (within the past 3 months) have been completed AND the patient is negative for HIV, Hepatitis B, and Hepatitis C infections AND the patient is not pregnant AND patient has a documented side effect, allergy, treatment failure or contraindication to at least three preferred drugs AND dosing does not exceed any of the following: 2 tablets per day, 10 tablets per cycle, 2 treatment cycles per course, 1 course per year. Following the administration of 2 treatment courses, Mavenclad may not be administered during the next 2 years.</p> <p>Mayzent, Ponvory, Zeposia: <i>Diagnosis of relapsing-remitting MS, Clinical Isolated Syndrome, or Active Secondary Progressive MS (SPMS):</i></p> <ul style="list-style-type: none"> • Patient is ≥ 18 years AND • Patient CYP2C9 variant status has been tested to determine genotyping (Mayzent only; required for dosing; therapy is contraindicated in CYP2C9*3/*3) AND • Baseline CBC, electrocardiogram (ECG), and ophthalmic evaluation have been completed AND • Patient has a documented side effect, allergy, treatment failure or contraindication to at least two preferred drugs, one of which must be Fingolimod. <p>Briumvi, Lemtrada, Ocrevus: Patient is ≥18 years AND has a diagnosis of relapsing multiple sclerosis AND has a documented side effect, allergy, treatment failure or contraindication to at least two preferred drugs, one of which must be Tysabri, unless contraindicated. OR Patient is ≥18 years AND has a diagnosis of primary progressive multiple sclerosis (Ocrevus only).</p> <p>Plegridy: Patient is ≥ 18 years AND has a diagnosis of relapsing form of Multiple Sclerosis AND has a documented side effect, allergy, treatment failure or contraindication to at least three preferred drugs including at least one preferred form of interferon.</p> <p>Tascenso ODT: patient has a medical necessity for a non-solid oral dosage form.</p> <p>Tysabri: Patient is ≥ 18 years AND has a diagnosis of relapsing multiple sclerosis (including clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease).</p>

MUSCLE RELAXANTS, SKELETAL

MUSCULOSKELETAL AGENTS

SINGLE AGENTS

CYCLOBENZAPRINE 5 mg, 10mg tablets (compare to

Flexeril[®])

QTY LIMIT: 5 mg = 6 tablets/day, 10 mg = 3 tablets/day

METHOCARBAMOL tablets (compare to Robaxin[®])

QTY LIMIT: 8 tablets/day

ORPHENADRINE CITRATE ER 100 mg tablet

QTY LIMIT: 2 tablets/day

COMBINATION PRODUCT

All products require PA

ASA = aspirin

ANTISPASTICITY AGENTS

BACLOFEN tablets

DANTROLENE (compare to Dantrium[®])

TIZANIDINE (compare to Zanaflex[®]) tablets

Amrix[®] (cyclobenzaprine sustained-release) capsule

QTY LIMIT: 1 capsule/day

Carisoprodol tablets

QTY LIMIT: 8 tablets/day

Chlorzoxazone tablets

QTY LIMIT: 4 tablets/day

Cyclobenzaprine 7.5 mg tab (compare to Fexmid[®])

QTY LIMIT: 3 tablets/day

Cyclobenzaprine ER (compare to Amrix[®])

QTY LIMIT: 1 capsule/day

Fexmid[®] (cyclobenzaprine) 7.5 mg tablet

QTY LIMIT: 3 tablets/day

Lorzone[®] (chlorzoxazone) tablets

QTY LIMIT: 4 tablets/day

Metaxalone tablets (compare to Skelaxin[®])

QTY LIMIT: 4 tablets/day

Soma[®] (carisoprodol) tablets

QTY LIMIT: 4 tablets/day

Baclofen oral solution

Dantrium[®] (dantrolene)

Fleqsuvy[™] (baclofen) oral suspension

Lyvispah[™] (baclofen) oral granule packet

Tizanidine (compare to Zanaflex[®]) capsules

Zanaflex[®] (tizanidine) capsules

Zanaflex[®] (tizanidine) tablets

Amrix, Cyclobenzaprine 7.5 mg, Cyclobenzaprine ER, Fexmid: The prescriber must provide a clinically valid reason why a preferred generic cyclobenzaprine 5mg or 10mg cannot be used. For approval of Fexmid, the patient must also have a documented intolerance to the generic equivalent.

Baclofen oral solution Fleqsuvy: Patient has a medical necessity for a non-solid oral dosage form AND the patient has a documented intolerance to Lyvispah.

Carisoprodol, Chlorzoxazone, Lorzone, Soma, Metaxalone: The patient has had a documented side effect, allergy or treatment failure with two different preferred musculoskeletal agents. Additionally, if a brand name product is requested where an AB rated generic exists, the patient must also have had a documented intolerance to the generic product.

Dantrium, Zanaflex Tablets: The patient must have a documented intolerance with the AB rated generic product.

Lyvispah: Patient has a medical necessity for the non-solid oral dosage form.

Tizanidine capsules, Zanaflex capsules: The prescriber must provide a clinically valid reason why generic tizanidine tablets cannot be used. AND If the request is for Zanaflex capsules, the patient must have a documented intolerance to generic tizanidine capsules.

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

MUSCULAR DYSTROPHY AGENTS

Preferred After Clinical Criteria Are Met

Agamree® (vamorolone) suspension
Amondys®45 (casimersen)

Emflaza, Agamree, deflazacort:

EMFLAZA® (deflazacort)

Deflazacort
Elevidys® (delandistrogene
moxeparvovec-rokl)
Exondys 51™ (eteplirsen)
Viltepso® (viltolarsen)
Vyondys 53™ (golodirsen)

- The patient must be ≥ 2 years of age AND
- The patient must have a diagnosis of Duchenne Muscular Dystrophy AND
- There is documented improvement in muscle function or strength with use of prednisone, but the patient has experienced weight gain >10% of body weight within 3 months or >25% within 1 year.
- For Agamree or deflazacort, the patient must have tried and failed or been intolerant to preferred therapies (Emflaza)

Amondys, Exondys, Viltepso, Vyondys:

- The patient must have a diagnosis of Duchenne Muscular Dystrophy with a confirmed mutation of the DMD gene that is amenable to exon 45 skipping (for Amondys) or exon 51 skipping (for Exondys) or exon 53 skipping (for Viltepso, Vyondys) (results of genetic testing must be submitted) AND
- The prescriber is, or has consulted with, a neuromuscular disorder specialist AND
- The dose does not exceed 30mg/kg once weekly (for Amondys, Exondys, Vyondys) or 80mg/kg once weekly (for Viltepso) AND
- The patient is currently on a stable corticosteroid dose for at least 6 months. AND
- Baseline documentation of the members voluntary motor and cardiac function has been provided and results have shown member retains meaningful voluntary motor function:

Optional

- 6-minute walk test (6MWT) or other timed functions tests (e.g time to stand [TTSTAND], time to run/walk 10 meters [TTRW], time to climb 4 stairs [TTCLIMB])
- Brooks Upper Extremity Test
- North Star Ambulatory Assessment (NSAA)
Required
- Forced Vital Capacity (FVC) percent predicted
- Ejection Fraction Percentage

Elevidys:

- The patient is 4 to 5 years of age AND
- The patient is ambulatory, and the results of the North Star Ambulatory Assessment (NSAA) have been provided AND

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<ul style="list-style-type: none"> • The patient must have a diagnosis of Duchenne Muscular Dystrophy with either a confirmed frameshift mutation, or a premature stop codon mutation between exons 18 to 58 in the DMD gene AND • The patient does not have a deletion in exon 8 or 9 of the DMD gene. AND The baseline anti-AAVrh74 antibody titer results are <1:400, using a Total Binding Antibody enzyme-linked immunosorbent assay (ELISA). AND • The client is not on concomitant DMD antisense oligonucleotide therapy (e.g. golodirsen, casimersen, viltolarsen, eteplirsen) AND • Prescriber attests to complete the following: <ul style="list-style-type: none"> • Assess liver function (clinical exam, GGT, and total bilirubin) at baseline and weekly for the first 3 months. Continue monitoring if clinically indicated, until results are unremarkable (normal clinical exam, GGT, and total bilirubin levels return to near baseline levels). • Obtain platelet counts at baseline and weekly for the first two weeks. • Obtain troponin-I at baseline and weekly for the first month. • Approval will be granted for a maximum of one dose per lifetime and may not be renewed. <p>Note: Initial approval will be granted for 6 months. For re-approval after 6 months, the patient must demonstrate a response to therapy compared to baseline as evidenced by stable, improved, or slowed rate of either motor function or cardiac function degradation. Evidence may include one or more of the following (not all-inclusive):</p> <ul style="list-style-type: none"> • 6MWT or other timed function tests (e.g., time to stand [TTSTAND], time to run/walk 10 meters [TTRW], time to climb 4 stairs [TTCLIMB]) • Brooks Upper Extremity Test • North Star Ambulatory Assessment (NSAA) • Forced Vital Capacity (FVC) percent predicted • Ejection Fraction Percentage • Improvement in quality of life.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

NEUROGENIC ORTHOSTATIC HYPOTENSION

<p>FLUDROCORTISONE MIDODRINE</p>	<p>Northera®</p>	<p>Quantity Limits:</p> <ul style="list-style-type: none"> Initial 2 weeks approval Continued therapy approvals based on documentation of continued benefit clinically and as evidenced by positional blood pressure readings <p>Clinical Criteria:</p> <ul style="list-style-type: none"> diagnosis of neurogenic orthostatic hypotension caused by primary autonomic failure (Parkinson’s disease, multiple system atrophy, or pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy, AND the presentation of symptoms including dizziness, lightheadedness, and the feeling of “blacking out” AND Failure of multiple non-pharmacologic measures as appropriate (e.g. removal of offending medications, compression stockings, increased fluid and salt intake) AND Failure, intolerance or contra-indication to fludrocortisone AND midodrine
--------------------------------------	------------------	---

NEUROPATHIC PAIN & FIBROMYALGIA AGENTS

Oral		
<p>DULOXETINE (compare to Cymbalta®) <i>QTY LIMIT:</i> 2 capsules/day</p> <p>LYRICA® (pregabalin) capsules <i>QTY LIMIT:</i> 3 capsules/day</p> <p>LYRICA® (pregabalin) solution</p> <p>PREGABALIN (compare to Lyrica®) capsules <i>QTY LIMIT:</i> 3 capsules/day</p> <p>SAVELLA® (milnacipran) tablet, titration pack <i>QTY LIMIT:</i> 2 tablets/day</p>	<p>Cymbalta® (duloxetine) <i>QTY LIMIT:</i> 2 capsules/day</p> <p>Gralise® (gabapentin) tablet, starter pack <i>QTY LIMIT:</i> 3 tablets/day Maximum 30-day supply per fill</p> <p>Horizant® (gabapentin enacarbil) ER Tablet FDA maximum recommended dose = 1200 mg/day</p> <p>Lyrica® CR (pregabalin, extended release) FDA maximum recommended dose = 330 mg/day (DPN), 660 MG/day (PHN)</p> <p>Pregabalin (compare to Lyrica®) solution</p> <p>Pregabalin extended release (compare to Lyrica® CR) FDA maximum recommended dose = 330 mg/day (DPN), 660 mg/day (PHN)</p>	<p>Cymbalta: The patient has had a documented intolerance with the generic equivalent.</p> <p>Gralise, Horizant: The patient has a diagnosis of post-herpetic neuralgia (PHN) AND The patient has had a documented side effect, allergy, contraindication or treatment failure with at least one drug from the tricyclic antidepressant class AND The patient has had an inadequate response to the generic gabapentin immediate-release.</p> <p>Pregabalin ER, Lyrica CR: The patient has a diagnosis of post-herpetic neuralgia (PHN) or diabetic peripheral neuropathy (DPN) AND patient has not been able to be adherent to a twice daily dosing schedule of pregabalin immediate release resulting in a significant clinical impact AND for approval of pregabalin ER, the patient has a documented intolerance to brand Lyrica CR. Note: The efficacy of Lyrica® CR has not been established for the management of fibromyalgia or as adjunctive therapy for adult patients with partial onset seizures.</p> <p>Pregabalin solution: The patient is unable to use pregabalin capsules (i.e. swallowing disorder) AND has a documented intolerance to brand Lyrica solution.</p>

NUTRITIONALS, LIQUID ORAL SUPPLEMENTS

All products require PA

Note: Nutritional supplements administered via tube feeds may be provided through the Medical Benefit

EleCare, EleCare Jr: The patient is an infant or child who needs an amino acid-based medical food or who cannot tolerate intact or hydrolyzed protein. AND The product is being requested for the dietary management of protein maldigestion, malabsorption, severe food allergies, short-bowel syndrome, eosinophilic GI disorders, GI-tract impairment, or other conditions for which an amino acid-based diet is required.

All Others: Requested nutritional supplement will be administered via tube feeding. OR Patient has one of the following conditions where feeding is difficult or malabsorption or maldigestion occurs: AIDS, Cancer, Cerebral Palsy, Cystic Fibrosis, Dementia resulting in loss of motor skills, Neuromuscular Disease, Short Gut. OR Patient has experienced unplanned weight loss or is extremely low weight (see further definitions below) OR Patient has demonstrated nutritional deficiency identified by low serum protein levels (albumin or pre-albumin levels to be provided) (albumin <3.5 g/dL /pre-albumin <15 mg/dL)

Unplanned Weight Loss/Low Weight Table:

Adult: Involuntary loss of > 10 % of body weight within 6 months OR Involuntary loss of > 5% of body weight within 1 month OR Loss of > 2% of body weight within one week OR BMI of < 18.5 kg/m²

Elderly: (>65): Involuntary loss of > 10 % of body weight within 6 months OR Involuntary loss of > 5 % of body weight within 3 months OR Loss of > 2 % of body weight within one month OR BMI of < 18.5 kg/m²

Children: Anatomic causes for malnutrition have been evaluated and treated AND clinical diagnosis and documentation supports the need for enteral nutrition (See Below)

- Members weight is below the 5th percentile for sex and corrected age AND weight-to-length ratio is below the 10th percentile OR
- Sustained decrease in growth velocity as demonstrated by weight-for-age or weight-for-length fall by two major percentiles (percentile markers 95, 90, 75, 50, 25, 10, and 5) over time (defined by the WHO for children less than 2 years of age and the CDC for children greater than 2 years of age)

Limitations: Approvals will be based on medical necessity for supplemental nutrition. Approval will NOT be granted for individuals whose need is nutritional rather than medical, including an unwillingness to consume solid or pureed foods. For nonmedical needs contact WIC at 800-464-4343

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

ONCOLOGY: DRUGS (select)

Clinical Criteria: Medication is being used for an FDA approved indication AND age, dose, duration, required concurrent therapy, and past treatment failures (if applicable) are consistent with prescribing information AND the patient does not have any contraindications prohibiting use of the medication OR medication is being used in accordance with the National Comprehensive Cancer Network® (NCCN®) Clinical Practice Guidelines. Requests outside of these parameters require medical director review. This includes all cell and gene therapies, including CAR-T therapies, regardless of site of administration. For physician-administered drugs, please refer to the Fee Schedule for which codes require a PA: <http://vtmedicaid.com/#/feeSchedule/hcps>

OPHTHALMICS

ANTIBIOTICS

QUINOLONES

BESIVANCE® (besifloxacin) suspension
 CILOXAN® ointment
 CIPROFLOXACIN HCL (compare to Ciloxan®) solution
 MOXIFLOXACIN 0.5% solution (compare to Vigamox®)
 OFLOXACIN (compare to Ocuflax®) solution

MACROLIDES

AZASITE® (azithromycin) solution
 ERYTHROMYCIN ointment

Gatifloxacin 0.5% solution (compare to Zymaxid®)
 Levofloxacin 0.5 % solution
 Moxifloxacin 0.5% (compare to Moxeza®) (preservative free) solution
 Ocuflax® (ofloxacin) solution
 Vigamox® (moxifloxacin 0.5%) (preservative free) solution
 Zymaxid® (gatifloxacin 0.5%) solution

All other brands

Criteria for All Non-Preferred: The patient has had a documented side effect, allergy, or treatment failure with at least TWO preferred ophthalmic antibiotics or ophthalmic antibiotic combination agents, one of which must be in the same therapeutic class. (If a product has an AB rated generic, there must have also been a trial of the generic formulation.)

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>AMINOGLYCOSIDES</u></p> <p><u>SINGLE AGENT</u> GENTAMICIN solution TOBRAMYCIN solution (compare to Tobrex®)</p> <p><u>COMBINATION</u> TOBRAMYCIN W/DEXAMETHASONE suspension ZYLET® (tobramycin/loteprednol) suspension</p> <p><u>MISCELLANEOUS</u> <u>SINGLE AGENT</u> All products require PA</p> <p><u>Combination</u> BACITRACIN ZINC W/POLYMYXIN B ointment NEOMYCIN/BACITRACIN/POLYMYXIN ointment NEOMYCIN/POLYMYXIN W/DEXAMETHASONE (compare to Maxitrol®) ointment, suspension NEOMYCIN/POLYMYXIN/BACITRACIN/HYDROCORTISONE ointment POLYMYXIN B W/TRIMETHOPRIM (compare to Polytrim®) solution SULFACETAMIDE W/PREDNISOLONE SOD PHOSPHATE solution</p>	<p>Tobrex® ointment (tobramycin)</p> <p>Tobradex ST® (tobramycin/dexamethasone) suspension</p> <p>Bacitracin ointment Sulfacetamide sodium (compare to Bleph-10®) solution Sulfacetamide sodium ointment</p> <p>Maxitrol® (neomycin/polymyxin/dexamethasone) suspension, ointment Neomycin/Polymyxin W/Gramicidin solution Neomycin/Polymyxin w/Hydrocortisone ointment, suspension Polytrim® (polymyxin B/trimethoprim) soln</p>	
ANTI-HISTAMINES		
<p>AZELASTINE KETOTIFEN 0.025 % OLOPATADINE 0.1%, 0.2%</p>	<p>Bepotastine (compare to Bepreve®) Bepreve® (bepotastine besilate) Epinastine Zerviate® (cetirizine 0.24%) <i>QTY LIMIT: 60 vials/30 days</i></p>	<p>Bepotastine, Bepreve, Epinastine: The patient has had a documented side effect, allergy, or treatment failure to a preferred ophthalmic antihistamine AND for approval of Bepotastine, the patient must have a documented intolerance to brand Bepreve.</p> <p>Zerviate: The patient has had a documented side effect, allergy, or treatment failure to TWO preferred ophthalmic antihistamines.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
ANTIPARASITICS		
	<p>Xdemvy® (lotilaner ophthalmic solution) QTY LIMIT: 20 ml per approval (6 weeks of treatment)</p>	<p>Xdemvy: Patient is ≥ 18 years of age AND has a diagnosis of Demodex blepharitis AND has the presence of the following in at least one eye: more than 10 lashes with collarettes present on the upper lid (collarette scale grade 2 or worse) AND at least mild erythema in at least one eye of the upper eyelid margin</p>
ANTI-VEGF AND MISCELLANEOUS AGENTS		
<p>EYLEA® (aflibercept) EYLEA® HD (aflibercept) LUCENTIS® (ranibizumab)</p>	<p>Beovu® (brolocizumab-dbll) Byooviz™ (ranibizumab-nuna) biosimilar to Lucentis® Cimerli® (ranibizumab-eqrn) biosimilar to Lucentis® Susvimo® (ranibizumab) implant Syfovre® (pegcetacoplan) QTY LIMIT: 15mg (0.1mL) per dose (each affected eye) every 25 days Vabysmo® (faricimab-svoa)</p>	<p>Beovu, Vabysmo: The patient has a documented side effect, allergy, or treatment failure with Eylea and Lucentis. Byooviz, Cimerli: Patient must be unable to use Lucentis. Susvimo: Patient has had a positive clinical response to an intravitreal formulation of ranibizumab AND Medical necessity for a specialty dosage form has been provided. Syfovre: Medication is being used for the treatment of Geographic Atrophy (GA) secondary to age-related macular degeneration (AMD) and the patient is not considered legally blind (visual acuity score of 20/200 or worse). Initial approval will be granted for 6 months. For re-approval, documentation is required showing patient has not progressed to or beyond a visual acuity score of 20/200.</p>
CORTICOSTEROIDS: TOPICAL		
<p>ALREX® (loteprednol) 0.2% suspension DEXAMETHASONE sodium phosphate 0.1% solution DUREZOL® (difluprednate) 0.05% emulsion FML Forte® (fluorometholone) 0.25% suspension FLUROMETHOLONE 0.1% suspension FML® (fluorometholone) 0.1% ointment LOTEMAX® (loteprednol) 0.5% suspension, Ointment, gel MAXIDEX® (dexamethasone) suspension PRED MILD® (prednisolone acetate) 0.12% suspension PREDNISOLONE ACETATE 1% suspension PREDNISOLONE SODIUM PHOSPHATE 1% solution</p>	<p>Difluprednate (compare to Durezol®) Flarex® (fluorometholone acetate) 0.1% suspension FML Liquifilm® (fluorometholone) 0.1% suspension Inveltys™ (loteprednol) suspension Lotemax SM (loteprednol) 0.038% gel drops Loteprednol suspension Pred Forte® (prednisolone acetate) 1% suspension All other brands</p>	<p>Non-preferred agents: The patient has had a documented side effect, allergy, or treatment failure with TWO preferred ophthalmic corticosteroids. (If a product has an AB rated generic, there must have been a trial of the generic formulation)</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
CYSTEAMINE		
All products require PA	Cystadrops® (cysteamine) 0.37% ophthalmic solution QTY LIMIT: 4 bottles (20 ml)/28 days Maximum day supply/Rx = 28 days Cystaran® (cysteamine) 0.44% ophthalmic solution QTY LIMIT: 4 bottles (60 ml)/ 28 days Maximum day supply/RX = 28 days	Cystadrops, Cystaran: The indication for use is corneal cystine accumulation in patients with cystinosis.
DRY EYE SYNDROME		
<p>OCULAR LUBRICANTS Please refer to the DVHA website for covered OTC ocular lubricants https://dvha.vermont.gov/sites/dvha/files/documents/OTCWebList_0.pdf</p> <p>IMMUNOMODULATORS EYSUVIS® (loteprednol etabonate ophthalmic suspension) 0.25% RESTASIS® (cyclosporine ophthalmic emulsion) 0.05% droperette (NDC 00023916330 and 00023916360 are the only preferred NDC's) QTY LIMIT: 180 vials per 90 days XIIDRA® (lifitegrast) solution QTY LIMIT: 180 vials per 90 days</p>	Cequa™ (cyclosporine ophthalmic solution) 0.09% Cyclosporin ophthalmic emulsion 0.05% droperette (compare to Restasis®) QTY LIMIT: 180 vials per 90 days Miebo® (perfluorohexyloctane ophthalmic solution) Restasis® (cyclosporine ophthalmic emulsion) 0.05% multidose bottle QTY LIMIT: 1 bottle (5.5ml) per 25 days Tyrvaya™ (varenicline) nasal spray QTY LIMIT: 2 bottles (8.4 ml) per 30 days Verkazia® (cyclosporine ophthalmic emulsion) 0.1% single dose vials Vevye® (cyclosporine ophthalmic solution) 0.1%	<p>Cequa, Vevye: The patient has a diagnosis of Dry Eye Disease AND has a documented side effect, allergy, or treatment failure to two ophthalmic immunomodulators, one of which must be Restasis. For Approval of Vevye: the patient must have had a treatment failure with Cequa.</p> <p>Cyclosporin emulsion, Tyrvaya: The patient has a diagnosis of Dry Eye Disease AND has a documented side effect, allergy or treatment failure to Restasis.</p> <p>Miebo: The patient has a diagnosis of Dry Eye Disease AND is ≥ 18 years of age AND has a documented side effect, allergy, treatment failure or contraindication to Restasis and Xiidra AND Miebo will not be used in combination with cyclosporine, lifitegrast or varenicline.</p> <p>Restasis Multidose: Both package sizes of the droperettes must be on a long-term backorder and unavailable from the manufacturer.</p> <p>Verkazia: The patient has a diagnosis of vernal keratoconjunctivitis (VKC) AND the patient has had a documented side effect, allergy, or treatment failure with a mast cell stabilizer (e.g. cromolyn sodium) or a dual acting antihistamine/mast cell stabilizer (e.g. olopatadine, azelastine)</p>
GLAUCOMA AGENTS/MIOTICS		

PREFERRED AGENTS
(No PA required unless otherwise noted)

ALPHA-2 ADRENERGIC SINGLE AGENT

ALPHAGAN P[®] 0.1 %, 0.15 % (brimonidine tartrate)
BRIMONIDINE TARTRATE 0.2 %

COMBINATION

COMBIGAN[®] (brimonidine tartrate/timolol maleate)
SIMBRINZA[®] (brinzolamide 1% and brimonidine 0.2%) Suspension

BETA BLOCKER

CARTEOLOL HCL
LEVOBUNOLOL HCL
TIMOLOL MALEATE

PROSTAGLANDIN INHIBITORS

LATANOPROST (compare to Xalatan[®])
LUMIGAN[®] (bimatoprost)
TRAVATAN Z[®] (travoprost) (BAK free)
ZIOPTAN[®] (tafluprost)

RHO KINASE INHIBITORS

SINGLE AGENT

RHOPRESSA[®] (netarsudil)

COMBINATION

ROCKLATAN[®] (netarsudil/latanoprost)

CARBONIC ANHYDRASE INHIBITOR

SINGLE AGENT

AZOPT[®] (brinzolamide 1%)
DORZOLAMIDE 2 % (compare to Trusopt[®])

NON-PREFERRED AGENTS
(PA required)

Apraclonidine (compare to Iopidine[®])
Brimonidine tartrate 0.1%, 0.15 % (compare to Alphagan P[®])
Iopidine[®] (apraclonidine)

Brimonidine tartrate/timolol maleate (compare to Combigan[®])

Betaxolol HCl solution
Betoptic S[®] (betaxolol suspension)
Istalol[®] (timolol)
Timolol maleate PF (compare to Timoptic[®] Ocudose) droperette
Timoptic[®] Ocudose (timolol maleate) preservative free droperette
Timolol maleate gel (formerly Timotic XE[®])

Bimatoprost 0.03% (Lumigan[®])
Durysta[®] (bimatoprost) 10 mcg implant
Iyuzeh[™] (latanaprost)
Tafluprost PF solution (compare to Zioptan[®])
Travoprost BAK Free (compare to Travatan Z[®])
Vyzulta[®] (latanoprostene bunod)
Xelpros[®] (latanoprost) (BAK free)

Brinzolamide 1% (compare to Azopt[®])

PA CRITERIA

ALPHA 2 ADRENERGIC AGENTS: Single Agent: The patient has had a documented side effect, allergy, or treatment failure with at least one preferred ophthalmic alpha 2 adrenergic agent. If the request is for brimonidine tartrate 0.1% or 0.15%, the patient must have a documented intolerance of brand name Alphagan P.

Brimonidine/timolol: the patient must have a documented intolerance to brand Combigan.

BETA BLOCKERS: The patient has had a documented side effect, allergy, or treatment failure with at least one preferred ophthalmic beta blocker OR the patient has a documented intolerance to the preservatives in generic timolol maleate.

PROSTAGLANDIN INHIBITORS

Bimatoprost, Iyuzeh, Tafluprost, Travoprost, Vyzulta, Xalatan, Xelpros:

The patient has had a documented side effect, allergy, or treatment failure with at least 2 preferred prostaglandin inhibitors AND if a product has an AB rated preferred formulation, there must have also been a trial of the preferred formulation.

Durysta: The patient has had a documented side effect, allergy, or treatment failure with at least 2 preferred prostaglandin inhibitors OR the patient is not a candidate for topical drop therapy AND the patient does not have any of the following contraindications:

- History of prior corneal transplantation or endothelial cell transplants (e.g. Descemet's Stripping Automated Endothelial Keratoplasty)
- Diagnosis of corneal endothelial dystrophy (e.g. Fuchs' Dystrophy)
- Absent or ruptured posterior lens capsule

Approval will be limited to a single implant per eye without retreatment.

CARBONIC ANHYDRASE INHIBITORS

Brinzolamide: the patient has a documented intolerance to a preferred carbonic anhydrase inhibitor.

Cosopt PF, Dorzolamide w/timolol PF: The patient has had a documented intolerance to the preservatives in the generic combination product.

Miscellaneous: The patient has had a documented side effect, allergy or treatment failure with a preferred miscellaneous ophthalmic agent. If a product has an AB rated generic, there must have also been a trial of the generic formulation)

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>COMBINATION</u> DORZOLAMIDE w/TIMOLOL (compare to Cosopt®)</p> <p><u>MISCELLANEOUS</u> PILOCARPINE HCL</p>	<p>Cosopt® (dorzolamide w/timolol) Cosopt PF® (dorzolamide w/timolol) (pres-free) Dorzolamide w/timolol PF (compare to Cosopt PF®)</p> <p>Miochol-E® (acetylcholine) Phospholine iodide® (echothiophate)</p>	
MAST CELL STABILIZERS		
CROMOLYN SODIUM	Alocril® (nedocromil sodium) Alomide® (lodoxamide)	Criteria for Approval: The patient has had a documented side effect, allergy, or treatment failure with generic cromolyn sodium
NEUROTROPHIC KERATITIS		
All products require PA	Oxervate™ (cenegermin-bkjb) ophthalmic solution 0.002% <i>QTY LIMIT:</i> 1 vial (1mL) per eye per day Maximum of 8 weeks therapy	Oxervate: Medication is being prescribed by, or in consultation with, an ophthalmologist AND Patient has a diagnosis of Stage 2 or 3 neurotrophic keratitis (in one or both eyes) as evidenced by persistent epithelial defect or corneal ulceration AND patient has evidence of decreased corneal sensitivity in at least one corneal quadrant AND patient has failed one or more conventional non-surgical treatments such as artificial tears, gels, or ointments.
NON-STEROIDAL ANTI-INFLAMMATORY DRUGS (NSAIDs)		
<p>ACULAR LS® (ketorolac 0.4% ophthalmic solution) DICLOFENAC 0.1% ophthalmic solution FLURBIPROFEN 0.03% ophthalmic solution KETOROLAC 0.4 % ophthalmic solution (compare to</p>	Acular® (ketorolac 0.5% ophthalmic solution)	Acuvail: The patient has had a documented side effect, allergy, or treatment failure to Acular OR ketorolac 0.5% OR The patient has a documented
<p>Acular LS®) KETOROLAC 0.5 % ophthalmic solution (compare to Acular®) NEVANAC® ophthalmic suspension (nepafenac 0.1%)</p>	<p>Acuvail (ketorolac 0.45 %) Ophthalmic Solution <i>QTY LIMIT:</i> 30-unit dose packets/15 days Bromfenac 0.09 % ophthalmic solution BromSite™ (bromfenac 0.075%) solution Ilevro® ophthalmic suspension (nepafenac 0.3%) Prolensa® ophthalmic solution (bromfenac 0.07%)</p>	<p>hypersensitivity to the preservative benzalkonium chloride. All other non-preferred agents: The patient has had a documented side effect, allergy, or treatment failure to TWO preferred agents. In addition, if a product has an AB rated generic, there must have also been a trial of the generic formulation.</p>
PRESBYOPIA AGENTS		
All products require PA	Vuity™ (pilocarpine) 1.25% solution	Vuity: The patient has a diagnosis of presbyopia AND the patient is between the ages of 40-55 at the time of therapy initiation AND the medication is being prescribed by or in consultation with an optometrist or ophthalmologist AND the patient has failed corrective eyeglasses or contact lenses, unless contraindicated.

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

OTIC ANTI-INFECTIVES/ANTI-INFLAMMATORIES

**ANTI-INFECTIVE
SINGLE AGENT**

OFLOXACIN 0.3% Otic solution

Ciprofloxacin 0.2% otic solution
QTY LIMIT: 14-unit dose packages/ 7 days

**ANTI-INFECTIVE/CORTICOSTEROID
COMBINATION**

CIPRO-HC® (ciprofloxacin 0.2%/hydrocortisone 1%)
Otic suspension
NEOMYCIN/POLYMYXIN B SULFATE/
HYDROCORTISONE SOLUTION, SUSPENSION

Cortisporin-TC®
(neomycin/colistin/thonzium/hydrocortisone)

Ciprofloxacin/Dexamethasone (formerly Ciprodex®)
otic suspension

CORTICOSTEROID

FLUOCINOLONE OIL 0.01%

Ciprofloxacin/Fluocinolone otic solution
QTY LIMIT: 28-units dose packages/7days

MISCELLANEOUS AGENTS

ACETIC ACID Otic solution

DermOtic® Oil (fluocinolone acetonide) 0.01%
Flac® Oil (fluocinolone acetonide) 0.01%

Acetic Acid/Hydrocortisone Otic Solution

Anti-infective single and combination agents: The patient has had a documented side effect, allergy, or treatment failure to two preferred products.

DermOtic, Flac Oil: the patient has a documented intolerance to generic fluocinolone oil.

OVER THE COUNTER (OTC) MEDICATIONS

Please refer to the DVHA website for covered OTC categories not already managed on the PDL. Many categories limited to generics ONLY and other categories not covered. No PA process for non-covered OTCs.

https://dvha.vermont.gov/sites/dvha/files/documents/OTCWebList_0.pdf

PANCREATIC ENZYME PRODUCTS

CREON® DR Capsule
ZENPEP® DR Capsule

Pertzye® DR Capsule
Viokace® DR Capsule

Pertzye, Viokace: The patient has been started and stabilized on the requested product. OR The patient has had treatment failure or documented intolerance with both Creon and Zenpep.

PARATHYROID AGENTS

CALCITRIOL (compare to Rocaltrol®)

Doxercalciferol (compare to Hectoral®)

Doxercalciferol, Drisdol, Hectoral, Royaldee, Rocaltrol, Zemplar: The

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
CINACALCET (compare to Sensipar®) ERGOCALCIFEROL (compare to Drisdol®) PARICALCITOL (compare to Zemplar®)	Drisdol® (ergocalciferol) Hectoral® (doxercalciferol) Parsabiv™ (etelcalcetide) Rayaldee® (calcifediol ER) Rocaltrol® (calcitriol) Sensipar® (cinacalcet) Zemplar® (paricalcitol)	<p>patient must have a documented side effect, allergy, or treatment failure to two preferred agents. If a product has an AB rated generic, one trial must be the generic formulation.</p> <p>Parsabiv: indication is for the treatment of secondary hyperparathyroidism in a patient with Chronic Kidney Disease (CKD) receiving hemodialysis AND the patient has a documented side effect, allergy, or treatment failure with Sensipar. Note: treatment failure is defined as < 30% reduction from baseline in mean pre-dialysis PTH concentrations.</p> <p>Sensipar: the patient has a documented intolerance to the generic equivalent.</p>

PARKINSON'S MEDICATIONS

<p><u>DOPAMINE PRECURSOR</u> CARBIDOPA/LEVODOPA (compare to Sinemet®) CARBIDOPA/LEVODOPA ER (compare to Sinemet® CR) CARBIDOPA/LEVODOPA ODT</p> <p><u>Preferred After Clinical Criteria Are Met</u> DHIVY® (carbidopa/levodopa)</p> <p><u>DOPAMINE AGONISTS (ORAL)</u> BROMOCRIPTINE (compare to Parlodel®) PRAMIPEXOLE (compare to Mirapex®) ROPINIROLE (compare to Requip®)</p> <p><u>DOPAMINE AGONISTS (TRANSDERMAL)</u> All products require PA</p> <p><u>COMT INHIBITORS</u> ENTACAPONE (compare to Comtan®)</p> <p><u>MAO-B INHIBITORS</u> SELEGILINE</p>	<p>Inbrija® (levodopa capsule for inhalation) <i>QTY LIMIT:</i> 10 caps/day Rytary® (carbidopa/levodopa ER caps) Sinemet® (carbidopa/levodopa)</p> <p>Mirapex ER® (pramipexole ER) <i>QTY LIMIT:</i> 1 tab/day Pramipexole ER (compare to Mirapex ER®) <i>QTY LIMIT:</i> 1 tab/day Ropinirole XL <i>QTY LIMIT:</i> 12 mg = 2 tabs/day, All other strengths = 1 tab/day</p> <p>Neupro® (rotigotine) transdermal patch <i>QTY LIMIT:</i> 2, 4, 6, and 8 mg = 1 patch/day</p> <p>Comtan® (entacapone) Ongentys® (opicapone) Tasmar® (tolcapone) Tolcapone (compare to Tasmar®)</p> <p>Azilect® (rasagiline) <i>QTY LIMIT:</i> 1 mg/day Rasagiline (compare to Azilect®) <i>QTY LIMIT:</i> 1 mg/day</p>	<p>Dhivy: the patient has had a documented side effect, allergy, or treatment failure with a generic formulation of Carbidopa/Levodopa OR the patient has medical necessity for a dose that can only be achieved by splitting tablets.</p> <p>Inbrija: The patient has a diagnosis of Parkinson's disease with intermittent presence of OFF episodes AND the patient is currently taking Carbidopa/Levodopa AND the patient has had a documented side effect, allergy, or treatment failure with Apokyn®</p> <p>Comtan, Sinemet, Parlodel, Stalevo: The patient has had a documented intolerance to the generic product.</p> <p>Ongentys: The diagnosis or indication is Parkinson's disease AND the patient has had a documented side effect, allergy, or treatment failure with entacapone.</p> <p>Rytary: The patient has a diagnosis of Parkinson's disease, post-encephalitic parkinsonism, or parkinsonism following intoxication from carbon monoxide or manganese AND the prescriber is a neurologist AND the patient is having breakthrough symptoms despite a combination of concurrent IR and ER formulations of carbidopa/levodopa</p> <p>Azilect, Rasagiline: The diagnosis or indication is Parkinson's disease. AND The patient has had a documented side effect, allergy, or treatment failure with selegiline. AND The dose requested does not exceed 1 mg/day</p> <p>Gocovri: diagnosis or indication is for the treatment of dyskinesia in a patient with Parkinson's Disease AND the patient is currently receiving levodopa-based therapy (with or without concomitant dopaminergic medications) AND the patient has a documented side effect, allergy, or treatment failure with immediate release amantadine. Note: treatment failure is defined by a decrease in effectiveness despite attempts to increase dosage to 300mg/day or by temporarily discontinuing amantadine for several weeks and restarting therapy.</p> <p>Kymobi: The patient has a diagnosis of Parkinson's disease with intermittent presence of OFF episodes AND the patient is receiving concomitant</p>
---	---	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>ANTICHOLINERGICS</u> BENZTROPINE TRIHEXYPHENIDYL</p> <p><u>ADENSOSINE RECEPTOR AGONIST</u> All products require PA</p> <p><u>OTHER</u> APOKYN® (apomorphine) AMANTADINE syrup AMANTADINE capsules, tablets (PA required for ≤ 10-day supply) CARBIDOPA/LEVODOPA/ENTACAPONE (compare to Stalevo®)</p>	<p>Xadago® (safinamide) <i>QTY LIMIT: 1 tab/day</i></p> <p>Zelapar® (selegiline ODT) <i>QTY LIMIT: 2.5 mg/day</i></p> <p>Nourianz (istradefylline) <i>QTY LIMIT: 1 tab/day</i></p> <p>Gocovri™ (amantadine extended release) <i>QTY LIMIT: 2 tabs/day</i></p> <p>Kynmobi® (apomorphine) sublingual film Osmolex® ER (amantadine extended-release) <i>QTY LIMIT: 1 tablet/strength/day</i> Stalevo® (carbidopa/levodopa/entacapone)</p>	<p>levodopa which has been at a stable dose for a minimum of 4 weeks AND the patient is not taking a 5HT3 antagonist (e.g ondansetron, alosetron) concurrently AND the patient has had a documented side effect, allergy or treatment failure with Apokyn.</p> <p>Mirapex ER, Pramipexole ER, Ropinirole XL: The diagnosis or indication is Parkinson’s disease. Requests will not be approved for Restless Leg Syndrome (RLS) AND The patient has had an inadequate response (i.e. wearing off effect or “off” time) with the immediate release product. OR The patient has not been able to be adherent to a three times daily dosing schedule of the immediate release product resulting in a significant clinical impact. AND If the requested product has an AB rated generic, the patient has a documented intolerance to the generic product.</p> <p>Neupro: The patient has a medical necessity for a specialty dosage form.</p> <p>Nourianz: The patient has a diagnosis of Parkinson’s disease with intermittent presence of OFF episodes AND the patient is currently taking Carbidopa/Levodopa AND the patient has had a documented side effect, allergy, or treatment failure with TWO preferred medications being used as adjunct therapy.</p> <p>Osmolex ER: patient has not been able to be adherent to the dosing schedule of amantadine immediate release resulting in a significant clinical impact.</p> <p>Tasmar, Tolcapone: The diagnosis or indication is Parkinson’s disease. AND The patient has had a documented side effect, allergy, or treatment failure with entacapone AND patient has provided written acknowledgement of risks per the package insert. For approval of brand Tasmar, the patient must have documented intolerance to the generic equivalent.</p> <p>Xadago: The diagnosis or indication is Parkinson’s disease AND The patient is on current therapy with levodopa/carbidopa AND The patient has had a documented side effect, allergy, or treatment failure with selegiline. Note: Xadago will not be approved for monotherapy.</p> <p>Zelapar: The diagnosis or indication is Parkinson’s disease. AND The patient is on current therapy with levodopa/carbidopa. AND Medical necessity for disintegrating tablet administration is provided (i.e. inability to swallow tablets or drug interaction with oral selegiline). AND the dose requested does not exceed 2.5 mg/day</p> <p>Limitations: To prevent the use of amantadine in influenza treatment/prophylaxis, days supply < 10 days will require PA.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>corticosteroids, immunoglobulins, or splenectomy. Note: For Alvaiz, the patient must be at least 6 years old</p> <p><i>Indication for use is chronic Hepatitis-C associated thrombocytopenia:</i> The patient is at least 18 years of age AND medication is used to initiate or maintain interferon-based therapy.</p> <p><i>Indication for use is Severe Aplastic Anemia:</i> patient has had an inadequate response to standard immunosuppressive therapy (e.g. cyclosporine).</p> <p>Tavalisse: The patient is at least 18 years of age AND The diagnosis is chronic immune thrombocytopenia (ITP) AND The patient's platelet count is less than <math>30 \times 10^9/L</math> AND The patient has had a documented side effect, allergy, treatment failure or a contraindication to therapy with corticosteroids AND the patient has failed at least one of the following additional treatments: immunoglobulins, rituximab, splenectomy, or a thrombopoietin receptor agonist (e.g. eltrombopag, romiplostim, etc.). Note: Initial approval will be granted for 12 weeks. For therapy continuation, the patient must have achieved and maintained a platelet count of at least $50 \times 10^9/L$ and/or have a documented decrease in rescue treatment(s) with platelet transfusions.</p>

PSEUDOBULBAR AFFECT AGENTS

All products require PA	Nuedexta® capsules (dextromethorphan/quinidine) <i>QTY LIMIT:</i> 2 capsules/day	<p>Nuedexta: The patient must have a diagnosis of pseudobulbar affect (PBA) secondary to a neurological condition AND the patient has had a trial and therapy failure at a therapeutic dose with a tricyclic antidepressant (TCA) or an SSRI AND the patient has documentation of a current EKG (within the past 3 months) without QT prolongation AND initial authorizations will be approved for 6 months with a baseline Center for Neurologic Studies Liability Scale (CNS-LS) questionnaire AND subsequent prior authorizations will be considered at 6 month intervals with documented efficacy as seen in an improvement in the CNS-LS questionnaire</p>
-------------------------	---	--

PSORIASIS

BIOLOGICS: Initial approval is 3 months, renewals are 1 year

Preferred After Clinical Criteria Are Met

INJECTABLE

ADALIMUMAB-ADBM (compare to Cyltezo®) biosimilar to Humira®

AVSOLA® (infliximab-axxq) biosimilar to Remicade®

ENBREL® (etanercept)

QTY LIMIT: 50 mg = 8 syringes/28 days for the first 3 months; then 4 syringes/28 days
25 mg = 8 syringes/28 days subsequently

HUMIRA® (adalimumab)

QTY LIMIT: 4 syringes/28 days for one month; 2 syringes/28 days subsequently

INFLECTRA® (infliximab-dyyb) biosimilar to Remicade®

Skyrizi™ (risankizumab-rzaa)

QTY LIMIT: 150 mg/28 days for the first month and 150mg/84 days thereafter

TALTZ® (ixekizumab)

QTY LIMIT: 3 syringes/28 days for the first month, 2 syringes/28 days months 2 and 3 and 1 syringe/28 days subsequently

ORAL

Otezla® tablet (apremilast) *QTY LIMIT:* Starter Pack = 55 tablets/28 days, 30 mg = 2 tablets/day

Abrilada™ (adalimumab-afzb) biosimilar to Humira®
Adalimumab-adaz (compare to Hyrimoz®) biosimilar to Humira®

Adalimumab-adbm (compare to Cyltezo®) biosimilar to Humira®

Adalimumab-fkjp (compare to Hulio®) biosimilar to Humira®
Amjevita™ (adalimumab-atto) biosimilar to Humira®

Bimzelx® (bimekizumab-bkzx)

QTY LIMIT: 320 mg (2 syringes or autoinjectors)/28 days for the first 4 months, then 160 mg (1ml) or 320 mg (2ml)/56 days thereafter. 320 mg (2ml)/28 days maintenance dose only permitted if patient weight > 120kg

Cimzia® (certolizumab pegol)

QTY LIMIT: 1 kit/28 days (starter X 1, then regular)

Cosentyx® (secukinumab)

Cyltezo® (adalimumab-adbm) biosimilar to Humira®

Hadlima™ (adalimumab-bwvd) biosimilar to Humira®

Hulio® (adalimumab-fkjp) biosimilar to Humira®

Hyrimoz® (adalimumab-adaz) biosimilar to Humira®

Idacio® (adalimumab-aacf) biosimilar to Humira®

Ilumya™ (tildrakizumab-asmn)

QTY LIMIT: 2 ml (2 syringes) for the first month then 1 ml (1 syringe)/84 days subsequently

Remicade® (infliximab)

Renflexis™ (infliximab-abda) biosimilar to Remicade®

Siliq™ (brodalumab) injection

QTY LIMIT: 6 ml (4 syringes) for the first month then 3 ml (2 syringes)/28 days subsequently

Simlandi® (adalimumab-ryvk) biosimilar to Humira®

Spevigo® (spesolimab-sbzo)

QTY LIMIT: 900 mg (15 ml) per dose

Stelara® (ustekinumab)

QTY LIMIT: 45 mg (0.5 ml) or 90 mg (1 ml) per dose (90mg dose only permitted if patient weight > 100kg)

One dose/28 days for the first month and one dose/84 days thereafter

Tremfya® (guselkumab)

QTY LIMIT: 1 syringe/28 days for the first month, then 1 syringe every 56 days thereafter

Clinical Criteria:

For all drugs (except Spevigo): The prescription must be written by a dermatologist or rheumatologist AND The patient has a documented diagnosis of moderate to severe plaque psoriasis and has already been stabilized on the drug being requested OR The prescription must be written by a dermatologist or rheumatologist AND The patient has a documented diagnosis of moderate to severe plaque psoriasis affecting > 10% of the body surface area (BSA) and/or has involvement of the palms, soles, head and neck, or genitalia and has had a documented side effect, allergy, inadequate treatment response, or treatment failure to at least 2 different categories of therapy [i.e. at least 2 topical agents and at least 1 oral systemic agent, (unless otherwise contraindicated)] from the following categories: Topical agents: emollients, keratolytics, corticosteroids, calcipotriene, tazarotene, etc. Systemic agents: methotrexate, sulfasalazine, azathioprine, cyclosporine, tacrolimus, mycophenolate mofetil, etc. Phototherapy: ultraviolet A and topical psoralens (topical PUVA), ultraviolet A and oral psoralens (systemic PUVA, narrow band ultraviolet B (NUVA), etc.

Additional Criteria for Cimzia, Sotyktu, Taltz: The prescriber must provide evidence of a trial and failure or contraindication to a preferred TNF Inhibitor.

Additional Criteria for, Cosentyx, Siliq, Tremfya: The prescriber must provide a clinically valid reason why both a preferred TNF Inhibitor and Taltz® cannot be used. **Note:** Cosentyx approvals for 300mg dose(s) must use “300DOSE” package (containing 2x150mg pens or syringes) Approval will not be granted for 2 separate 150mg packages.

Additional Criteria for Humira Biosimilars: the patient must be unable to use Humira.

Additional Criteria for Bimzelx, Illumya, Stelara: The prescriber must provide evidence of a trial and failure or contraindication to a preferred TNF Inhibitor, Taltz, and either Tremfya or Skyrizi.

Additional Criteria for Remicade, Renflexis: The prescriber must provide a clinically valid reason why Humira®, Taltz®, and Avsola/Inflectra cannot be used.

Spevigo:

- The patient is experiencing a moderate-to-severe intensity flare of generalized pustular psoriasis (GPP) as defined by:
 - A Generalized Pustular Psoriasis Physician Global Assessment (GPPPGA) total score of at least 3 (moderate) or greater AND The presence of fresh pustules (new appearance or worsening of

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

	<p>Yuflyma® (adalimumab-aaty) biosimilar to Humira® Yusimry™ (adalimumab-aqvh) biosimilar to Humira®</p> <p>Sotyktu® (deucravacitinib) <i>QTY LIMIT</i>: 1 tablet/day</p>	<p>pustules) AND At least 5% of body surface area (BSA) covered with erythema and the presence of pustules AND</p> <ul style="list-style-type: none"> • The patient will not use concomitantly with other systemic immunosuppressants or topical agents AND <ul style="list-style-type: none"> ○ Approval will be granted for a maximum of two 900mg doses, given 7 days apart.
--	---	--

NON-BIOLOGICS

<p>ORAL ACITRETIN capsules CYCLOSPORINE (generic) METHOTREXATE (generic)</p> <p>TOPICAL CALCIPOTRIENE Cream, Ointment, Solution Tazarotene Cream, Gel Vtama® (tapinarof) cream</p>	<p>Methoxsalen (compare to Oxsoralen-Ultra®) Oxsoralen-Ultra® (methoxsalen)</p> <p>Calcitriol (compare to Vectical®) Ointment <i>QTY LIMIT</i>: 200 g (2 tubes)/week Calcipotriene Foam (compare to Sorilux®) Calcipotriene/betamethasone ointment (compare to Taclonex®) <i>QTY LIMIT</i>: Initial fill = 60 grams Duobrii™ (halobetasol propionate/tazarotene) lotion Enstilar® (calcipotriene/betamethasone) foam Sorilux® (calcipotriene) foam Taclonex® (calcipotriene/betamethasone ointment/scalp suspension) <i>QTY LIMIT</i>: Initial fill = 60 grams Zoryve® (roflumilast)</p>	<p>Duobrii lotion: the patient has had an inadequate response to at least 2 different preferred high or very high potency corticosteroids AND tazarotene cream.</p> <p>Enstilar, Taclonex or Calcipotriene/betamethasone dipropionate Ointment or Scalp Suspension: The patient has had an inadequate response to a trial (defined as daily treatment for at least one month) of a betamethasone dipropionate product and Dovonex (or generic calcipotriene), simultaneously.</p> <p>Calcipotriene Foam, Calcitriol Ointment, Sorilux, Tazarotene, Vtama: The patient has a diagnosis of mild-to-moderate plaque psoriasis AND The patient has demonstrated inadequate response, (defined as daily treatment for at least one month), adverse reaction, or contraindication to a preferred formulation of calcipotriene.</p> <p>Methoxsalen, Oxsoralen Ultra: The patient has a documented diagnosis of moderate to severe psoriasis affecting > 10% of the body surface area (BSA) and/or has involvement of the palms, soles, head and neck, or genitalia and has had a documented side effect, allergy, inadequate treatment response, or treatment failure to at least 2 topical agents and at least 1 oral systemic agent, unless otherwise contraindicated.</p> <p>Zoryve: the patient has a diagnosis consistent with the FDA indication of the requested formulation AND has had an inadequate response (defined as daily treatment for at least one month), adverse reaction, or contraindication to at least 2 different preferred agents for the diagnosis.</p> <p>Limitations: Kits with non-drug or combinations of 2 drug products are not covered.</p>
--	--	--

PULMONARY AGENTS

ANTICOLINERGICS: INHALED

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>SHORT-ACTING BRONCHODILATORS</u> ATROVENT HFA® (ipratropium) COMBIVENT® RESPIMAT (ipratropium/albuterol) <i>QTY LIMIT:</i> 3 inhalers (12 grams)/90 days IPRATROPIUM NEBULIZER SOLN IPRATROPIUM/ALBUTEROL NEBULIZER SOLN</p> <p><u>LONG-ACTING BRONCHODILATORS (LAMA)</u> INCRUSE ELLIPTA® (umeclidinium bromide) <i>QTY LIMIT:</i> 3 inhalers/90 days SPIRIVA® HANDIHALER (tiotropium) <i>QTY LIMIT:</i> 1 capsule/day SPIRIVA® RESPIMAT (tiotropium) <i>QTY LIMIT:</i> 3 inhalers/90 days</p> <p><u>COMBINATION LONG-ACTING BRONCHODILATORS (LAMA & LABA)</u> ANORO® ELLIPTA (umeclidinium/vilanterol) <i>QTY LIMIT:</i> 3 inhalers (180 blisters)/90 days STIOLTO® RESPIMAT (tiotropium/olodaterol) <i>QTY LIMIT:</i> 3 inhalers/90 days</p> <p><u>LAMA/LABA/ICS COMBINATION</u> All products require PA</p>	<p>Tiotropium bromide (compare to Spiriva®) <i>QTY LIMIT: 1 CAPSULE/DAY</i></p> <p>Tudorza® Pressair® (aclidinium bromide) <i>QTY LIMIT:</i> 3 inhalers/90 days Yupelri™ (revefenacin) inhalation solution <i>QTY LIMIT:</i> 300 vials/30 days</p> <p>Bevespi Aerosphere® (glycopyrrolate/formoterol) <i>QTY LIMIT:</i> 3 inhalers/90 days Duaklir® Pressair (aclidinium bromide/ formoterol fumarate) <i>QTY LIMIT:</i> 3 inhalers/90 days</p> <p>Breztri® Aerosphere (budesonide/glycopyrrolate/formoterol fumarate) <i>QTY LIMIT:</i> 1 inhaler (120 blisters)/30 days Trelegy® Ellipta (fluticasone/umeclidinium/vilanterol) <i>QTY LIMIT:</i> 1 inhaler (60 blisters)/30 days</p>	<p>Tudorza: The patient has had documented side effect, allergy or treatment failure with a preferred LAMA.</p> <p>Bevespi Aerosphere, Duaklir Pressiar: The patient has a documented side effect, allergy, or treatment failure to TWO preferred LAMA/LABA combinations.</p> <p>Yupelri: patient has a diagnosis of COPD (not FDA approved for asthma) AND has a failure of nebulized ipratropium solution AND at least 3 inhaled LAMAs.</p> <p>Breztri: patient has a diagnosis of COPD (not FDA approved for asthma) AND patient has a treatment failure of at least 2 different combinations of a preferred Inhaled Corticosteroid, LABA, and LAMA used in combination for a minimum of 30 consecutive days AND patient has a documented side effect, allergy, treatment failure, or contraindication with Trelegy Ellipta.</p> <p>Tiotropium bromide: The patient has had a documented intolerance to brand Spiriva</p> <p>Trelegy Ellipta: patient has a treatment failure of at least 2 different combinations of a preferred Inhaled Corticosteroid, LABA, and LAMA used in combination for a minimum of 30 consecutive days.</p>

ANTI-HISTAMINES: INTRANASAL

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>SINGLE AGENT</u> AZELASTINE 0.1% Nasal Spray OLOPATADINE 0.6% (compare to Patanase®) Nasal Spray <i>QTY LIMIT:</i> 1 bottle (31 gm)/30 days</p> <p><u>COMBO WITH CORTICOSTEROID</u> DYMISTA® (azelastine/fluticasone) Nasal Spray <i>QTY LIMIT:</i> 1 bottle (23 gm)/30 days</p>	<p>Azelastine 0.15 % Nasal Spray <i>QTY LIMIT:</i> 1 bottle (30 ml)/25 days Patanase® (olopatadine 0.6%) Nasal Spray <i>QTY LIMIT:</i> 1 bottle (31 gm)/30 days</p> <p>Azelastine/fluticasone (compare to Dymista®) Nasal Spray <i>QTY LIMIT:</i> 1 bottle (23 gm)/30 days Ryaltris® (olopatadine/mometasone) <i>QTY LIMIT:</i> 1 bottle (29 gm)/30days</p>	<p>Azelastine 0.15%: The patient has a documented side effect, allergy, or treatment failure to Azelastine 0.1%</p> <p>Azelastine/Fluticasone: The patient has a documented side effect, allergy, or treatment failure to azelastine 0.1% AND The patient has a documented side effect, allergy, or treatment failure to a preferred nasal corticosteroid OR the patient has a documented intolerance to Dymista.</p> <p>Patanase: The patient has a documented side effect, allergy, or treatment failure to Olopatadine 0.6%.</p> <p>Ryaltris: The patient has a documented side effect, allergy, or treatment failure to Olopatadine 0.6% AND The patient has a documented side effect, allergy, or treatment failure to a preferred nasal corticosteroid OR the patient has a documented intolerance to Dymista.</p>
ANTI-HISTAMINES:		
<p>Please refer to the DVHA website for covered OTC antihistamines https://dvha.vermont.gov/sites/dvha/files/documents/OTCWebList_0.pdf</p>	<p>Clarinetx® (desloratadine) 5 mg tablet Clarinetx-D® 12 hr (desloratadine/pseudoephedrine 2.5 mg/120 mg) Desloratadine (compare to Clarinetx®) 5 mg tablet Desloratadine ODT (compare to Clarinetx Reditabs®) 2.5 mg, 5 mg Levocetirizine Solution</p>	<p>LIMITATIONS: Over-the-counter antihistamines are not covered for Members Age 21 and Older.</p> <p>Clarinetx tablets, Desloratadine tablets: The patient is ≤ 20 years of age AND The patient has had a documented side effect, allergy, or treatment failure to 2 second generation antihistamines, at least one of which must be loratadine AND if the request is for Clarinetx, the patient must also have a documented intolerance to the generic equivalent tablets.</p> <p>Desloratadine ODT: The patient is ≤ 20 years of age AND The patient has had a documented side effect, allergy, or treatment failure to cetirizine oral solution and one of the following loratadine formulations: chewable tablet, rapidly disintegrating tablet, or oral solution.</p> <p>Levocetirizine solution: The patient is ≤ 20 years of age AND the patient has had a documented side effect, allergy, or treatment failure to loratadine syrup AND cetirizine syrup.</p> <p>Clarinetx-D: The patient has had a documented side effect, allergy, or treatment failure to loratadine-D and cetirizine-D.</p>
BETA-ADRENERGIC AGENTS		

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>METERED-DOSE INHALERS (SHORT-ACTING)</u> ALBUTEROL HFA (Preferred labeler codes: Teva labeler code 00093 and Sandoz labeler code 00781) PROAIR[®] Respiclick (albuterol) VENTOLIN[®] HFA (albuterol) XOPENEX[®] HFA (levalbuterol)</p> <p><u>METERED-DOSE INHALERS (LONG-ACTING)</u> <i>Preferred After Clinical Criteria Are Met</i> SEREVENT[®] DISKUS (salmeterol xinafoate) <i>QTY LIMIT: 3 inhalers (180 blisters)/90 days</i></p> <p><u>NEBULIZER SOLUTIONS (SHORT-ACTING)</u> ALBUTEROL neb solution (all strengths) LEVALBUTEROL neb solution (age ≤ 12 years)</p> <p><u>NEBULIZER SOLUTIONS (LONG-ACTING)</u> All products require PA</p> <p><u>TABLETS/SYRUP (SHORT-ACTING)</u> ALBUTEROL tablets/syrup</p>	<p>Albuterol HFA (all other labelers) Levalbuterol Aerosol (compare to Xopenex[®] HFA)</p> <p>Striverdi Respimat[®] (olodaterol)</p> <p>Levalbuterol neb solution (compare to Xopenex[®]) (age > 12 years)</p> <p>Arformoterol (compare to Brovana[®]) <i>QTY LIMIT: 2 vials/day</i> Brovana[®] (arformoterol) <i>QTY LIMIT: 2 vials/day</i> Formoterol (compare to Perforomist[®]) <i>QTY LIMIT: 2 vials/day</i> Perforomist[®] (formoterol) <i>QTY LIMIT: 2 vials/day</i></p> <p>Terbutaline tablets</p>	<p>Albuterol HFA: The patient has a documented side effect, allergy, or treatment failure to two preferred short acting metered dose inhalers.</p> <p>Levalbuterol HFA: The patient has a documented intolerance to brand Xopenex HFA.</p> <p>Serevent: The patient has a diagnosis of asthma and is prescribed an inhaled corticosteroid (pharmacy claims will be evaluated to assess compliance with long term controller therapy) OR the patient has a diagnosis of COPD.</p> <p>Striverdi: The patient has a diagnosis of COPD (not FDA approved for asthma). AND The patient has a documented side effect, allergy, or treatment failure to Serevent.</p> <p>Levalbuterol nebulizer solution (age > 12 years): The patient must have had a documented side effect, allergy, or treatment failure to albuterol nebulizer. Arformoterol, Brovana, Formoterol, Perforomist Nebulizer Solution: The patient must have a diagnosis of COPD. AND The patient must be unable to use a non-nebulized long-acting bronchodilator or anticholinergic (Serevent or Spiriva) due to a physical limitation AND for approval of Brovana, Formoterol, or Perforomist, the patient must also have a documented intolerance or treatment failure with arformoterol.</p> <p>Terbutaline tablets: The medication is not being prescribed for the prevention/treatment of preterm labor.</p>

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

METERED DOSE INHALERS (SINGLE AGENT)

ARNUITY ELLIPTA (fluticasone furoate)
QTY LIMIT: 90 blisters/90 days
ASMANEX[®] Twisthaler[®] (mometasone furoate)
QTY LIMIT: 3 inhalers/90 days
PULMICORT FLEXHALER[®] (budesonide)
QTY LIMIT: 6 inhalers/90 days
QVAR REDIHALER[®] (beclomethasone dipropionate) 40mcg/inh
QTY LIMIT: 2 inhalers (21.2 gm)/90 days
QVAR REDIHALER[®] 80mcg/inh
QTY LIMIT: 3 inhalers (31.8 gm)/90 days
ASMANEX[®] HFA (mometasone furoate)
QTY LIMIT: 3 inhalers (39gm)/90 days
FLUTICASONE PROPIONATE[®] HFA (compare to Flovent HFA) Age ≤ 5 years:
QTY LIMIT: 3 inhalers (36gm)/90 days

METERED DOSE INHALERS (COMBINATION PRODUCT)

ADVAIR[®] DISKUS (fluticasone/salmeterol) (Age ≥ 4 years)
QTY LIMIT: 3 inhalers/90 days
ADVAIR[®] HFA (fluticasone/salmeterol) (Age ≥ 12 years)
QTY LIMIT: 3 inhalers (36 gm)/90 days
AIRDUO RESPICLICK[®] (fluticasone/salmeterol)
QTY LIMIT: 3 inhalers/90 days
DULERA[®] (mometasone/formoterol)
QTY LIMIT: 9 inhalers (117 gm)/90 days
SYMBICORT[®] (budesonide/formoterol)
QTY LIMIT: 9 inhalers (91.8gm)/90 days

Armonair[®] Digihaler (fluticasone propionate)
QTY LIMIT = 3 inhalers/90 days
Alvesco[®] (ciclesonide)
QTY LIMIT: 80 mcg = 3 inhalers/90 days
Airsupra[®] (albuterol/budesonide inhalation)
QTY LIMIT: 3 inhalers (32.1 gm)/30 days
Fluticasone propionate Diskus (compare to Flovent[®] Diskus)
QTY LIMIT: 180 listers/90 days)
Fluticasone propionate HFA (compare to Flovent[®] HFA) Age ≥ 6 years
QTY LIMIT: 3 inhalers (36 gm)/90 days

AirDuo[®] Digihaler (fluticasone/salmeterol)
QTY LIMIT: 3 inhalers/90 days
Breo Ellipta[®] (fluticasone furoate/vilanterol)
QTY LIMIT: 3 inhalers (180 blisters)/ 90 days
Breyna[™] (budesonide/formoterol) *QTY LIMIT*: 9 inhalers (92.7gm)/90 days
Budesonide/formoterol (compare to Symbicort[®])
QTY LIMIT: 9 inhalers (91.8gm)/90 days
Fluticasone furoate/vilanterol (compare to Breo Ellipta[®])
QTY LIMIT: 3 inhalers (180 blisters)/90 days
Fluticasone/salmeterol (compare to AirDuo Resplick[®])
QTY LIMIT: 3 inhalers/90 days
Fluticasone/salmeterol inhalation Powder (compare to Advair[®] Diskus)
QTY LIMIT: 3 inhalers/90 days
Wixela[™] Inhub[™] (fluticasone/salmeterol inhalation powder) (compare to Advair[®] Diskus)
QTY LIMIT: 3 inhalers/90 days

Armonair Digihaler, Alvesco, Fluticasone Diskus: The patient has had a documented side effect, allergy, or treatment failure to at least two preferred agents.
Fluticasone HFA (Age ≥ 6 years): The patient has had a documented side effect, allergy, or treatment failure to at least two preferred agents, one of which must be Asmanex HFA. OR Medical necessity for the use of an HFA formulation has been provided and the patient has a documented side effect, allergy, or treatment failure to Asmanex HFA.
Advair HFA (age < 12 years): The patient has had a documented side effect, allergy, or treatment failure to Dulera or Symbicort.
AirDuo Digihaler, Breo Ellipta, Fluticasone Furoate/Vilanterol, Fluticasone/Salmeterol (non-authorized generics): The patient has had a documented side effect, allergy, or treatment failure to any 2 of the following: Advair HFA, Advair Diskus, Airduo Resplick, Dulera, or Symbicort AND for approval of Fluticasone Furoate/Vilanterol, the patient must also have a documented intolerance to Breo Ellipta.
Airsupra: The patient is ≥ 18 years of age AND the patient has had a documented side effect, allergy, treatment failure or a contraindication to Symbicort and Dulera being used as needed for asthma exacerbations (SMART therapy) AND the patient is unable to use Albuterol and Budesonide as individual agents.
Breyna, Budesonide/formoterol: the patient has a documented intolerance to brand Symbicort.
Fluticasone/salmeterol powder (authorized generic), Wixela Inhub: A clinically compelling reason must be provided detailing why the patient is unable to use Advair HFA or Advair Diskus.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>NEBULIZER SOLUTIONS BUDESONIDE INH SUSPENSION 0.25mg, 0.5mg (Age ≤ 12 yrs)</p>	<p>Budesonide Inh Suspension 1mg (all ages), 0.25mg and 0.5mg (age >12 years) Pulmicort Respules[®] (budesonide)</p>	<p>Budesonide Inh Suspension: Medical necessity for the use of a nebulized solution has been provided AND if the dose is 1mg, the patient must be unable to use two 0.5 mg vials Pulmicort Respules: medical necessity for the use of a nebulized solution has been provided AND if the dose is 1 mg, the patient must be unable to use two 0.5 mg vials AND the patient has a documented intolerance to the generic.</p>
CORTICOSTEROIDS: INTRANASAL		
<p>BUDESONIDE <i>QTY LIMIT:</i> 1 inhaler (8.43 ml)/30 days FLUTICASONE PROPIONATE <i>QTY LIMIT:</i> 1 inhaler (16 gm)/30 days MOMETASONE <i>QTY LIMIT:</i> 1 inhaler (17 gm)/30 days OMNARIS[®] (ciclesonide) <i>QTY LIMIT:</i> 1 inhaler (12.5 gm)/30 days TRIAMCINOLONE <i>QTY LIMIT:</i> 1 inhaler (16.9 ml)/30 days ZETONNA[®] (ciclesonide) <i>QTY LIMIT:</i> 1 inhaler (6.1 gm)/30 days</p>	<p>Beconase AQ[®] (beclomethasone) <i>QTY LIMIT:</i> 2 inhalers (50 gm)/30 days Flunisolide 25 mcg/spray <i>QTY LIMIT:</i> 2 inhalers (50 ml)/30 days QNASL[®] (beclomethasone dipropionate) <i>QTY LIMIT:</i> 1 inhaler (10.6 gm)/30 days Xhance[™] (fluticasone propionate) <i>QTY LIMIT:</i> 1 inhaler (16 ml)/30 days</p>	<p>Beconase AQ, Flunisolide 25 mcg/spray, QNASL: The patient has had a documented side effect, allergy, or treatment failure of two preferred nasal glucocorticoids. If a product has an AB rated generic, one trial must be the generic. Xhance: The patient has had a documented side effect, allergy, or treatment failure of three preferred nasal glucocorticoids, one of which must be fluticasone. Limitations: Nasacort Allergy OTC and Flonase are not covered as no Federal Rebate is offered.</p>
LEUKOTRIENE MODIFIERS		
<p><i>Preferred After Age Criteria Are Met</i> MONTELUKAST SODIUM (compare to Singulair[®]) tablets, 10mg for ages ≥ 15 MONTELUKAST SODIUM (compare to Singulair[®]) chews, 4 mg for ages 2-5, 5 mg for age 6-14 MONTELUKAST SODIUM (compare to Singulair[®]) granules, ages 6 months-23 months</p>	<p>Accolate[®] (zafirlukast) <i>QTY LIMIT:</i> 2 tablets/day Singulair[®] (montelukast sodium) tablets, chew tabs, granules <i>QTY LIMIT:</i> 1 tablet or packet per day Zafirlukast (compare to Accolate[®]) Zileuton ER (compare to Zyflo CR[®]) <i>QTY LIMIT:</i> 4 tablets/day Zyflo (zileuton) <i>QTY LIMIT:</i> 4 tablets/day</p>	<p>Montelukast: Clinical rationale must be provided for prescribing a dose and formulation that differs from age recommendations AND If the request is for brand Singulair, the patient has a documented intolerance to the generic equivalent montelukast preparation. Zafirlukast, Accolate: The diagnosis or indication for the requested medication is asthma. AND If the request is for Accolate, the patient has a documented intolerance to generic zafirlukast. Zileuton ER, Zyflo: The diagnosis or indication for the requested medication is asthma. AND The patient has had a documented side effect, allergy, or treatment failure to Accolate/Zafirlukast or Singulair/Montelukast</p>
PHOSPHODIESTERASE-4 (PDE-4) INHIBITORS		
<p>All products require PA</p>	<p>Daliresp[®] tablet (roflumilast) <i>QTY LIMIT:</i> 1 tablet/day Roflumilast (compare to Daliresp) tablet <i>QTY LIMIT:</i> 1 tablet/day * Maximum days' supply per fill = 30 *</p>	<p>Daliresp, Roflumilast: The indication for the requested medication is treatment to reduce the risk of COPD exacerbations in patients with severe COPD associated with chronic bronchitis and a history of exacerbations. AND The patient has had a documented side effect, allergy, treatment failure, or a contraindication to at least one inhaled long-acting anticholinergic AND at least one inhaled long-acting beta-agonist AND at least one inhaled corticosteroid AND for approval of brand name Daliresp, the patient has had a documented intolerance to the generic equivalent.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
SYNAGIS	SYNAGIS® (palivizumab) <i>QTY LIMIT:</i> 50 mg = 1 vial/month, 100 mg = 2 vials/month	CRITERIA FOR APPROVAL: <ul style="list-style-type: none"> • If infant is in their first RSV season, the healthcare provider must provide clinical reasoning as to why infant cannot receive Beyfortus NOTE: Beyfortus is managed through the Vaccines for Children (VFC) program • If infant is 8-19 months old AND entering their second RSV season AND has ONE of the following risk factors: <ul style="list-style-type: none"> • Chronic lung disease of prematurity who required medical support any time during the 6-month period before the start of the second RSV season • Severely immunocompromised children • Children with cystic fibrosis who have either manifestation or severe lung disease OR weight for length < 10th percentile • American Indian or Alaska Native children, Provider must provide clinical reasoning why they cannot receive a second dose of Beyfortus • The prescriber must confirm the member has not already received Beyfortus for the current RSV season. Concomitant use with Beyfortus will not be approved. • Infants born at 28 weeks of gestation or earlier (i.e., ≤ 28 weeks, 6 days) and under twelve months of age at the start of the RSV season (maximum 5 doses). • Infants born at 29-32 weeks (i.e., between 29 weeks, 0 days and 31 weeks, 6 days) of gestation and under 1 year of age at the start of the RSV season who develop chronic lung disease of prematurity defined as a requirement for >21% oxygen for at least the first 28 days after birth (maximum 5 doses). • Children under 24 months of age with chronic lung disease of prematurity defined as born at 31 weeks, 6 days or less who required >21% oxygen for at least the first 28 days after birth and continue to require medical support (chronic corticosteroid therapy, diuretic therapy, or supplemental oxygen) during the 6-month period before the start of the second RSV season (maximum 5 doses). • Children under 12 months of age with hemodynamically significant congenital heart disease (CHD) (dosing continues in the RSV season through the end of the month the infant reaches 12 months old - maximum 5 doses): Acyanotic heart disease and receiving medication to control congestive heart failure and will require cardiac surgical procedures, Moderate to severe pulmonary hypertension, Cyanotic heart disease and recommended for Synagis therapy by Pediatric Cardiologist

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<ul style="list-style-type: none"> • Infants under 12 months of age with either: (dosing continues in the RSV season through the end of the month the infant reaches 12 months old -maximum 5 doses) Congenital abnormalities of the airways that impairs the ability to clear secretions from the upper airway because of ineffective cough, Neuromuscular condition that impairs the ability to clear secretions from the upper airway because of ineffective cough • Infants and children less than 24 months of age who will undergo a heart transplant during the RSV season • Infants and children less than 24 months of age who are profoundly immunocompromised during the RSV season (e.g. undergoing organ or stem cell transplant or receiving chemotherapy). <p>EXCLUDED FROM APPROVAL:</p> <ul style="list-style-type: none"> • Infants and children with hemodynamically insignificant heart disease. • Infants with cardiac lesions adequately corrected by surgery, unless they continue to require medication for congestive heart failure. • Infants with mild cardiomyopathy who are not receiving medical therapy. • Breakthrough hospitalization for RSV disease (Synagis therapy should be discontinued for the season once hospitalization for RSV has occurred). • Infants and children with Down syndrome unless other indications above are present. • Infants and children with cystic fibrosis unless other specific conditions are present

PULMONARY ARTERIAL HYPERTENSION MEDICATIONS

<p><u>ENDOTHELIN RECEPTOR ANTAGONISTS</u> AMBRISENTAN (compare to Letairis®) <i>QTY LIMIT:</i> 1 tablet/day BOSENTAN (compare to Tracleer) <i>QTY LIMIT:</i> 2 tablets/day</p>	<p>Letairis® (ambrisentan) Tablet <i>QTY LIMIT:</i> 1 tablet/day Opsumit® (macitentan) Tablet <i>QTY LIMIT:</i> 1 tablet/day Opsynvi® (macitentan/tadalafil) tablets Tracleer® tablets for oral suspension (32 mg) Tracleer® (bosentan) tablet (62.5 mg, 125 mg) <i>QTY LIMIT:</i> 2 tablets/day</p>	<p>Adempas: The patient has a diagnosis of pulmonary arterial hypertension (PAH) with New York Heart Association (NYHA) Functional Class II or III. OR The patient has a diagnosis of chronic thromboembolic pulmonary hypertension (CTEPH, WHO Group 4) AND the patient has persistent or recurrent disease after surgical treatment (e.g., pulmonary endarterectomy) or has CTEPH that is inoperable AND The patient is 18 years of age or older AND The patient will not use Adempas concomitantly with the following: Nitrates or nitric oxide donors (such as amyl nitrate) in any form Phosphodiesterase (PDE) inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline) AND The patient is not pregnant AND Female patients are enrolled in the Adempas REMS Program</p> <p>Flolan, Letairis, Tracleer: patient has a documented intolerance to the generic equivalent.</p>
---	---	---

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p><u>PROSTACYCLIN AGONISTS</u> <u>INJECTION</u> EPOPROSTENOL (compare to Flolan®) REMODULIN® (treprostinil sodium injection) VELETRI® (epoprostinil)</p> <p><u>INHALATION</u> All products require PA</p> <p><u>ORAL</u> ORENITRAM® (treprostinil) ER Tablet</p> <p><u>sGC STIMULATOR</u> All products require PA</p> <p>**Maximum days supply for all drugs is 30 days**</p>	<p>Flolan® (epoprostenol) Treprostinil sodium injection (compare to Remodulin®) Winrevair™ (sotatercept-csrk)</p> <p>Tyvaso® (Treprostinil) inhalation solution Tyvaso® DPI (treprostinil) powder for inhalation Ventavis® (iloprost) inhalation solution</p> <p>Uptravi® (selexipag) tablets <i>QTY LIMIT:</i> 200 mcg = 140 tablets/30 days for the first 2 months, then 2 tablets/day thereafter All other strengths = 2 tablets/day</p> <p>Adempas® (riociguat) Tablets <i>QTY LIMIT:</i> 3 tablets/day</p>	<p>Tracleer tablets for oral suspension: Patient has a diagnosis of PAH with NYHA Functional Class II or III AND patient is ≤ 12 years of age and <40kg.</p> <p>Opsumit, Opsyvni: Patient has a diagnosis of PAH with NYHA Functional Class II or III AND Patient is not pregnant AND Female patients have been enrolled in the REMS Program AND the patient has a documented side effect, allergy, or treatment failure with bosentan or ambrisentan. Additional criteria for Opsyvni: the patient is unable to tolerate the individual ingredient medications.</p> <p>Treprostinil: Patient has a diagnosis of pulmonary arterial hypertension AND The patient has had a documented intolerance to the brand Remodulin.</p> <p>Tyvaso, Ventavis: The patient has a diagnosis of pulmonary arterial hypertension (PAH) with New York Heart Association (NYHA) Functional Class II or III heart failure AND the patient is unable to tolerate or has failed 2 different preferred medications.</p> <p>Uptravi: The patient has a diagnosis of pulmonary arterial hypertension (PAH) with New York Heart Association (NYHA) Functional Class II or III heart failure AND the patient is unable to tolerate or has failed 2 different preferred medications, one of which must be Orenitram</p> <p>Winrevair: The patient has a diagnosis of pulmonary arterial hypertension (PAH) Group 1 and WHO functional class II or III AND</p> <ul style="list-style-type: none"> • The patient must have LV end diastolic pressure of ≤ 15mmHg AND • The patient must be on stable doses of standard of care therapy (patient-specific dose goal for each PAH therapy achieved) AND • The patient has 6MWT between 150-500 meters AND • The patient has used dual combination therapy from two other PAH-indicated drug classes, unless contraindicated, for at least 60 days and has experienced continued decline in pulmonary hemodynamics and exercise capacity
<p>PHOSPHODIESTERASE-5 (PDE-5) INHIBITORS Effective 7/1/06, phosphodiesterase-5 (PDE-5) inhibitors are no longer a covered benefit for all Vermont Pharmacy Programs for the treatment of erectile dysfunction. This change is resultant from changes set into effect January 1, 2006 and as detailed in Section 1903 (i)(21)(K) of the Social Security Act (the Act), precluding Medicaid Federal Funding for outpatient drugs used for the treatment of sexual or erectile dysfunction. Sildenafil will remain available for coverage via prior-authorization for the treatment of Pulmonary Arterial Hypertension.</p>		
<p><i>Preferred After Clinical Criteria Are Met</i> SILDENAFIL CITRATE (compare to Revatio®) tablet <i>QTY LIMIT:</i> 3 tablets/day SILDENAFIL suspension TADALAFIL (compare to Adcirca®) <i>QTY LIMIT:</i> 2 tablets/day</p>	<p>Adcirca® (tadalafil) <i>QTY LIMIT:</i> 2 tablets/day Revatio® (sildenafil) tabs <i>QTY LIMIT:</i> 3 tablets/day Revatio® (sildenafil citrate) vial <i>QTY LIMIT:</i> 3 vials/day Maximum 14-day supply per fill Sildenafil (compare to Revatio®) vial</p>	<p>Sildenafil tablet, Tadalafil tablet: Clinical Diagnosis of Pulmonary Arterial Hypertension Adcirca (tadalafil) 20 mg, Revatio (sildenafil citrate) 20 mg: Clinical diagnosis of pulmonary arterial hypertension AND No concomitant use of organic nitrate-containing products AND patient has a documented intolerance to the generic equivalent. Sildenafil Suspension: Clinical diagnosis of pulmonary arterial hypertension AND medical necessity for a liquid formulation is provided Revatio IV, Sildenafil IV: Clinical diagnosis of pulmonary arterial</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
	Tadliq® (tadalafil) suspension	<p>hypertension AND no concomitant use of organic nitrate-containing products AND The patient has a requirement for an injectable dosage form. AND Arrangements have been made for IV bolus administration outside of an inpatient hospital setting.</p> <p>Tadliq: Clinical diagnosis of pulmonary arterial hypertension AND medical necessity for a liquid formulation is provided AND the patient has a documented side effect, allergy, or treatment failure with sildenafil suspension.</p>

RENAL DISEASE: PHOSPHATE BINDERS

<p>CALCIUM ACETATE capsule CALCIUM ACETATE tablet SEVELAMER CARBONATE (compare to Renvela®) tablets</p> <p><u>ORAL SOLUTIONS</u> PHOSLYRA® (calcium acetate) oral solution</p>	<p>Auryxia® (ferric citrate) <i>QTY LIMIT:</i> 12/day Fosrenol® (lanthanum carbonate) Lanthanum carbonate (compare to Fosrenol) Renagel® (sevelamer) Renvela® (sevelamer carbonate) Oral Suspension Packet <i>QTY LIMIT:</i> 0.8 g = 2 packs/day Renvela® (sevelamer carbonate) tablets Sevelamer carbonate Oral Suspension Packet (compare to Renvela®) <i>QTY LIMIT:</i> 0.8 g = 2 packs/day Sevelamer hydrochloride (compare to Renagel®) Velphoro® (sucroferric oxyhydroxide) Chew Tablet Xphozah® (tenapanor) tablet</p>	<p>Renvela Oral Suspension Packet, Sevelamer Packet: The patient has a requirement for a liquid dosage form.</p> <p>Auryxia, lanthanum carbonate, Renagel, Renvela tablets, sevelamer hydrochloride tablets, Velphoro Chew Tablet: The patient must have a documented side effect, allergy, or inadequate response to one preferred phosphate binder.</p> <p>Xphozah: The patient must have a documented side effect, allergy, or inadequate response to two non-calcium-containing binders (sevelamer, lanthanum, ferric citrate, and sucroferric oxyhydroxide)</p>
---	---	---

RESTLESS LEG SYNDROME MEDICATIONS

<p><u>DOPAMINE AGONISTS (ORAL)</u> PRAMIPEXOLE (compare to Mirapex®) ROPINIROLE (compare to Requip®)</p> <p><u>DOPAMINE AGONISTS (TRANSDERMAL)</u></p>	<p>Mirapex® (pramipexole)</p> <p>Neupro® (rotigotine) transdermal patch <i>QTY LIMIT:</i> 1, 2, and 3 mg ONLY = 1 patch/day</p>	<p>Mirapex: The patient has had a documented intolerance to the generic product. Neupro: The patient has a medical necessity for a specialty dosage form. Horizant: The patient has a diagnosis of restless legs syndrome (RLS). AND The patient has had a documented side effect, allergy, contraindication or treatment failure to two preferred dopamine agonists AND gabapentin IR.</p>
<p>All products require PA</p> <p><u>GAMMA-AMINOBUTYRIC ACID ANALOG</u> GABAPENTIN IR</p>	<p>Horizant® (gabapentin enacarbil) ER Tablet <i>QTY LIMIT:</i> 1 tablet/day</p>	<p>Limitations: Requests for Mirapex ER and Requip XL will not be approved for Restless Leg Syndrome (RLS).</p>

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

RHEUMATOID, JUVENILE & PSORIATIC ARTHRITIS: IMMUNOMODULATORS

Length of Authorization: Initial PA 3 months; 12 months thereafter

Preferred After Clinical Criteria Are Met
INJECTABLE

ADALIMUMAB-ADBM (compare to Cyltezo®) biosimilar to Humira®

AVSOLA® (infliximab-axxq) biosimilar to Remicade®

ENBREL® (etanercept)
QTY LIMIT: 50 mg = 4 syringes/28 days, 25 mg = 8 syringes/28 days

INFLECTRA® (infliximab-dyyb) biosimilar to Remicade®

KINERET® (anakinra)
QTY LIMIT: 1 syringe/day

HUMIRA® (adalimumab)
QTY LIMIT: 4 syringes/28 days

SKYRIZI™ (risankizumab-rzaa)
QTY LIMIT: 150 mg/28 days for the first month and 150mg/84 days thereafter

TALTZ® (ixekizumab)
QTY LIMIT: 80 mg prefilled syringe or autoinjector = 2/28 days for the first month and 1/28 days subsequently

TYENNE® (tocilizumab-aazg) biosimilar to Actemra®

Abrilada™ (adalimumab-afzb) biosimilar to Humira®

Actemra® (tocilizumab) Intravenous Infusion
QTY LIMIT: 80 mg vial = 4 vials/28 days, 200 mg vial = 3 vials/28 days, 400 mg vial = 2 vials/28 days

Actemra® (tocilizumab) Subcutaneous Prefilled Syringe
QTY LIMIT: 4 prefilled syringes (3.6ml)/28 days

Actemra® (tocilizumab) ACTPen

QTY LIMIT: 4 pens (3.6ml)/28 days

Adalimumab-adaz (compare to Hyrimoz®) biosimilar to Humira®

Adalimumab-adbm (compare to Cyltezo®) biosimilar to Humira®

Adalimumab-fkjp (compare to Hulio®) biosimilar to Humira®

Amjevita™ (adalimumab-atto) biosimilar to Humira®
Cimzia® (certolizumab pegol)
QTY LIMIT: 1 kit/28 days

Cosentyx® (secukinumab)

Cyltezo® (adalimumab-adbm) biosimilar to Humira®

Hadlima™ (adalimumab-bwvd) biosimilar to Humira®

Hulio® (adalimumab-fkjp) biosimilar to Humira®

Hyrimoz® (adalimumab-adaz) biosimilar to Humira®

Idacio® (adalimumab-aacf) biosimilar to Humira®

Kevzara® (sarilumab)
QTY LIMIT: 2 syringes/28 days

Ilaris® (canakinumab)

Orencia® (abatacept) Subcutaneous Injection

QTY LIMIT: 4 syringes/28 days

Orencia® (abatacept) Intravenous Infusion

Remicade (infliximab)

Renflexis™ (Infliximab-abda) biosimilar to Remicade®

Simlandi® (adalimumab-ryvk) biosimilar to Humira®

Simponi® (golimumab) Subcutaneous
QTY LIMIT: 50 mg = 1 prefilled syringe or autoinjector/28 days

Simponi Aria® (golimumab) 50 mg/4 ml Vial for

Clinical Criteria for all drugs: Patient has a diagnosis of rheumatoid arthritis (RA), juvenile idiopathic arthritis* or psoriatic arthritis and has already been stabilized on the drug being requested OR Diagnosis is RA, juvenile idiopathic arthritis or psoriatic arthritis, and methotrexate therapy resulted in an adverse effect, allergic reaction, inadequate response, or treatment failure. If methotrexate is contraindicated, another DMARD should be tried prior to approving therapy. Other DMARDs include leflunomide, sulfasalazine, gold, antimalarials, minocycline, D-penicillamine, azathioprine, cyclophosphamide and cyclosporine

Taltz, Xeljanz, Xeljanz XR additional criteria: patient must be ≥ 18 years of age AND the prescriber must provide evidence of a trial and failure or contraindication to a preferred TNF Inhibitor. **Note:** Xeljanz 10mg BID and XR 22mg are NOT recommended for Rheumatoid Arthritis or Psoriatic Arthritis. Please refer to Gastrointestinal: Inflammatory Bowel Disease Biologics for Ulcerative Colitis criteria.

Cosentyx additional criteria: the prescriber must provide evidence of a trial and failure or contraindication to a preferred TNF Inhibitor and Taltz.

Actemra, Kevzara, Orencia, Tofidence, and Tremfya additional criteria: The prescriber must provide clinically valid reason why at least 2 preferred agents cannot be used. **For approval of Actemra, patient has had a documented side effect, allergy, or treatment failure to Tyenne.**

Humira Biosimilars Additional Criteria: The patient must be unable to use Humira.

Ilaris: The diagnosis is systemic juvenile idiopathic arthritis (sJIA) with active systemic features and varying degrees of synovitis with continued disease activity after initial therapy (initial therapy defined as 1 month of anakinra (Kineret), 2 weeks of glucocorticoid monotherapy (oral or IV) or one month of NSAIDs). AND patient is > 2 years of age.

Remicade, Renflexis additional criteria: The prescriber must provide a clinically valid reason why at least 2 preferred agents cannot be used AND the patient must be unable to use Avsola or Inflectra.

Simponi Aria additional criteria: The patient has not responded adequately to Simponi subcutaneous. AND The prescriber must provide a clinically valid reason why at least 2 preferred agents cannot be used.

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>ORAL</p> <p>OTEZLA® tablet (apremilast) <i>QTY LIMIT:</i> Starter Pack = 55 tablets/28 days, 30 mg = 2 tablets/day Maximum 30 days supply</p> <p>XELJANZ® (tofacitinib) 5 mg tablet <i>QTY LIMIT:</i> 2 tablets/day Maximum 30 days supply</p> <p>XELJANZ® XR (tofacitinib) tablet <i>QTY LIMIT:</i> 1 tablet/day</p> <p>XELJANZ® (tofacitinib) oral solution</p>	<p>Intravenous Infusion</p> <p>Stelara® (ustekinumab) <i>QTY LIMIT:</i> 45 mg (0.5 ml) or 90 mg (1 ml) per dose (90 mg dose only permitted for pt weight > 100 kg) One dose/28 days for the first month and one dose/84 days thereafter</p> <p>Tofidence (tocilizumab-bavi) biosimilar to Actemra Intravenous Infusion <i>QTY LIMIT:</i> 80mg vial=4 vials/28 days, 200mg vial=3 vials/28 days, 400mg vial=2 vials/28 days</p> <p>Tyenne® (tocilizumab-aazg) biosimilar to Actemra®</p> <p>Tremfya® (guselkumab) <i>QTY LIMIT:</i> 1 syringe/28 days for the first month, then 1 syringe every 56 days thereafter</p> <p>Yuflyma® (adalimumab-aaty) biosimilar to Humira®</p> <p>Yusimry™ (adalimumab-aqvh) biosimilar to Humira®</p> <p>Olumiant® (baricitinib) tablets <i>QTY LIMIT:</i> 1 tablet/day Maximum 30 days supply</p> <p>Rinvoq® (upadactinib) extended-release tablet <i>QTY LIMIT:</i> 1 tablet/day Maximum 30 days supply</p>	<p>Stelara additional criteria: the prescriber must provide evidence of a trial and failure or contraindication to a preferred TNF Inhibitor, Taltz, and Tremfya</p> <p>Olumiant, Rinvoq additional criteria: The patient must be ≥ 18 years of age AND The prescriber must provide a clinically valid reason why at least two preferred agents cannot be used, one of which must be Xeljanz or Xeljanz XR.</p> <p>Note: Patients with systemic juvenile arthritis (SJRA/SJIA) and fever are not required to have a trial of a DMARD, including methotrexate. Patients with systemic juvenile arthritis without fever should have a trial of methotrexate, but a trial of another DMARD in the case of a contraindication to methotrexate is not required. * Patients with psoriatic arthritis with a documented diagnosis of active axial involvement should have a trial of NSAID therapy, but a trial with DMARD is not required before a TNF-blocker is approved. If no active axial skeletal involvement, then an NSAID trial and a DMARD trial are required (unless otherwise contraindicated).</p>
SICKLE CELL DISEASE THERAPIES		
<p>DROXIA® (hydroxyurea) 200 mg, 300 mg, 400 mg cap</p> <p>HYDROXYUREA (compare to Hydrea®) 500 mg cap</p> <p><u>Preferred After Clinical Criteria Are Met</u></p> <p>ENDARI® (L-glutamine powder for oral solution) <i>QTY LIMIT:</i> maximum of 30-day supply</p>	<p>Adakveo® (crizanlizumab-tmca)</p> <p>Casgevy™ (exagamglogene autotemcel)</p> <p>Hydrea® (hydroxyurea) 500 mg cap</p> <p>Lyfgenia™ (lovotibeglogene autotemcel)</p> <p>Siklos® (hydroxyurea) 100 mg, 1000 mg tablet</p>	<p>Adakveo: Patient has a diagnosis of Sickle Cell Disease AND patient is at least 16 years of age or older AND patient has had an inadequate response to a 6-month trial of hydroxyurea dosed at 15-35 mg/kg/day, unless contraindicated AND patient has experienced at least 2 vaso-occlusive crises in the previous 12 months despite compliance with hydroxyurea. Initial approval will be granted for 6 months. For re-approval, the patient must have a decrease in the frequency or severity of VOC compared to baseline.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
		<p>Casgevy, Lyfgenia: Patient has a diagnosis of Transfusion-Dependent β-thalassemia (Casgevy only) OR Patient has a diagnosis of Sickle Cell Disease AND patient is at least 12 years of age or older AND patient has had an inadequate response to a 6-month trial of hydroxyurea dosed at 15-35 mg/kg/day, unless contraindicated AND patient has experienced at least 2 vaso-occlusive crises in the previous 12 months despite compliance with hydroxyurea AND Patient has not previously received gene therapy for sickle cell disease AND Patient meets approved clinical parameters for use (i.e. no contraindications, appropriate monitoring has been completed.) AND for approval of Lyfgenia, the patient must have a contraindication to use of Casgevy</p> <p>Endari: Indication for use is to reduce the acute complications of Sickle Cell Anemia AND medication will be approved with quantity limits based on patient weight (<30kg = 2 packets/day, 30-65kg = 4 packets/day, > 65kg = 6 packets/day).</p> <p>Hydrea: Patient has had a documented intolerance to the generic equivalent.</p> <p>Siklos: Patient has a diagnosis of Sickle Cell Anemia with recurrent moderate to severe pain crises AND the required dose is < 200mg OR Patient has a diagnosis of Sickle Cell Anemia with recurrent moderate to severe pain crises AND has a documented intolerance to a preferred hydroxyurea formulation. For re-approval, the patient must have a documented decrease in vaso-occlusive episodes, acute chest syndrome, SCD related hospitalizations, or blood transfusions.</p>

SEDATIVE/HYPNOTICS

BENZODIAZEPINE

TEMAZEPAM 7.5mg, 15 mg, 30 mg (compare to Restoril®)
TRIAZOLAM

Estazolam
 Flurazepam
 Halcion® (triazolam)
 Restoril® (temazepam)
 Temazepam 22.5 mg (compare to Restoril®)

Criteria for Approval: The patient has had a documented side effect, allergy, or treatment failure with Temazepam **or Triazolam**. If a product has an AB rated generic, one trial must be the generic.

NON BENZODIAZEPINE, NON BARBITURATE

ESZOPICLONE (compare to Lunesta)
QTY LIMIT: 1 tab/day
 ZALEPLON
QTY LIMIT: 5 mg = 1 cap/day, 10 mg = 2 caps/day
 ZOLPIDEM (compare to Ambien®)
QTY LIMIT: 1 tab/day

Ambien® (zolpidem)
QTY LIMIT: 1 tab/day
 Ambien CR® (zolpidem)
QTY LIMIT: 1 tab/day
 Belsomra® (suvorexant)
QTY LIMIT: 1 tab/day
 Dayvigo® (lemborexant) tablet
QTY LIMIT: 1 tab/day

Ambien, Ambien CR, Lunesta: The patient has had a documented intolerance to the generic equivalent.

Belsomra: The patient has had a documented side effect, allergy, or treatment failure to one preferred sedative/hypnotic.

Dayvigo, Quviviq: The patient has had a documented side effect, allergy, or treatment failure to two preferred sedative/hypnotics and Belsomra.

Edluar, Zolpidem sublingual: The patient has a medical necessity for a

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>ZOLPIDEM CR (compare to Ambien CR®) <i>QTY LIMIT:</i> 1 tab/day</p>	<p>Doxepin 3mg tablets (compare to Silenor) <i>QTY LIMIT:</i> 1 tab/day Edluar® (zolpidem) sublingual tablet <i>QTY LIMIT:</i> 1 tab/day Hetlio^z® (tasimelteon) 20 mg oral capsule <i>QTY LIMIT:</i> 1 capsule/day Maximum days supply per fill is 30 days Lunesta® (eszopiclone) <i>QTY LIMIT:</i> 1 tab/day Quviviq™ (daridorexant) <i>QTY LIMIT:</i> 1 tab/day Ramelteon (compare to Rozerem®) <i>QTY LIMIT:</i> 1 tab/day Rozerem® (ramelteon) <i>QTY LIMIT:</i> 1 tab/day Silenor® (doxepin) <i>QTY LIMIT:</i> 1 tab/day Tasimelteon (compare to Hetlio^z®) <i>QTY LIMIT:</i> 1 tab/day Zolpidem sublingual tablet <i>QTY LIMIT:</i> 1 tab/day</p>	<p>disintegrating tablet formulation (i.e. swallowing disorder). Hetlio^z, Tasimelteon: Patient has documentation of Non-24-Hour Sleep-Wake Disorder (Non24) or Insomnia due to Smith-Magenis Syndrome AND Patient has had a documented side effect, allergy or treatment failure with Ramelteon and at least one OTC melatonin product. For approval of Tasimelteon: Patient must have documented intolerance to Hetlio^z. Ramelteon, Rozerem: The patient has had a documented side effect, allergy, contraindication, or treatment failure to one preferred sedative/hypnotic OR the patient has had a treatment failure after a minimum 2-week trial of melatonin. OR There is a question of substance abuse with the patient or family of the patient. If the request is for Rozerem there must also have been a documented intolerance to Ramelteon. Silenor: The patient has had a documented side effect, allergy, contraindication, or treatment failure to two preferred sedative/hypnotics AND The patient has had a documented intolerance with a preferred generic doxepin formulation.</p>

SMOKING CESSATION THERAPIES

NICOTINE REPLACEMENT: maximum duration is 16 weeks (2 x 8 weeks)/365 days for non-preferred. For approval of therapy beyond the established maximum duration, the prescriber must provide evidence that the patient is engaged in a smoking cessation counseling program.

NICOTINE GUM
 NICOTINE LOZENGE
 NICOTINE PATCH OTC
 NICOTROL® (nicotine) NASAL SPRAY

ORAL THERAPY

BUPROPION SR (compare to Zyban®)
 VARENICLINE (Limited to 18 years and older)
QTY LIMIT: 2 tabs/day
 Max duration 24 weeks (2x12 weeks)/365 days

Smoking Cessation Counseling is encouraged with the use of smoking cessation therapies

The combined prescribing of long acting (patch) and faster acting (gum or lozenge) nicotine replacement therapy is encouraged for greater likelihood of quit success

Vermont QUIT LINE/802 Quits (available free to all patients) 1-800-QUIT-NOW (1- 800-784-8669) <https://802quits.org/>

PREFERRED AGENTS
(No PA required unless otherwise noted)

NON-PREFERRED AGENTS
(PA required)

PA CRITERIA

SUBSTANCE USE DISORDER TREATMENTS

ALCOHOL USE DISORDER

ACAMPROSATE
DISULFIRAM
NALTREXONE
VIVITROL® (naltrexone for extended-release
injectable suspension)
QTY LIMIT: 1 injection (380 mg) per 28 days

OPIOID USE DISORDER

Oral

NALTREXONE tablet
BUPRENORPHINE/NALOXONE TABLET
QTY LIMIT: 8 mg = 3 tablets/day, 2mg N/A
(Maximum Daily Dose = 24 mg/day,
PA required for over 24 mg)

SUBOXONE® sublingual FILM
(buprenorphine/naloxone)
QTY LIMIT: 4mg = 1 film per day, 8 mg = 3 films
per day, 12mg = 2 films per day, 2mg N/A
(Maximum daily Dose = 24 mg/day, PA required
for over 24 mg)

***Maximum days supply for Suboxone Films,
Buprenorphine/naloxone tablets is 30 days***

Injectable

BRIXADI® (buprenorphine extended-release) injection
WEEKLY
QTY LIMIT: 1 syringe per week; maximum days'
supply 28 days (Note: Two 8 mg syringes may
be approved for initial titration purposes in
patients not currently receiving buprenorphine)

BRIXADI® (buprenorphine extended-release) injection
MONTHLY
QTY LIMIT: 1 syringe per 28 days

Buprenorphine sublingual tablet
QTY LIMIT: 2 mg N/A, 8 mg = 3
tablets/day
Maximum Daily Dose = 24 mg/day
Buprenorphine/naloxone (compare to Suboxone®)
sublingual film
QTY LIMIT: 4mg = 1 film per day, 8 mg = 3 films per
day, 12mg = 2 films per day, 2mg N/A
Maximum daily Dose = 24 mg/day
Zubsolv® (buprenorphine/naloxone) sublingual tablet
QTY LIMIT: 1 tablet per day of all strengths

****Maximum days supply for oral
buprenorphine/naloxone films or buprenorphine is 30
days****

CLINICAL CONSIDERATIONS: These products are not FDA approved for
alleviation of pain. For this indication, please refer to the Opioid Analgesics PDL
category. **Note:** As of 1/1/23, a completed Buprenorphine safety checklist (page 2
of the buprenorphine Spoke (OBOT) prior authorization form) must be submitted
with all PA requests.

Buprenorphine/naloxone films, Zubsolv, Buprenorphine tablets: The patient
has experienced a current or past intolerance to the preferred products that
cannot be resolved or mitigated through alternative efforts AND the
Buprenorphine Safety Checklist has been completed (see PA form for detailed
requirements).

Requests to exceed quantity limits or maximum daily dose: Documentation must
be submitted explaining medical necessity for requested dosage regimen AND
the Buprenorphine Safety Checklist has been completed (see PA form for
detailed requirements).

Requests for treatment of pain AND opioid use disorder: The Buprenorphine
Safety Checklist has been completed (see PA form for detailed requirements and
for documentation required) AND other non-opioid medications and pain
management modalities have been trialed prior to increasing the buprenorphine
dose for pain AND split dosing (multiple daily administrations) on current dose
have been trialed for pain control as recommended in the ASAM 2020 practice
guidelines AND clinical rationale has been provided if the request is for a dose
increase > 25% the current daily dose.

Sublocade (to exceed quantity limits): A maintenance dose increase to 300mg
will be considered for those patients who are able to tolerate the 100mg dose but
do not demonstrate a satisfactory clinical response (including supplemental oral

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>SUBLOCADE® (buprenorphine extended-release) injection <i>QTY LIMIT:</i> 300mg 1 injection per 28 days for a maximum of 2 months, then 100mg 1 injection per 28 days thereafter</p> <p>VIVITROL® (naltrexone for extended-release injectable suspension) <i>QTY LIMIT:</i> 1 injection (380 mg) per 28 days</p> <p>Note: Methadone for opioid use disorder can only be prescribed through a Methadone Maintenance Clinic</p>		<p>buprenorphine dosing, documentation of self-reported illicit opioid use, or urine drug screens positive for illicit opioid use). Once the patient is established on a maintenance dose, concurrent use of Sublocade and supplemental oral buprenorphine dosing will not be permitted. Sublocade must be dispensed directly to a healthcare provider and will not be approved for dispensing to the patient.</p>
OPIOID WITHDRAWAL TREATMENT		
<p>Central Alpha Agonists CLONIDINE IR tablets (compare to Catapres®)</p> <p>Note: Methadone for opiate dependency or withdrawal can only be prescribed through a Methadone Maintenance Clinic</p>	<p>Lucemyra® (lofexidine) Maximum length of therapy = 14 days</p>	<p>Lucemyra: Indication for use is the mitigation of opioid withdrawal symptoms to facilitate abrupt opioid discontinuation AND the patient is ≥ 18 years of age AND the patient is unable to tolerate clonidine due to significant side effects.</p>
OVERDOSE TREATMENT		
<p>KLOXXADO™ (naloxone HCl) 8mg Nasal Spray <i>QTY LIMIT:</i> 4 single-use sprays/28days</p> <p>NALOXONE HCl OTC 4 mg Nasal Spray <i>QTY LIMIT:</i> 4 single-use sprays/28days</p> <p>NALOXONE HCl Prefilled luer-lock needleless syringe plus intranasal mucosal atomizing device (Rescue kit)</p> <p>NARCAN® OTC (naloxone hcl) 4mg Nasal Spray <i>QTY LIMIT:</i> 4 single-use sprays/28days</p>	<p>Naloxone HCl RX (compare to Narcan® 4 mg Nasal Spray) <i>QTY LIMIT:</i> 4 single-use sprays/28days</p> <p>Opvee® (nalmefene HCl) Nasal Spray</p> <p>Rextovy® (naloxone HCl) Nasal Spray</p> <p>Zimhi™ (naloxone HCl) 5mg Prefilled Syringe</p>	<p>Naloxone Nasal Spray (RX version): Narcan or OTC Naloxone nasal spray must be on a backorder and unavailable from the manufacturer.</p> <p>Opvee, Rextovy, Zimhi: The prescriber must provide a clinically compelling reason why the preferred agents would not be suitable alternatives.</p>
TESTOSTERONE REPLACEMENT THERAPY		
TOPICAL		
<p>TESTOSTERONE 1.62% Gel Packets <i>QTY LIMIT:</i> 1.25 gm packet (1.62%) = 1 packet/day, 2.5 gm packet (1.62%) = 2 packets/day</p> <p>TESTOSTERONE 1.62% Gel Pump (compare to Androgel®) <i>QTY LIMIT:</i> 2 bottles/30 days</p>	<p>Androgel® pump 1.62% (testosterone pump bottles) <i>QTY LIMIT:</i> 2 bottles/30 days</p> <p>Fortesta® (testosterone 2 % Gel) 60 gm Pump Bottle <i>QTY LIMIT:</i> 2 bottles/30 days</p> <p>Testim® Gel 5 gm (testosterone 1% gel tube) <i>QTY LIMIT:</i> 2 tubes/day</p>	<p>Non-preferred agents: The patient has a documented side effect, allergy, or treatment failure to at least two preferred topical products.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
<p>TESTOSTERONE 1% Gel Packets (compare to Androgel®, Vogelxo®) <i>QTY LIMIT:</i> 2.5 gm packet = 1 packet/day, 5 gm packet = 2 packets/day TESTOSTERONE 2% solution 90ml Pump Bottle</p>	<p>Testosterone 1% gel tube (compare to Testim® Gel 5 gm, Vogelxo®, Androgel®) <i>QTY LIMIT:</i> 2 tubes/day Testosterone 1% Gel Pump (compare to Vogelxo®) <i>QTY LIMIT:</i> 4 bottles/30 days Testosterone 2% gel 60 gm pump bottle (compare to Fortesta®) <i>QTY LIMIT:</i> 2 bottles/30 days Vogelxo® 1% (testosterone 1%) gel, pump <i>QTY LIMIT:</i> 2 tubes/day (5 gm gel tubes), 4 bottles/30 days (gel pump bottle)</p>	
NASAL		
<p>All products require PA</p>	<p>Natesto® (testosterone) nasal gel <i>QTY LIMIT:</i> 3 bottles/30 days</p>	<p>Natesto: The patient has had a documented side effect, allergy, or treatment failure to TWO preferred testosterone products (topical and/or injectable formulations)</p>
ORAL		
<p>All products require PA</p>	<p>Methitest (methyltestosterone) tablet 10 mg Methyltestosterone capsule 10 mg Tlando (testosterone undecanoate) capsule</p> <p>*Maximum day supply all products is 30 days*</p>	<p>Oral non-preferred agents: The patient has had a documented side effect, allergy, or treatment failure to TWO preferred testosterone products (topical and/or injectable formulations) AND if the request is for Methitest or methyltestosterone, the patient has had a documented side effect, allergy, or treatment failure with Tlando.</p>
INJECTABLE		
<p>TESTOSTERONE CYPIONATE IM (compare to Depo®-Testosterone) TESTOSTERONE ENANTHATE IM</p>	<p>Aveed® (testosterone undecanoate) IM Depo®-Testosterone (testosterone cypionate) IM Testopel® (testosterone) implant pellets Xyosted™ (testosterone enanthate) SC</p>	<p>Depo-Testosterone: The patient has a documented intolerance to generic testosterone cypionate. Aveed, Testopel, Xyosted: The patient has had a documented side effect, allergy, or treatment failure to TWO preferred testosterone products, one of which must be an injectable formulation. Treatment failure is defined as inability to achieve testosterone values in the 300-1,000ng/dL range despite adjustments to dose and frequency of injection.</p>

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
---	---------------------------------------	-------------

URINARY ANTISPASMODICS

<p><u>SHORT-ACTING AGENTS</u> BETHANECHOL OXYBUTYNIN TOLTERODINE (compare to Detrol®) TROSPIUM</p> <p><u>LONG-ACTING AGENTS</u> FESOTERODINE ER (compare to Toviaz®) OXYBUTYNIN XL (compare to Ditropan® XL) <i>QTY LIMIT: 1/day</i> SOLIFENACIN (compare to Vesicare®) <i>QTY LIMIT: 1/day</i> TOLTERODINE SR (compare to Detrol LA®)</p> <p><u>TRANSDERMAL/TOPICAL</u> GELNIQUE 10%® (oxybutynin topical gel) <i>QTY LIMIT: 1 sachet/day</i> OXYTROL® (oxybutynin transdermal) <i>QTY LIMIT: 8 patches/28 days</i></p> <p><u>BETA-3 ADRENERGIC AGONISTS</u> MYRBETRIQ® (mirabegron) ER Tablet <i>QTY LIMIT: 1 tablet/day</i></p>	<p>Detrol® (tolterodine) Detrol® LA (tolterodine SR) Flavoxate</p> <p>Darifenacin ER (compare to Enablex®) Ditropan XL® (oxybutynin XL) Toviaz® (fesoterodine ER) <i>QTY LIMIT: 1/day</i> Trospium ER Vesicare® (solifenacin) Vesicare LS™ (solifenacin) oral suspension</p> <p>Gemtesa® (vibegron) tablet <i>QTY LIMIT: 1 tablet/day</i> Myrbetriq® ER Granules for Suspension</p>	<p>Darifenacin ER, Detrol, Detrol LA, Ditropan XL, Flavoxate, Toviaz, trospium ER (generic), Vesicare: The patient has had a documented side effect, allergy, or treatment failure with two preferred long-acting agents. If a medication has an AB rated generic, there must have also been a trial of the generic formulation.</p> <p>Gemtesa: The patient has had a documented side effect, allergy, treatment failure, or contraindication with one preferred urinary antimuscarinic agent and Myrbetriq.</p> <p>Myrbetriq Granules, Vesicare LS: The patient has a diagnosis of neurogenic detrusor overactivity AND the patient has a documented side effect, allergy, or treatment failure with oxybutynin or Toviaz AND for patients ≥ 18 years of age, medical necessity has been provided for a liquid formulation.</p> <p>Limitations: Oxytrol (for Women) OTC not covered</p>
--	--	---

VAGINAL ANTI-INFECTIVES

<p>CLEOCIN® Vaginal Ovules (clindamycin vaginal suppositories) CLEOCIN® (clindamycin vaginal cream 2%) CLOTRIMAZOLE Vaginal cream MICONAZOLE Nitrate Vaginal cream, suppositories MICONAZOLE 1 Vaginal Kit XACIATO™ (clindamycin vaginal gel 2%)</p>	<p>Clindamycin Vaginal (clindamycin vaginal cream 2%) CLINDESSE® (clindamycin vaginal cream 2%) Gynazole-1® (butoconazole vaginal cream 2%) Nuversa™ (metronidazole 1.3% Vaginal Gel) Solosec™ (secnidazole) oral granules packet Terconazole (compare to Terazol®) vaginal cream 0.4%, 0.8%, vaginal suppositories 80 mg Vandazole (metronidazole vaginal 0.75%)</p>	<p>Clindesse, Clindamycin: The patient has had a documented side effect, allergy, or treatment failure to a preferred Cleocin vaginal cream.</p> <p>Nuversa, Vandazole: The patient has had a documented side effect, allergy, or treatment failure to preferred metronidazole vaginal gel.</p> <p>Solosec: The patient has had a documented side effect, allergy, or treatment failure to a preferred topical anti-infective and oral metronidazole.</p> <p>Gynazole, Terconazole: The patient has a documented side effect, allergy, or</p>
---	---	--

PREFERRED AGENTS (No PA required unless otherwise noted)	NON-PREFERRED AGENTS (PA required)	PA CRITERIA
MICONAZOLE 3 Vaginal Kit, cream MICONAZOLE 7 Vaginal cream, suppositories METRONIDAZOLE VAGINAL GEL 0.75%	Xiaciato™ (clindamycin vaginal gel 2%)	treatment failure to a preferred miconazole or clotrimazole formulation.
VASOPRESSIN RECEPTOR ANTAGONIST		
	Jynarque® tablets (tolvaptan) <i>QTY LIMIT:</i> 56 tablets/28 days Samsca® tablets (tolvaptan) <i>QTY LIMIT:</i> 15 mg = 1 tablet/day, 30 mg 2 tablets/day	<p>Jynarque: The patient must be ≥ 18 years of age AND the patient is at risk of rapidly progressing Autosomal Polycystic Kidney Disease (ADPKD) AND the patient has normal serum sodium concentrations before starting the medication (results must be submitted) AND the patient and provider are enrolled in the Jynarque® REMS program</p> <p>Samsca: The agent is being used for the treatment of euvolemic or hypervolemic hyponatremia AND Despite optimal fluid restriction, the patient’s serum sodium < 120 mEq/L or the patient is symptomatic with a serum sodium < 125 mEq/L. AND The treatment will be initiated or is being reinitiated in a hospital setting where serum sodium can be monitored</p>
VITAMINS: PRENATAL MULTIVITAMINS		
C-NATE DHA M-NATAL PLUS NIVA-PLUS PRENATAL PLUS IRON PRENATAL VITAMINS PLUS SE-NATAL CHEW WESTAB PLUS	All others	<p>All Non-Preferred: The prescriber must provide a clinically valid reason for the use of the requested medication including reasons why any of the preferred products would not be a suitable alternative.</p>